

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number  
**WO 03/004467 A2**

- (51) International Patent Classification<sup>7</sup>: **C07D** **Michael, B.** [US/US]; 3766 Tomahawk Lane, San Diego, CA 92117 (US). **YANG, Yi** [US/US]; 8976 Gainsborough Avenue, San Diego, CA 92129 (US).
- (21) International Application Number: **PCT/US02/21280**
- (22) International Filing Date: **5 July 2002 (05.07.2002)** (74) **Agent: HSU, Wendy, Lei;** Agouron Pharmaceuticals, Inc., Legal Division, Patent Dept., 10350 North Torrey Pines Road, La Jolla, CA 92037 (US).
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
60/303,679 6 July 2001 (06.07.2001) **US**  
60/305,274 13 July 2001 (13.07.2001) **US**
- (71) Applicant (for all designated States except US): **AGOURON PHARMACEUTICALS, INC.** [US/US]; 10350 North Torrey Pines Road, La Jolla, CA 92037 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **CHU, Shao, Song** [US/US]; 2075-J Corte del Nogal, Carlsbad, CA 92009 (US). **ALEGRIA, Larry, Andrew** [US/US]; 9632 Galvin Avenue, San Diego, CA 92126 (US). **BLECKMAN, Ted, Michael** [US/US]; 355 Coast Boulevard #22, La Jolla, CA 92037 (US). **CHONG, Wesley, K., M.** [US/US]; 134 Honeycomb Court, Encinitas, CA 92024 (US). **DUVADIE, Rohit, K.** [NP/US]; 11815-B Spruce Run Drive, San Diego, CA 91231 (US). **LI, Lin** [CN/US]; 10632 Nesenda Acuario, San Diego, CA 92131 (US). **REICH, Siegfried, H.** [US/US]; 311 Glenmont Drive, Solana Beach, CA 92075 (US). **ROMINES, William, H.** [US/US]; 12665 Futura Street, San Diego, CA 92130 (US). **WALLACE,**
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.**
- (84) Designated States (regional): **ARIPO** patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **THIAZOLE BENZAMIDE DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS FOR INHIBITING CELL PROLIFERATION, AND METHODS FOR THEIR USE**



(I)

(57) Abstract: Aminothiazole compounds with mono-/di-substituted benzamide are represented by the Formula (I), and their pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, pharmaceutically active metabolites, and pharmaceutically acceptable salts of said metabolites are described. These agents modulate and/or inhibit the cell proliferation and activity of protein kinases and are useful as pharmaceuticals for treating malignancies and other disorders.

**THIAZOLE BENZAMIDE DERIVATIVES AND PHARMACEUTICAL  
COMPOSITIONS FOR INHIBITING CELL PROLIFERATION,  
AND METHODS FOR THEIR USE**

5

TECHNICAL FIELD AND INDUSTRIAL APPLICABILITY  
OF THE INVENTION

This invention relates to compounds with {4-aminothiazol-2-ylamino}-benzamide  
10 nuclei that demonstrate anti-proliferative activity such as antitumor activity, to processes for  
preparing these compounds and to pharmaceutical compositions containing such compounds.  
The invention also relates to the therapeutic or prophylactic use of such compounds and  
compositions, and to methods of treating cancer, viral, microbial, and/or parasitic  
colonization/infection, as well as other disease states associated with unwanted cellular  
15 proliferation, by administering effective amounts of such compounds.

BACKGROUND OF THE INVENTION

Cell proliferation occurs in response to various stimuli and may stem from de-  
regulation of the cell division cycle (or cell cycle), the process by which cells multiply and  
20 divide. Hyperproliferative disease states, including cancer, are characterized by cells  
rampantly winding through the cell cycle with uncontrolled vigor due to, for example, damage  
to the genes that directly or indirectly regulate progression through the cycle. Thus, agents that  
modulate the cell cycle, and thus hyperproliferation, could be used to treat various disease  
states associated with uncontrolled or unwanted cell proliferation. In addition to cancer  
25 chemotherapeutic agents, cell cycle inhibitors are also proposed as antiparasitics (see Gray et  
al., *Curr. Med. Chem.*, 6, 859-875 (1999)) and recently demonstrated as potential antivirals  
(see Schang et al., *J. Virol.*, 74, 2107-2120 (2000)). Moreover, the applicability of  
antiproliferative agents may be expanded to treating cardiovascular maladies such as  
arteriosclerosis or restenosis (see Braun-Dullaeus et al., *Circulation*, 98, 82-89 (1998)), and  
30 states of inflammation, such as arthritis (see Taniguchi et al., *Nature Med.*, 5, 760-767(1999))  
or psoriasis.

Mechanisms of cell proliferation are under active investigation at cellular and  
molecular levels. At the cellular level, deregulation of signaling pathways, loss of cell cycle  
controls, unbridled angiogenesis and stimulation of inflammatory pathways are under scrutiny,  
35 while at the molecular level, these processes are modulated by various proteins, among which

protein kinases are prominent suspects. Overall abatement of proliferation may also result from programmed cell death, or apoptosis, which is also regulated via multiple pathways, some involving proteolytic enzyme proteins.

Among the candidate regulatory proteins, protein kinases are a family of enzymes that catalyze phosphorylation of the hydroxyl group of specific tyrosine, serine or threonine residues in proteins. Typically, such phosphorylation dramatically perturbs the function of the protein, and thus protein kinases are pivotal in the regulation of a wide variety of cellular processes.

Cyclin-dependent kinases (CDKs) are serine-threonine protein kinases that play critical roles in regulating the transitions between different phases of the cell-cycle, such as the progression from a quiescent stage in G<sub>1</sub> (the gap between mitosis and the onset of DNA replication for a new round of cell division) to S (the period of active DNA synthesis), or the progression from G<sub>2</sub> to M phase, in which active mitosis and cell-division occurs. (See, e.g., the articles compiled in *Science*, 274, 1643-1677 (1996); and *Ann. Rev. Cell Dev. Biol.*, 13, 261-291 (1997)). CDK complexes are formed through association of a regulatory cyclin subunit (e.g., cyclin A, B1, B2, D1, D2, D3, and E) and a catalytic kinase subunit (e.g., CDK1, CDK2, CDK4, CDK5, and CDK6). As the name implies, the CDKs display an absolute dependence on the cyclin subunit in order to phosphorylate their target substrates, and different kinase/cyclin pairs function to regulate progression through specific phases of the cell-cycle.

Aberrations in this control system, particularly those that affect the function of CDK4 and CDK2, have been implicated in the advancement of cells to the highly proliferative state characteristic of malignancies, particularly familial melanomas, esophageal carcinomas, and pancreatic cancers (see, e.g., Hall et al., *Adv. Cancer Res.*, 68, 67-108 (1996); Kamb, *Trends in Genetics*, 11, 136-140 (1995); Kamb et al., *Science*, 264, 436-440 (1994)).

Because CDK4 may serve as a general activator of cell division in most cells and complexes of CDK4/cyclin D and CDK2/cyclin E govern the early G<sub>1</sub> phase of the cell cycle, CDK4 or CDK2 inhibitors may be used as anti-proliferative agents. Also, the pivotal roles of cyclin E/CDK2 and cyclin B/CDK1 in the G<sub>1</sub>/S phase and G<sub>2</sub>/M transitions, respectively, offer additional targets for therapeutic intervention in suppressing deregulated cell cycle progression.

A large number of small molecule ATP-site antagonists have been identified as CDK inhibitors (see, Webster, *Exp. Opin. Invest. Drugs*, 7, 865-887 (1998); Stover et al., *Curr. Opin. Drug Disc. Dev.*, 2, 274-285 (1999); Gray et al., *Curr. Med. Chem.*, 6, 859-875 (1999); Sielecki et al., *J. Med. Chem.*, 43, 1-18 (2000); Crews et al., *Curr. Opin. Chem. Biol.*, 4, 47-53 (2000); Buolamwini, *Curr. Pharm. Des.*, 6, 379-392 (2000); and Rosania et al., *Exp. Opin. Ther. Pat.*, 10, 215-230 (2000)).

In addition to the protein kinases identified above, many other protein kinases have been considered to be therapeutic targets, and numerous publications disclose inhibitors of kinase activity, as reviewed in the following: McMahon et al., *Curr. Opin. Drug Disc. Dev.*, 1, 131-146 (1998); Strawn et al., *Exp. Opin. Invest. Drugs*, 7, 553-573 (1998); Adams et al., *Curr. Opin. Drug Disc. Dev.*, 2, 96-109 (1999); Stover et al., *Curr. Opin. Drug Disc. Dev.*, 2, 274-285 (1999); Toledo et al., *Curr. Med. Chem.*, 6, 775-805 (1999); and García-Echeverría et al., *Med. Res. Rev.*, 20, 28-57 (2000).

Among others, the following patent publications disclose thiazole compounds: International Publication No. WO 99/21845 discloses certain 2,4-diaminothiazoles as CDK inhibitors; International Publication No. WO 99/62890 discloses certain isothiazoles as anticancer agents; International Publication No. WO 98/04536 describes certain thiazoles as protein kinase C inhibitors; and European Publication No. EP 816362A(1998) discloses certain thiazoles useful as dopamine D4 receptor antagonists. Certain aminothiazoles are reported in International Publication No. WO 99/65844 and International Publication No. WO 99/24416, and certain aminobenzothiazoles are disclosed in International Publication No. WO 99/24035. International Publication No. WO 00/17175 describes certain other aminothiazoles as p38 mitogen-activated protein (MAP) kinase inhibitors, and International Publication No. WO 00/26202, International Publication No. WO 00/26203, and US Patent No. 6,114,365 describe certain aminothiazoles and certain ureidothiazoles as anti-tumor agents. International Publication No. WO 99/21845 discloses certain 4-aminothiazole derivatives containing unsubstituted nitrogen or primary benzamides.

There is still a need, however, for more potent inhibitors of protein kinases. Moreover, as is understood by those skilled in the art, it is desirable for kinase inhibitors to possess both high affinity for the target kinase as well as high selectivity versus other protein kinases.

#### SUMMARY OF THE INVENTION


An object of the invention is to discover potent anti-proliferative agents. Another object of the invention is to discover effective inhibitors of protein kinases.

These and other objects of the invention, which will become apparent from the following description, have been achieved through the discovery of 4-aminothiazole compounds with mono- or di-N-substituted benzamides. The invention also relates to pharmaceutically acceptable prodrugs, pharmaceutically active metabolites, and pharmaceutically acceptable salts of such compounds (such compounds, prodrugs, metabolites and salts are collectively referred to as "agents") which modulate and/or inhibit cell growth.

Thus, the inventive agents and pharmaceutical compositions containing such agents are useful in treating various diseases or disorder states associated with uncontrolled or unwanted cellular proliferation, such as cancer, autoimmune diseases, viral diseases, fungal diseases, neurodegenerative disorders, and cardiovascular diseases.

Further, the agents modulate and/or inhibit the activity of protein kinases, for example one or more CDKs, such as CDK1, CDK2, CDK4 and/or CDK6, or cyclin complexes thereof, and/or one or more LCKs, VEGF or FGFs. Thus, the pharmaceutical compositions containing such agents are useful in treating diseases mediated by kinase activity, such as cancer.

In a general aspect, the invention relates to compounds represented by Formula (I):



wherein:

$R^1$  and  $R^2$  are each independently hydrogen, or an alkyl, alkenyl, alkynyl, heteroalkyl, alkoxy, amino alkyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl group unsubstituted or substituted with one or more substituents independently selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N}-\text{OH}$ ,  $\text{N}-\text{OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{OR}_b$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{OR}_c$ , keto ( $=\text{O}$ ), thioketo ( $=\text{S}$ ),  $-\text{SO}_2-\text{R}_c$ ,  $-\text{SO}-\text{R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{SO}_2-\text{R}_c$ ,  $-\text{O}-\text{SO}-\text{R}_c$ ,  $-\text{O}-\text{S}-\text{R}_c$ ,  $-\text{S}-\text{CO}-\text{R}_c$ ,  $-\text{SO}-\text{CO}-\text{OR}_c$ ,  $-\text{SO}_2-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{SO}_3$ ,  $-\text{NR}_c-\text{SR}_d$ ,  $-\text{NR}_c-\text{SO}-\text{R}_d$ ,  $\text{NR}_c-\text{SO}_2-\text{R}_d$ ,  $-\text{CO}-\text{SR}_c$ ,  $-\text{CO}-\text{SO}-\text{R}_c$ ,  $-\text{CO}-\text{SO}_2-\text{R}_c$ ,  $-\text{CS}-\text{R}_c$ ,  $-\text{CSO}-\text{R}_c$ ,  $-\text{CSO}_2-\text{R}_c$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{O}-\text{CS}-\text{R}_c$ ,  $-\text{O}-\text{CSO}-\text{R}_c$ ,  $-\text{O}-\text{CSO}_2-\text{R}_c$ ,  $-\text{SO}_2-\text{NR}_d\text{R}_e$ ,  $-\text{SO}-\text{NR}_d\text{R}_e$ ,  $-\text{S}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_d-\text{CSO}_2-\text{R}_d$ ,  $-\text{NR}_c-\text{CSO}-\text{R}_d$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{SH}$ ,  $-\text{S}-\text{R}_b$ , and  $-\text{PO}_2-\text{OR}_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_f$ ,  $-\text{O}-\text{CO}-\text{R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl

group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ , - $\text{NH}_2$ , - $\text{CN}$ ,  $-(\text{CH}_2)_z\text{-CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl, - $\text{OH}$ , keto ( $=\text{O}$ ), - $\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $\text{-NR}_d\text{R}_e$ ,  $\text{-CO-NR}_d\text{R}_e$ ,  $\text{-CO-OR}_c$ ,  $\text{-CO-R}_c$ ,  $\text{-NR}_c\text{-CO-NR}_d\text{R}_e$ ,  $\text{-C-CO-OR}_c$ ,  $\text{-NR}_c\text{-CO-R}_d$ ,  $\text{-O-CO-O-R}_c$ ,  $\text{-O-CO-NR}_d\text{R}_e$ , - $\text{SH}$ , - $\text{O-R}_b$ ,  $\text{-O-R}_a\text{-O-}$ , - $\text{S-R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are defined above, where  $R^1$  and  $R^2$  are not both hydrogen, or

$R^1$  or  $R^2$ , together with the  $\text{N}-\overset{\text{O}}{\underset{\parallel}{\text{C}}}$  and two adjacent carbon atoms of the phenyl ring of Formula (I), forms a 5- or 6-membered ring structure fused to the phenyl ring of Formula (I) and unsubstituted or substituted with one or more substituents independently selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $\text{-NO}_2$ ,  $\text{-NH}_2$ ,  $\text{-N-OH}$ ,  $\text{-N-OR}_c$ , - $\text{CN}$ ,  $-(\text{CH}_2)_z\text{-CN}$  where  $z$  is 0-4, halogen, - $\text{OH}$ ,  $\text{-O-R}_a\text{-O-}$ ,  $\text{-OR}_b$ ,  $\text{-CO-R}_c$ ,  $\text{-O-CO-R}_c$ ,  $\text{-CO-OR}_c$ ,  $\text{-O-CO-OR}_c$ ,  $\text{-O-CO-O-CO-R}_c$ ,  $\text{-O-OR}_c$ , keto ( $=\text{O}$ ), thioketo ( $=\text{S}$ ),  $\text{SO}_2\text{-R}_c$ ,  $\text{-SO-R}_c$ ,  $\text{-NR}_d\text{R}_e$ ,  $\text{-CO-NR}_d\text{R}_e$ ,  $\text{-O-CO-NR}_d\text{R}_e$ ,  $\text{-NR}_c\text{-CO-NR}_d\text{R}_e$ ,  $\text{-NR}_c\text{-CO-R}_e$ ,  $\text{-NR}_c\text{-CO-OR}_e$ ,  $\text{-CO-NR}_c\text{-CO-R}_d$ ,  $\text{-O-SO}_2\text{-R}_c$ ,  $\text{-O-SO-R}_c$ ,  $\text{-O-S-R}_c$ ,  $\text{-S-CO-R}_c$ ,  $\text{-SO-CO-OR}_c$ ,  $\text{-SO}_2\text{-CO-OR}_c$ ,  $\text{-O-SO}_3$ ,  $\text{-NR}_c\text{-SR}_d$ ,  $\text{-NR}_c\text{-SO-R}_d$ ,  $\text{NR}_c\text{-SO}_2\text{-R}_d$ ,  $\text{-CO-SR}_c$ ,  $\text{-CO-SO-R}_c$ ,  $\text{-CO-SO}_2\text{-R}_c$ ,  $\text{-CS-R}_c$ ,  $\text{-CSO-R}_c$ ,  $\text{-CSO}_2\text{-R}_c$ ,  $\text{-NR}_c\text{-CS-R}_d$ ,  $\text{-O-CS-R}_c$ ,  $\text{-O-CSO-R}_c$ ,  $\text{-O-CSO}_2\text{-R}_c$ ,  $\text{-SO}_2\text{-NR}_d\text{R}_e$ ,  $\text{-SO-NR}_d\text{R}_e$ ,  $\text{-S-NR}_d\text{R}_e$ ,  $\text{-NR}_d\text{-CSO}_2\text{-R}_d$ ,  $\text{-NR}_c\text{-CSO-R}_d$ ,  $\text{-NR}_c\text{-CS-R}_d$ , - $\text{SH}$ , - $\text{S-R}_b$ , and  $\text{-PO}_2\text{-OR}_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $\text{-CO-R}_c$ ,  $\text{-CO-OR}_c$ ,  $\text{-O-CO-O-R}_c$ ,  $\text{-O-CO-R}_c$ ,  $\text{-NR}_c\text{-CO-R}_d$ ,  $\text{-CO-NR}_d\text{R}_e$ , - $\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl,  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $\text{-COR}_f$ ,  $\text{-COOR}_f$ ,  $\text{-O-CO-O-R}_f$ ,  $\text{-O-CO-R}_f$ , - $\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group

consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, keto (=O), -N-OH, N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are defined above; or

R<sup>1</sup> and R<sup>2</sup>, taken together with the nitrogen atom to which they are bonded, form a monocyclic or fused or non-fused polycyclic structure which may contain one to three additional heteroatoms, unsubstituted or substituted with one or more substituents independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, O-R<sub>a</sub>-O-, -OR<sub>b</sub>, -CO-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-OR<sub>c</sub>, -O-CO-O-CO-R<sub>c</sub>, -O-OR<sub>c</sub>, keto (=O), thioketo (=S), -SO<sub>2</sub>-R<sub>c</sub>, -SO-R<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-R<sub>e</sub>, -NR<sub>c</sub>-CO-OR<sub>e</sub>, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>, -O-SO<sub>2</sub>-R<sub>c</sub>, -O-SO-R<sub>c</sub>, -O-S-R<sub>c</sub>, -S-CO-R<sub>c</sub>, -SO-CO-OR<sub>c</sub>, -SO<sub>2</sub>-CO-OR<sub>c</sub>, -O-SO<sub>3</sub>-, -NR<sub>c</sub>-SR<sub>d</sub>, -NR<sub>c</sub>-SO-R<sub>d</sub>, -NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>, -CO-SR<sub>c</sub>, -CO-SO-R<sub>c</sub>, -CO-SO<sub>2</sub>-R<sub>c</sub>, -CS-R<sub>c</sub>, -CSO-R<sub>c</sub>, -CSO<sub>2</sub>-R<sub>c</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -O-CS-R<sub>c</sub>, -O-CSO-R<sub>c</sub>, -O-CSO<sub>2</sub>-R<sub>c</sub>, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>c</sub>-CSO-R<sub>d</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, -COR<sub>f</sub>, -COOR<sub>f</sub>, -O-CO-O-R<sub>f</sub>, -O-CO-R<sub>f</sub>, -OH, Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where R<sub>d</sub> and R<sub>e</sub> can cyclize to form a heteroaryl or heterocycloalkyl group, and R<sub>f</sub> is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, keto (=O), -N-OH, -N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl,

unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are defined above;

$R^3$  is an aryl, heteroaryl, alkyl, or cycloalkyl group, unsubstituted or substituted with one or more substituents independently selected from the group consisting of

5 hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N}-\text{OH}$ ,  $-\text{N}-\text{OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{OR}_b$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{OR}_c$ , keto ( $=\text{O}$ ), thioketo ( $=\text{S}$ ),  $-\text{SO}_2-\text{R}_c$ ,  $-\text{SO}-\text{R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{OR}_e$ ,  $-\text{CO}-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{SO}_2-\text{R}_c$ ,  $-\text{O}-\text{SO}-\text{R}_c$ ,  $-\text{O}-\text{S}-\text{R}_c$ ,  $-\text{S}-\text{CO}-\text{R}_c$ ,  $-\text{SO}-\text{CO}-\text{OR}_c$ ,  $-\text{SO}_2-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{SO}_3$ ,  $-\text{NR}_c-\text{SR}_d$ ,  $-\text{NR}_c-\text{SO}-\text{R}_d$ ,  $-\text{NR}_c-\text{SO}_2-\text{R}_d$ ,  $-\text{CO}-\text{SR}_c$ ,  $-\text{CO}-\text{SO}-\text{R}_c$ ,  $-\text{CO}-\text{SO}_2-\text{R}_c$ ,  $-\text{CS}-\text{R}_c$ ,  $-\text{CSO}-\text{R}_c$ ,  $-\text{CSO}_2-\text{R}_c$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{O}-\text{CS}-\text{R}_c$ ,  $-\text{O}-\text{CSO}-\text{R}_c$ ,  $-\text{O}-\text{CSO}_2-\text{R}_c$ ,  $-\text{SO}_2-\text{NR}_d\text{R}_e$ ,  $-\text{SO}-\text{NR}_d\text{R}_e$ ,  $-\text{S}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_d-\text{CSO}_2-\text{R}_d$ ,  $-\text{NR}_c-\text{CSO}-\text{R}_d$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{SH}$ ,  $-\text{S}-\text{R}_b$ , and  $-\text{PO}_2-\text{OR}_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_f$ ,  $-\text{O}-\text{CO}-\text{R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with


25 one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ , keto ( $=\text{O}$ ),  $-\text{N}-\text{OH}$ ,  $-\text{N}-\text{OR}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{C}-\text{CO}-\text{OR}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{SH}$ ,  $-\text{O}-\text{R}_b$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{S}-\text{R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are defined above; and

Y is hydrogen, alkyl, heteroalkyl, haloalkyl, halocycloalkyl, haloheterocycloalkyl, cycloalkyl, heterocycloalkyl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N}-\text{OH}$ ,  $-\text{N}-\text{OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{OR}_b$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{OR}_c$ , keto ( $=\text{O}$ ), thioketo ( $=\text{S}$ ),  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{O}-\text{CO}-$




$\text{NR}_d\text{R}_e$ ,  $-\text{NR}_e-\text{CO}-\text{R}_e$ ,  $-\text{NR}_e-\text{CO}_2-\text{OR}_e$ ,  $-\text{CO}-\text{NR}_e-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{SO}_2-\text{R}_e$ ,  $-\text{O}-\text{SO}-\text{R}_e$ ,  $-\text{O}-\text{S}-\text{R}_e$ ,  $-\text{S}-\text{CO}-\text{R}_e$ ,  $-\text{SO}-\text{CO}-\text{OR}_e$ ,  $-\text{SO}_2-\text{CO}-\text{OR}_e$ ,  $-\text{O}-\text{SO}_3$ ,  $-\text{NR}_e-\text{SR}_d$ ,  $-\text{NR}_e-\text{SO}-\text{R}_d$ ,  $\text{NR}_e-\text{SO}_2-\text{R}_d$ ,  $-\text{CO}-\text{SR}_e$ ,  $-\text{CO}-\text{SO}-\text{R}_e$ ,  $-\text{CO}-\text{SO}_2-\text{R}_e$ ,  $-\text{CS}-\text{R}_e$ ,  $-\text{CSO}-\text{R}_e$ ,  $-\text{CSO}_2-\text{R}_e$ ,  $-\text{NR}_e-\text{CS}-\text{R}_d$ ,  $-\text{O}-\text{CS}-\text{R}_e$ ,  $-\text{O}-\text{CSO}-\text{R}_e$ ,  $-\text{O}-\text{CSO}_2-\text{R}_e$ ,  $-\text{SO}_2-\text{NR}_d\text{R}_e$ ,  $-\text{SO}-\text{NR}_d\text{R}_e$ ,  $-\text{S}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_d-\text{CSO}_2-\text{R}_d$ ,  $-\text{NR}_e-\text{CSO}-\text{R}_d$ ,  $-\text{NR}_e-\text{CS}-\text{R}_d$ ,  $-\text{SH}$ ,  $-\text{S}-\text{R}_b$ , and  $-\text{PO}_2-\text{OR}_e$ , where  $\text{R}_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $\text{R}_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-\text{R}_e$ ,  $-\text{CO}-\text{OR}_e$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_e$ ,  $-\text{O}-\text{CO}-\text{R}_e$ ,  $-\text{NR}_e-\text{CO}-\text{R}_d$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{OH}$ , heterocycloalkyl, and cycloalkyl, and  $\text{R}_e$ ,  $\text{R}_d$  and  $\text{R}_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_f$ ,  $-\text{O}-\text{CO}-\text{R}_f$ ,  $-\text{OH}$ , cycloalkyl, and heterocycloalkyl, where  $\text{R}_d$  and  $\text{R}_e$  can cyclize to form a heteroaryl or heterocycloalkyl group and  $\text{R}_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl.

The nitrogen-containing ring optionally formed by  $\text{R}^1$  and  $\text{R}^2$  may be monocyclic, or fused or un-fused polycyclic (i.e. spiral), and may contain one to three additional heteroatoms selected from N, O or S. Examples of such a ring include




According to one preferred embodiment of the invention,  $(\text{C}(\text{O})\text{N}(\text{R}^1)(\text{R}^2))$  moiety is connected meta or para to the secondary amine linking the phenyl and thiazole rings, and Y can be at any position on the phenyl ring. More preferably, the  $(\text{C}(\text{O})\text{N}(\text{R}^1)(\text{R}^2))$  moiety is para and Y is meta to the secondary amine linking the phenyl and thiazole rings.



In one preferred embodiment of the invention, in the compounds of Formula (I),  $\text{R}^3$  is an aryl or heteroaryl group having one or more substituents selected from the group consisting of a halogen, alkoxy,  $-\text{OH}$ , alkyl and  $-\text{NO}_2$  groups.

The invention is also directed to pharmaceutically acceptable salts of compounds represented by the Formula (I), pharmaceutically acceptable prodrugs, pharmaceutically active metabolites of compounds represented by the Formula (I), and pharmaceutically acceptable salts of such metabolites. Advantageous methods of making the compounds of the Formula (I) are also described.

In a preferred general embodiment, the invention relates to compounds represented by Formula (II):



wherein:

$R^1$  is selected from the group consisting of: alkyl, alkenyl, alkynyl, heteroalkyl, halogen, aryl, heteroaryl, cycloalkyl, and heterocycloalkyl groups unsubstituted or substituted with one or more substituents independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z\text{-CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O-R}_a\text{-O-}$ ,  $-\text{OR}_b$ ,  $-\text{CO-R}_c$ ,  $-\text{O-CO-R}_c$ ,  $-\text{CO-OR}_c$ ,  $-\text{O-CO-OR}_c$ ,  $-\text{O-CO-O-CO-R}_c$ ,  $-\text{O-OR}_c$ , keto ( $=\text{O}$ ), thioketo ( $=\text{S}$ ),  $-\text{SO}_2\text{-R}_c$ ,  $-\text{SO-R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{O-CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c\text{-CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c\text{-CO-R}_e$ ,  $-\text{NR}_c\text{-CO}_2\text{-OR}_c$ ,  $-\text{CO-NR}_c\text{-CO-R}_d$ ,  $-\text{O-SO}_2\text{-R}_c$ ,  $-\text{O-SO-R}_c$ ,  $-\text{O-S-R}_c$ ,  $-\text{S-CO-R}_c$ ,  $-\text{SO-CO-OR}_c$ ,  $-\text{SO}_2\text{-CO-OR}_c$ ,  $-\text{O-SO}_3$ ,  $-\text{NR}_c\text{-SR}_d$ ,  $-\text{NR}_c\text{-SO-R}_d$ ,  $\text{NR}_c\text{-SO}_2\text{-R}_d$ ,  $-\text{CO-SR}_c$ ,  $-\text{CO-SO-R}_c$ ,  $-\text{CO-SO}_2\text{-R}_c$ ,  $-\text{CS-R}_c$ ,  $-\text{CSO-R}_c$ ,  $-\text{CSO}_2\text{-R}_c$ ,  $-\text{NR}_c\text{-CS-R}_d$ ,  $-\text{O-CS-R}_c$ ,  $-\text{O-CSO-R}_c$ ,  $-\text{O-CSO}_2\text{-R}_c$ ,  $-\text{SO}_2\text{-NR}_d\text{R}_e$ ,  $-\text{SO-NR}_d\text{R}_e$ ,  $-\text{S-NR}_d\text{R}_e$ ,  $-\text{NR}_d\text{-CSO}_2\text{-R}_d$ ,  $-\text{NR}_c\text{-CSO-R}_d$ ,  $-\text{NR}_c\text{-CS-R}_d$ ,  $-\text{SH}$ ,  $-\text{S-R}_b$ , and  $-\text{PO}_2\text{-OR}_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO-R}_c$ ,  $-\text{CO-OR}_c$ ,  $-\text{O-CO-O-R}_c$ ,  $-\text{O-CO-R}_c$ ,  $-\text{NR}_c\text{-CO-R}_d$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O-CO-O-R}_f$ ,  $-\text{O-CO-R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl,

heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, keto (=O), -N-OH, N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are defined above; and

R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are each independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, -O-R<sub>a</sub>-O-, -OR<sub>b</sub>, -CO-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-OR<sub>c</sub>, -O-CO-O-CO-R<sub>c</sub>, -O-OR<sub>c</sub>, keto (=O), thioketo (=S), -SO<sub>2</sub>-R<sub>c</sub>, -SO-R<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-R<sub>e</sub>, -NR<sub>c</sub>-CO<sub>2</sub>-OR<sub>e</sub>, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>, -O-SO<sub>2</sub>-R<sub>c</sub>, -O-SO-R<sub>e</sub>, -O-S-R<sub>c</sub>, -S-CO-R<sub>c</sub>, -SO-CO-OR<sub>c</sub>, -SO<sub>2</sub>-CO-OR<sub>c</sub>, -O-SO<sub>3</sub>, -NR<sub>c</sub>-SR<sub>d</sub>, -NR<sub>c</sub>-SO-R<sub>d</sub>, NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>, -CO-SR<sub>c</sub>, -CO-SO-R<sub>c</sub>, -CO-SO<sub>2</sub>-R<sub>c</sub>, -CS-R<sub>c</sub>, -CSO-R<sub>c</sub>, -CSO<sub>2</sub>-R<sub>c</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -O-CS-R<sub>c</sub>, -O-CSO-R<sub>c</sub>, -O-CSO<sub>2</sub>-R<sub>c</sub>, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>c</sub>-CSO-R<sub>d</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, -COR<sub>f</sub>, -COOR<sub>f</sub>, -O-CO-O-R<sub>f</sub>, -O-CO-R<sub>f</sub>, -OH, Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where R<sub>d</sub> and R<sub>e</sub> can cyclize to form a heteroaryl or heterocycloalkyl group, and R<sub>f</sub> is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, keto (=O), -N-OH, N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are defined above.

Preferred compounds of the invention include those represented by Formula (II) wherein R<sup>1</sup> is an alkyl, heteroalkyl or heterocycloalkyl group substituted with one or more substituents independently selected from the group consisting of hydrogen, alkyl, heteroalkyl,

haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, -OR<sub>a</sub>-O-, -OR<sub>b</sub>-, -CO-R<sub>c</sub>-, -O-CO-R<sub>c</sub>-, -CO-OR<sub>c</sub>-, -O-CO-OR<sub>c</sub>-, -O-CO-O-CO-R<sub>c</sub>-, -O-OR<sub>c</sub>-, keto (=O), thioketo (=S), -SO<sub>2</sub>-R<sub>c</sub>-, -SO-R<sub>c</sub>-, -NR<sub>d</sub>R<sub>e</sub>-, -CO-NR<sub>d</sub>R<sub>e</sub>-, -O-CO-NR<sub>d</sub>R<sub>e</sub>-, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>-, -NR<sub>c</sub>-CO-R<sub>e</sub>-, -NR<sub>c</sub>-CO<sub>2</sub>-OR<sub>c</sub>-, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>-, -O-SO<sub>2</sub>-R<sub>c</sub>-, -O-SO-R<sub>c</sub>-, -O-S-R<sub>c</sub>-, -S-CO-R<sub>c</sub>-, -SO-CO-OR<sub>c</sub>-, -SO<sub>2</sub>-CO-OR<sub>c</sub>-, -O-SO<sub>3</sub>-, -NR<sub>c</sub>-SR<sub>d</sub>-, -NR<sub>c</sub>-SO-R<sub>d</sub>-, NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>-, -CO-SR<sub>c</sub>-, -CO-SO-R<sub>c</sub>-, -CO-SO<sub>2</sub>-R<sub>c</sub>-, -CS-R<sub>c</sub>-, -CSO-R<sub>c</sub>-, -CSO<sub>2</sub>-R<sub>c</sub>-, -NR<sub>c</sub>-CS-R<sub>d</sub>-, -O-CS-R<sub>c</sub>-, -O-CSO-R<sub>c</sub>-, -O-CSO<sub>2</sub>-R<sub>c</sub>-, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>-, -SO-NR<sub>d</sub>R<sub>e</sub>-, -S-NR<sub>d</sub>R<sub>e</sub>-, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>-, -NR<sub>c</sub>-CSO-R<sub>d</sub>-, -NR<sub>c</sub>-CS-R<sub>d</sub>-, -SH, -S-R<sub>b</sub>-, and -PO<sub>2</sub>-OR<sub>c</sub>-, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>-, -CO-OR<sub>c</sub>-, -O-CO-O-R<sub>c</sub>-, -O-CO-R<sub>c</sub>-, -NR<sub>c</sub>-CO-R<sub>d</sub>-, -CO-NR<sub>d</sub>R<sub>e</sub>-, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, -COR<sub>f</sub>-, -COOR<sub>f</sub>-, -O-CO-O-R<sub>f</sub>-, -O-CO-R<sub>f</sub>-, -OH, Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where R<sub>d</sub> and R<sub>e</sub> can cyclize to form a heteroaryl or heterocycloalkyl group, and R<sub>f</sub> is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, keto (=O), -N-OH, N-OR<sub>c</sub>-, -NR<sub>d</sub>R<sub>e</sub>-, -CO-NR<sub>d</sub>R<sub>e</sub>-, -CO-OR<sub>c</sub>-, -CO-R<sub>c</sub>-, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>-, -C-CO-OR<sub>c</sub>-, -NR<sub>c</sub>-CO-R<sub>d</sub>-, -O-CO-O-R<sub>c</sub>-, -O-CO-NR<sub>d</sub>R<sub>e</sub>-, -SH, -O-R<sub>b</sub>-, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>-, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are defined above. Preferably, the substituted alkyl for R<sup>1</sup> is -N(R<sup>7</sup>)(R<sup>8</sup>) where R<sup>7</sup> and R<sup>8</sup> are each independently an alkyl, alkyl- (heterocycloalkyl group unsubstituted or substituted with one or more substituents independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>-, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, -O-R<sub>a</sub>-O-, -OR<sub>b</sub>-, -CO-R<sub>c</sub>-, -O-CO-R<sub>c</sub>-, -CO-OR<sub>c</sub>-, -O-CO-OR<sub>c</sub>-, -O-CO-O-CO-R<sub>c</sub>-, -O-OR<sub>c</sub>-, keto (=O), thioketo (=S), -SO<sub>2</sub>-R<sub>c</sub>-, -SO-R<sub>c</sub>-, -NR<sub>d</sub>R<sub>e</sub>-, -CO-NR<sub>d</sub>R<sub>e</sub>-, -O-CO-NR<sub>d</sub>R<sub>e</sub>-, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>-, -NR<sub>c</sub>-CO-R<sub>e</sub>-, -NR<sub>c</sub>-CO-OR<sub>c</sub>-, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>-, -O-SO<sub>2</sub>-R<sub>c</sub>-, -O-SO-R<sub>c</sub>-, -O-S-R<sub>c</sub>-, -S-CO-R<sub>c</sub>-, -SO-CO-OR<sub>c</sub>-, -SO<sub>2</sub>-CO-OR<sub>c</sub>-, -O-SO<sub>3</sub>-, -NR<sub>c</sub>-SR<sub>d</sub>-, -NR<sub>c</sub>-SO-R<sub>d</sub>-, NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>-, -CO-SR<sub>c</sub>-, -CO-SO-R<sub>c</sub>-, -CO-SO<sub>2</sub>-R<sub>c</sub>-, -CS-R<sub>c</sub>-, -CSO-R<sub>c</sub>-, -CSO<sub>2</sub>-R<sub>c</sub>-, -NR<sub>c</sub>-CS-R<sub>d</sub>-, -O-CS-R<sub>c</sub>-, -O-CSO-R<sub>c</sub>-, -O-CSO<sub>2</sub>-R<sub>c</sub>-, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>-, -SO-NR<sub>d</sub>R<sub>e</sub>-, -S-NR<sub>d</sub>R<sub>e</sub>-, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>-, -NR<sub>c</sub>-CSO-R<sub>d</sub>-, -NR<sub>c</sub>-CS-R<sub>d</sub>-, -SH, -S-R<sub>b</sub>-, and -PO<sub>2</sub>-OR<sub>c</sub>-, where R<sub>a</sub> is selected from the group consisting of alkyl,

heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-R_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-R_c$ ,  $-\text{O}-\text{CO}-R_c$ ,  $-\text{NR}_c-\text{CO}-R_d$ ,  $-\text{CO}-\text{NR}_dR_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl,

5 haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-R_f$ ,  $-\text{O}-\text{CO}-R_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further

10 substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ , keto ( $=\text{O}$ ),  $-\text{N}-\text{OH}$ ,  $\text{N}-\text{OR}_c$ ,  $-\text{NR}_dR_e$ ,  $-\text{CO}-\text{NR}_dR_e$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-R_c$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_dR_e$ ,  $-\text{C}-\text{CO}-\text{OR}_c$ ,  $-\text{NR}_c-\text{CO}-R_d$ ,  $-\text{O}-\text{CO}-\text{O}-R_c$ ,  $\text{O}-\text{CO}-\text{NR}_dR_e$ ,  $-\text{SH}$ ,  $-\text{O}-R_b$ ,  $-\text{O}-R_a-\text{O}-$ ,  $-\text{S}-R_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted

15 heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are defined above, or alkyl-(substituted aryl) group unsubstituted or substituted with one or more substituents independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N}-\text{OH}$ ,  $\text{N}-\text{OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O}-R_a-\text{O}-$ ,  $-\text{OR}_b$ ,  $-\text{CO}-R_c$ ,  $-\text{O}-\text{CO}-R_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{CO}-R_c$ ,  $-\text{O}-\text{OR}_c$ , keto ( $=\text{O}$ ), thioketo ( $=\text{S}$ ),  $-\text{SO}_2-R_c$ ,  $-\text{SO}-R_c$ ,  $-\text{NR}_dR_e$ ,  $-\text{CO}-\text{NR}_dR_e$ ,  $-\text{O}-\text{CO}-\text{NR}_dR_e$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_dR_e$ ,  $-\text{NR}_c-\text{CO}-R_c$ ,  $-\text{NR}_c-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{NR}_c-\text{CO}-R_d$ ,  $-\text{O}-\text{SO}_2-R_c$ ,  $-\text{O}-\text{SO}-R_c$ ,  $-\text{O}-\text{S}-R_c$ ,  $-\text{S}-\text{CO}-R_c$ ,  $-\text{SO}-\text{CO}-\text{OR}_c$ ,  $-\text{SO}_2-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{SO}_3$ ,  $-\text{NR}_c-\text{SR}_d$ ,  $-\text{NR}_c-\text{SO}-R_d$ ,  $\text{NR}_c-\text{SO}_2-R_d$ ,  $-\text{CO}-\text{SR}_c$ ,  $-\text{CO}-\text{SO}-R_c$ ,  $-\text{CO}-\text{SO}_2-R_c$ ,  $-\text{CS}-R_c$ ,  $-\text{CSO}-R_c$ ,  $-\text{CSO}_2-R_c$ ,  $-\text{NR}_c-\text{CS}-R_d$ ,  $-\text{O}-\text{CS}-R_c$ ,  $-\text{O}-\text{CSO}-R_c$ ,  $-\text{O}-\text{CSO}_2-R_c$ ,  $-\text{SO}_2-\text{NR}_dR_e$ ,  $-\text{SO}-\text{NR}_dR_e$ ,  $-\text{S}-$

25  $\text{NR}_dR_e$ ,  $-\text{NR}_d-\text{CSO}_2-R_d$ ,  $-\text{NR}_c-\text{CSO}-R_d$ ,  $-\text{NR}_c-\text{CS}-R_d$ ,  $-\text{SH}$ ,  $-\text{S}-R_b$ , and  $-\text{PO}_2-\text{OR}_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-R_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-R_c$ ,  $-\text{O}-\text{CO}-R_c$ ,  $-\text{NR}_c-\text{CO}-R_d$ ,  $-\text{CO}-\text{NR}_dR_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the

30 group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-R_f$ ,  $-\text{O}-\text{CO}-R_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the

35 above substituents may be further substituted with one or more substituents independently

selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, keto (=O), -N-OH, N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are defined above; and

R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are each independently selected from the group consisting of hydrogen, halogen, -OH, C<sub>1</sub>-C<sub>3</sub>-alkoxy, C<sub>1</sub>-C<sub>3</sub>-alkyl, heteroalkyl, or -N-COR<sub>c</sub>, -SR<sub>c</sub>, -SO-R<sub>c</sub>, -SO<sub>2</sub>R<sub>c</sub>, and -CO-R<sub>c</sub>, where R<sub>c</sub> is as defined above.

The invention also relates to methods of treating proliferative diseases such as cancer, autoimmune diseases, viral diseases, fungal diseases, neurodegenerative disorders and cardiovascular diseases, comprising administering effective amounts of a compound of Formula (I) or a pharmaceutically acceptable salt, pharmaceutically acceptable prodrug, pharmaceutically active metabolite, or pharmaceutically acceptable salt of such compound or metabolite to a subject in need of such treatment.

The invention further relates to a method of modulating and/or inhibiting the kinase activity of one or more CDKs such as CDK1, CDK2, CDK4, and/or CDK6 or cyclin complexes thereof, VEGF, FGF and/or LCK by administering a compound of Formula (I) or a pharmaceutically acceptable salt, pharmaceutically acceptable prodrug, or pharmaceutically acceptable salt of such compound or metabolite thereof.

The invention also relates to pharmaceutical compositions, each comprising an effective amount of an agent selected from compounds of Formula (I) and pharmaceutically active metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable salts of such compounds and metabolites, and a pharmaceutically acceptable carrier or vehicle for such agent.

#### DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

The inventive compounds of Formula (I) are potent anti-proliferative agents. The compounds are also useful for mediating the activity of protein kinases. More particularly, the compounds are useful as agents for modulating and/or inhibiting the activity of various enzymes, for example protein kinases, thus providing treatments for cancer or other diseases associated with uncontrolled or abnormal cellular proliferation.

The diseases or disorders in association with uncontrolled or abnormal cellular proliferation include the following:

- a variety of cancers, including carcinoma, hematopoietic tumors of lymphoid lineage, hematopoietic tumors of myeloid lineage, tumors of mesenchymal origin, tumors of the central and peripheral nervous system and other tumors including melanoma, seminoma and Kaposi's sarcoma and the like.
- 5 - a disease process which features abnormal cellular proliferation, e.g., benign prostatic hyperplasia, familial adenomatosis polyposis, neuro-fibromatosis, atherosclerosis, pulmonary fibrosis, arthritis, psoriasis, glomerulonephritis, restenosis following angioplasty or vascular surgery, hypertrophic scar formation, inflammatory bowel disease, transplantation rejection, endotoxic shock, and fungal
- 10 infections.
- defective apoptosis-associated conditions, such as cancers (including those types mentioned hereinabove), viral infections (including herpes virus, pox virus, Epstein-Barr virus, Sindbis virus and adenovirus), prevention of AIDS
- 15 development in HIV-infected individuals, autoimmune diseases (including systemic lupus erythematosus, rheumatoid arthritis, psoriasis, autoimmune mediated glomerulonephritis, inflammatory bowel disease and autoimmune diabetes mellitus), neurodegenerative disorders (including Alzheimer's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, Parkinson's disease, AIDS-related dementia, spinal muscular atrophy and cerebellar degeneration), myelodysplastic syndromes,
- 20 aplastic anemia, ischemic injury associated with myocardial infarctions, stroke and reperfusion injury, arrhythmia, atherosclerosis, toxin-induced or alcohol related liver diseases, hematological diseases (including chronic anemia and aplastic anemia), degenerative diseases of the musculoskeletal system (including osteoporosis and arthritis), aspirin-sensitive rhinosinusitis, cystic fibrosis, multiple
- 25 sclerosis, kidney diseases and cancer pain.

The agents of the invention may also be useful in the inhibition of the development of invasive cancer, tumor angiogenesis and metastasis.

Moreover, the agents of the invention, as inhibitors of CDKs, can modulate the level of cellular RNA and DNA synthesis and therefore are useful in the treatment of viral infections

30 such as HIV, human papilloma virus, herpes virus, Epstein-Barr virus, adenovirus, Sindbis virus, pox virus and the like.

The terms "comprising" and "including" are used herein in their open, non-limiting sense.

The term "alkyl" as used herein refers to straight- and branched-chain alkyl groups having

35 from one to twelve carbon atoms. Exemplary alkyl groups include methyl (Me), ethyl, n-propyl,


isopropyl, butyl, isobutyl, sec-butyl, tert-butyl (tBu), pentyl, isopentyl, tert-pentyl, hexyl, isohexyl, and the like.

The term "heteroalkyl" as used herein refers to straight- and branched-chain alkyl groups having from one to twelve atoms containing one or more heteroatoms selected from S, O, and N.

5 The term "alkenyl" refers to straight- and branched-chain alkenyl groups having from two to twelve carbon atoms. Illustrative alkenyl groups include prop-2-enyl, but-2-enyl, but-3-enyl, 2-methylprop-2-enyl, hex-2-enyl, and the like.

The term "alkynyl" refers to straight- and branched-chain alkynyl groups having from two to twelve carbon atoms. Illustrative alkynyl groups include prop-2-ynyl, but-2-ynyl, but-3-ynyl, 2-methylbut-2-ynyl, hex-2-ynyl, and the like.


10 The term "aryl" (Ar) refers to monocyclic and polycyclic aromatic ring structures containing only carbon and hydrogen. Illustrative examples of aryl groups include the following moieties:




15

The term "heteroaryl" (heteroAr) refers to monocyclic, or fused polycyclic aromatic ring structures which include one or more heteroatoms selected from nitrogen, oxygen and sulfur and having from 3 to 12 ring atoms per ring. The polycyclic heteroaryl group may be fused or non-fused. More preferably, illustrative examples of heteroaryl groups have from 4 to 7 ring atoms per ring, such as the following moieties:


20









- The term "cycloalkyl" refers to saturated carbocycles having from three to twelve carbon atoms, including bicyclic and tricyclic cycloalkyl structures. Illustrative examples of cycloalkyl groups include the following moieties:



10

- A "heterocycloalkyl" group refers to a monocyclic or fused or spiro polycyclic ring structure radical which may be saturated or unsaturated and contains from three to twelve ring atoms, selected from carbon and heteroatoms, preferably 4 or 5 ring carbon atoms, and at least one heteroatom selected from nitrogen, oxygen and sulfur. The radicals may be fused with an aryl or heteroaryl. More preferably, illustrative examples of heterocycloalkyl groups have 4 to 7 ring atoms per ring, such as the following moieties,





The term "alkoxy" refers to the radical  $-O-R$  where R is an alkyl as defined above. Examples of alkoxy groups include methoxy, ethoxy, propoxy, and the like.

5        The term "halogen" represents chlorine, fluorine, bromine or iodine. The term "halo" represents chloro, fluoro, bromo or iodo.

The term "alcohol" refers to the radical  $-R-OH$  where R is alkyl, alkenyl, alkynyl, Ar, heteroaryl, heterocycloalkyl, or cycloalkyl as defined above. Examples of alcohols include methanol, ethanol, propanol, phenol and the like.

10       The term "acyl" represents  $-C(O)R$ ,  $-C(O)OR$ ,  $-OC(O)R$  or  $-OC(O)OR$  where R is alkyl, alkenyl, alkynyl, Ar, heteroaryl, heterocycloalkyl, or cycloalkyl as defined as above.

The term "amide" refers to the radical  $-C(O)N(R')(R'')$  where R' and R'' are each independently selected from hydrogen, alkyl, alkenyl, alkynyl,  $-OH$ , alkoxy, cycloalkyl, heterocycloalkyl, heteroaryl, aryl as defined above; or R' and R'' cyclize together with the  
15       nitrogen to form a heterocycloalkyl or heteroaryl as defined above.

The term "substituted" as used herein means that the group in question, e.g., alkyl group, etc., may bear one or more substituents.

The alkyl, cycloalkyl, aryl, heterocycloalkyl and heteroaryl groups and the substituents containing these groups, as defined hereinabove, may be optionally substituted by at least one other  
20       substituent. The term "optionally substituted" is intended to expressly indicate that the specified group is unsubstituted or substituted by one or more substituents from the list above. Various groups may be unsubstituted or substituted (i.e., they are optionally substituted) as indicated.

If the substituents themselves are not compatible with the synthetic methods of this invention, the substituent may be protected with a suitable protecting group that is stable to the  
25       reaction conditions used in these methods. The protecting group may be removed at a suitable point in the reaction sequence of the method to provide a desired intermediate or target compound. Suitable protecting groups and the methods for protecting and de-protecting different substituents using such suitable protecting groups are well known to those skilled in the art; examples of which may be found in T. Greene and P. Wuts, *Protecting Groups in Chemical Synthesis* (3rd ed.), John  
30       Wiley & Sons, NY (1999), which is incorporated herein by reference in its entirety. In some instances, a substituent may be specifically selected to be reactive under the reaction conditions used in the methods of this invention. Under these circumstances, the reaction conditions convert

the selected substituent into another substituent that is either useful in an intermediate compound in the methods of this invention or is a desired substituent in a target compound.

Some of the inventive compounds may exist in various stereoisomeric or tautomeric forms. The present invention encompasses all such cell proliferation-inhibiting compounds, including active compounds in the form of single pure enantiomers (i.e., essentially free of other stereoisomers), racemates, mixtures of enantiomers and/or diastereomers, and/or tautomers. Preferably, the inventive compounds that are optically active are used in optically pure form.

As generally understood by those skilled in the art, an optically pure compound having one chiral center (i.e., one asymmetric carbon atom) is one that consists essentially of one of the two possible enantiomers (i.e., is enantiomerically pure), and an optically pure compound having more than one chiral center is one that is both diastereomerically pure and enantiomerically pure.

Preferably, the compounds of the present invention are used in a form that is at least 90% optically pure, that is, a form that contains at least 90% of a single isomer (80% enantiomeric excess ("e.e.") or diastereomeric excess ("d.e.")), more preferably at least 95% (90% e.e. or d.e.), even more preferably at least 97.5% (95% e.e. or d.e.), and most preferably at least 99% (98% e.e. or d.e.).

Additionally, the formulae are intended to cover solvated as well as unsolvated forms of the identified structures. For example, Formula I includes compounds of the indicated structure in both hydrated and non-hydrated forms. Additional examples of solvates include the structures in combination with isopropanol, ethanol, methanol, DMSO, ethyl acetate, acetic acid, or ethanolamine.

In addition to compounds of Formula I, the invention includes pharmaceutically acceptable prodrugs, pharmaceutically active metabolites, and pharmaceutically acceptable salts of such compounds and metabolites.

The term "pharmaceutically acceptable" means pharmacologically acceptable and substantially non-toxic to the subject being administered the agent.

"A pharmaceutically acceptable prodrug" is a compound that may be converted under physiological conditions or by solvolysis to the specified compound or to a pharmaceutically acceptable salt of such compound. "A pharmaceutically active metabolite" is intended to mean a pharmacologically active product produced through metabolism in the body of a specified compound or salt thereof. Prodrugs and active metabolites of a compound may be identified using routine techniques known in the art. See, e.g., Bertolini et al., *J. Med. Chem.*, 40, 2011-2016 (1997); Shan et al., *J. Pharm. Sci.*, 86 (7), 765-767; Bagshawe, *Drug Dev. Res.*, 34, 220-230 (1995); Bodor, *Advances in Drug Res.*, 13, 224-331 (1984); Bundgaard, *Design of Prodrugs*

(Elsevier Press 1985); and Larsen, *Design and Application of Prodrugs, Drug Design and Development* (Krogsgaard-Larsen et al., eds., Harwood Academic Publishers, 1991).

“A pharmaceutically acceptable salt” is intended to mean a salt that retains the biological effectiveness of the free acids and bases of the specified compound and that is not biologically or otherwise undesirable. A compound of the invention may possess a sufficiently acidic, a  
5 sufficiently basic, or both functional groups, and accordingly react with any of a number of inorganic or organic bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt. Exemplary pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a mineral or organic acid or an inorganic base, such as  
10 salts including sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, monohydrogenphosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrate, caproates, heptanoates, propiolates, oxalates, malonates, succinates, suberates, sebacates, fumarates, maleates, butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates,  
15 methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, phenylpropionates, phenylbutyrates, citrates, lactates,  $\gamma$ -hydroxybutyrates, glycolates, tartrates, methane-sulfonates, propanesulfonates, naphthalene-1-sulfonates, naphthalene-2-sulfonates, and mandelates.

If the inventive compound is a base, the desired pharmaceutically acceptable salt may  
20 be prepared by any suitable method available in the art, for example, treatment of the free base with an inorganic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, sulfamic acid, nitric acid, phosphoric acid and the like, or with an organic acid, such as acetic acid, phenylacetic acid, propionic acid, stearic acid, lactic acid, ascorbic acid, maleic acid, hydroxymaleic acid, isethionic acid, succinic acid, mandelic acid, fumaric acid, malonic acid,  
25 pyruvic acid, oxalic acid, glycolic acid, salicylic acid, a pyranosidyl acid, such as glucuronic acid or galacturonic acid, an alpha-hydroxy acid, such as citric acid or tartaric acid, an amino acid, such as aspartic acid or glutamic acid, an aromatic acid, such as benzoic acid, 2-acetoxybenzoic acid or cinnamic acid, a sulfonic acid, such as p-toluenesulfonic acid, methanesulfonic acid or ethanesulfonic acid, or the like.

30 If the inventive compound is an acid, the desired pharmaceutically acceptable salt may be prepared by any suitable method, for example, treatment of the free acid with an inorganic or organic base, such as an amine (primary, secondary or tertiary), an alkali metal hydroxide or alkaline earth metal hydroxide, or the like. Illustrative examples of suitable salts include organic salts derived from amino acids, such as glycine and arginine, ammonia, carbonates,  
35 bicarbonates, primary, secondary, and tertiary amines, and cyclic amines, such as

benzylamines, pyrrolidines, piperidine, morpholine and piperazine, and inorganic salts derived from sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum and lithium.

Pharmaceutical compositions according to the invention may, alternatively or in addition to  
5 a compound of Formula I, comprise as an active ingredient pharmaceutically acceptable prodrugs, pharmaceutically active metabolites, and pharmaceutically acceptable salts of such compounds and metabolites. Such compounds, prodrugs, multimers, salts, and metabolites are sometimes referred to herein collectively as "active agents" or "agents."

In the case of agents that are solids, it is understood by those skilled in the art that the  
10 inventive compounds and salts may exist in different crystal or polymorphic forms, all of which are intended to be within the scope of the present invention and specified formulas.

Therapeutically effective amounts of the active agents of the invention may be used to treat diseases mediated by modulation or regulation of protein kinases. An "effective amount" is intended to mean that amount of an agent that significantly inhibits proliferation and/or prevents  
15 de-differentiation of a eukaryotic cell, e.g., a mammalian, insect, plant or fungal cell, and is effective for the indicated utility, e.g., specific therapeutic treatment.

The amount of a given agent that will correspond to such an amount will vary depending upon factors such as the particular compound, disease condition and its severity, the identity (e.g., weight) of the subject or host in need of treatment, but can nevertheless be routinely determined in  
20 a manner known in the art according to the particular circumstances surrounding the case, including, e.g., the specific agent being administered, the route of administration, the condition being treated, and the subject or host being treated. "Treating" is intended to mean at least the mitigation of a disease condition in a subject such as mammal (e.g., human), that is affected, at least in part, by the activity of one or more kinases, for example protein kinases such as tyrosine  
25 kinases, and includes: preventing the disease condition from occurring in a mammal, particularly when the mammal is found to be predisposed to having the disease condition but has not yet been diagnosed as having it; modulating and/or inhibiting the disease condition; and/or alleviating the disease condition.

Agents that potently regulate, modulate, or inhibit cell proliferation are preferred. For  
30 certain mechanisms, inhibition of the protein kinase activity associated with CDK complexes, among others, and those which inhibit angiogenesis and/or inflammation are preferred. The present invention is further directed to methods of modulating or inhibiting protein kinase activity, for example in mammalian tissue, by administering an inventive agent. The activity of agents as anti-proliferatives is easily measured by known methods, for example by using whole  
35 cell cultures in an MTT assay. The activity of the inventive agents as modulators of protein

kinase activity, such as the activity of kinases, may be measured by any of the methods available to those skilled in the art, including in vivo and/or in vitro assays. Examples of suitable assays for activity measurements include those described in International Publication No. WO 99/21845; Parast et al., *Biochemistry*, 37, 16788-16801 (1998); Connell-Crowley and Harpes, *Cell Cycle: Materials and Methods*, (Michele Pagano, ed. Springer, Berlin, Germany)(1995); International Publication No. WO 97/34876; and International Publication No. WO 96/14843. These properties may be assessed, for example, by using one or more of the biological testing procedures set out in the examples below.

The active agents of the invention may be formulated into pharmaceutical compositions as described below. Pharmaceutical compositions of this invention comprise an effective modulating, regulating, or inhibiting amount of a compound of Formula I or Formula II and an inert, pharmaceutically acceptable carrier or diluent. In one embodiment of the pharmaceutical compositions, efficacious levels of the inventive agents are provided so as to provide therapeutic benefits involving anti-proliferative ability. By "efficacious levels" is meant levels in which proliferation is inhibited, or controlled. These compositions are prepared in unit-dosage form appropriate for the mode of administration, e.g., parenteral or oral administration.

An inventive agent can be administered in conventional dosage form prepared by combining a therapeutically effective amount of an agent (e.g., a compound of Formula I) as an active ingredient with appropriate pharmaceutical carriers or diluents according to conventional procedures. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation.

The pharmaceutical carrier employed may be either a solid or liquid. Exemplary of solid carriers are lactose, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or diluent may include time-delay or time-release material known in the art, such as glyceryl monostearate or glyceryl distearate alone or with a wax, ethylcellulose, hydroxypropylmethylcellulose, methylmethacrylate and the like.

A variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier may vary, but generally will be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation will be in the form of syrup, emulsion, soft gelatin capsule, sterile injectable solution or suspension in an ampoule or vial or non-aqueous liquid suspension.

To obtain a stable water-soluble dose form, a pharmaceutically acceptable salt of an inventive agent can be dissolved in an aqueous solution of an organic or inorganic acid, such as

0.3M solution of succinic acid or citric acid. If a soluble salt form is not available, the agent may be dissolved in a suitable cosolvent or combinations of cosolvents. Examples of suitable cosolvents include, but are not limited to, alcohol, propylene glycol, polyethylene glycol 300, polysorbate 80, glycerin and the like in concentrations ranging from 0-60% of the total volume. In an exemplary embodiment, a compound of Formula I is dissolved in DMSO and diluted with water. The composition may also be in the form of a solution of a salt form of the active ingredient in an appropriate aqueous vehicle such as water or isotonic saline or dextrose solution.

It will be appreciated that the actual dosages of the agents used in the compositions of this invention will vary according to the particular complex being used, the particular composition formulated, the mode of administration and the particular site, host and disease being treated. Optimal dosages for a given set of conditions can be ascertained by those skilled in the art using conventional dosage-determination tests in view of the experimental data for an agent. For oral administration, an exemplary daily dose generally employed is from about 0.001 to about 1000 mg/kg of body weight, with courses of treatment repeated at appropriate intervals. Administration of prodrugs is typically dosed at weight levels which are chemically equivalent to the weight levels of the fully active form.

The compositions of the invention may be manufactured in manners generally known for preparing pharmaceutical compositions, e.g., using conventional techniques such as mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing. Pharmaceutical compositions may be formulated in a conventional manner using one or more physiologically acceptable carriers, which may be selected from excipients and auxiliaries that facilitate processing of the active compounds into preparations which can be used pharmaceutically.

Proper formulation is dependent upon the route of administration chosen. For injection, the agents of the invention may be formulated into aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the compounds with pharmaceutically acceptable carriers known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

Pharmaceutical preparations for oral use can be obtained using a solid excipient in admixture with the active ingredient (agent), optionally grinding the resulting mixture, and processing the mixture of granules after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable

excipients include: fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; and cellulose preparations, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as crosslinked  
5 polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, polyvinyl pyrrolidone, Carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for  
10 identification or to characterize different combinations of agents.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the agents in admixture with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate, and, optionally,  
15 stabilizers. In soft capsules, the agents may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration. For buccal administration, the compositions take the form of tablets or lozenges formulated in conventional manners.

For administration intranasally or by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g.,  
dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by  
25 providing a valve to deliver a metered amount. Capsules and cartridges of gelatin for use in an inhaler or insufflator and the like may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit-dosage  
30 form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the agents in water-soluble form. Additionally, suspensions of the agents may be prepared as  
35 appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils



such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For administration to the eye, the agent is delivered in a pharmaceutically acceptable ophthalmic vehicle such that the compound is maintained in contact with the ocular surface for a sufficient time period to allow the compound to penetrate the corneal and internal regions of the eye, for example, the anterior chamber, posterior chamber, vitreous body, aqueous humor, vitreous humor, cornea, iris/ciliary, lens, choroid/retina and sclera. The pharmaceutically acceptable ophthalmic vehicle may be an ointment, vegetable oil, or an encapsulating material. A compound of the invention may also be injected directly into the vitreous and aqueous humor.

Alternatively, the agents may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use. The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described above, the agents may also be formulated as a depot preparation. Such long-acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion-exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

An exemplary pharmaceutical carrier for hydrophobic compounds is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system may be a VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) contains VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g. polyvinyl pyrrolidone; and other sugars or polysaccharides may be substituted for dextrose.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered  
5 using a sustained-release system, such as semi-permeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein  
10 stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid- or gel-phase carriers or excipients. Examples of such carriers or excipients include calcium carbonate, calcium phosphate, sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Some of the compounds of the invention may be provided as salts with pharmaceutically  
15 compatible counter ions. Pharmaceutically compatible salts may be formed with many acids, including hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free-base forms.

The agents of the invention may be useful in combination with known anti-cancer treatments such as: DNA interactive agents such as cisplatin or doxorubicin; topoisomerase II  
20 inhibitors such as etoposide; topoisomerase I inhibitors such as CPT-11 or topotecan; tubulin interacting agents such as paclitaxel, docetaxel or the epothilones; hormonal agents such as tamoxifen; thymidilate synthase inhibitors such as 5-fluorouracil; and anti-metabolites such as methotrexate. They may be administered together or sequentially, and when administered sequentially, the agents may be administered either prior to or after administration of the known  
25 anticancer or cytotoxic agent.

The agents may be prepared using the reaction routes and synthesis schemes as described below, employing the general techniques known in the art using starting materials that are readily available. The preparation of preferred compounds of the present invention is described in detail in the following examples, but the artisan will recognize that the chemical  
30 reactions described may be readily adapted to prepare a number of other anti-proliferatives or protein kinase inhibitors of the invention. For example, the synthesis of non-exemplified compounds according to the invention may be successfully performed by modifications apparent to those skilled in the art, e.g., by appropriately protecting interfering groups, by changing to other suitable reagents known in the art, or by making routine modifications of

reaction conditions. Alternatively, other reactions disclosed herein or generally known in the art will be recognized as having applicability for preparing other compounds of the invention.

### EXAMPLES

5           In the examples described below, unless otherwise indicated, all temperatures are set forth in degrees Celsius and all parts and percentages are by weight. Reagents were purchased from commercial suppliers such as Sigma-Aldrich Chemical Company or Lancaster Synthesis Ltd. and were used without further purification unless otherwise indicated. Tetrahydrofuran (THF) and N, N-dimethylformamide (DMF) were purchased from Aldrich in Sure Seal bottles and used as  
10       received. All solvents were purified using standard methods known to those skilled in the art, unless otherwise indicated.

          The reactions set forth below were done generally under a positive pressure of argon at an ambient temperature (unless otherwise stated) in anhydrous solvents, and the reaction flasks were fitted with rubber septa for the introduction of substrates and reagents via syringe. Glassware was  
15       oven dried and/or heat dried. Analytical thin layer chromatography (TLC) was performed on glass-backed silica gel 60 F 254 plates from Analtech (0.25 mm), eluted with the appropriate solvent ratios (v/v), and are denoted where appropriate. The reactions were assayed by TLC, NMR, or analytical HPLC and terminated as judged by the consumption of starting material.

          Visualization of the TLC plates was done with iodine vapor, ultraviolet illumination,  
20       2%  $\text{Ce}(\text{NH}_4)_4(\text{SO}_4)_4$  in 20% aqueous sulfuric acid, 2% ninhydrin in ethanol, or p-anisaldehyde spray reagent, and activated with heat where appropriate. Work-ups were typically done by doubling the reaction volume with the reaction solvent or extraction solvent and then washing with the indicated aqueous solutions using 25% by volume of the extraction volume unless otherwise indicated. Product solutions were dried over anhydrous  $\text{Na}_2\text{SO}_4$  and/or  $\text{MgSO}_4$  prior to filtration  
25       and evaporation of the solvents under reduced pressure on a rotary evaporator and noted as solvents removed in vacuo. Flash column chromatography (Still et al., *J. Org. Chem.*, 43, 2923 (1978)) was done using Merck silica gel (47-61  $\mu\text{m}$ ) with a silica gel crude material ratio of about 20:1 to 50:1, unless otherwise stated. Hydrogenolysis was done at the pressure indicated in the examples or at ambient pressure.

30        $^1\text{H}$ -NMR spectra were recorded on a Bruker or Varian instrument operating at 300 MHz and  $^{13}\text{C}$ -NMR spectra were recorded operating at 75 MHz. NMR spectra were obtained as  $\text{CDCl}_3$  solutions (reported in ppm), using chloroform as the reference standard (7.27 ppm and 77.00 ppm) unless otherwise noted. When peak multiplicities are reported, the following abbreviations are used: s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), br (broadened multiplet), bs

(broadened singlet), dd (doublet of doublets), dt (doublet of triplets). Coupling constants, when given, are reported in Hertz (Hz).

Infrared (IR) spectra were recorded on a Perkin-Elmer FT-IR Spectrometer as neat oils, as KBr pellets, or as  $\text{CDCl}_3$  solutions, and when given are reported in wave numbers ( $\text{cm}^{-1}$ ). The mass spectra were obtained using LSIMS, FAB, MALDI, or electrospray (ESIMS). All melting points (mp) are uncorrected.


Mass spectrometry (MS) was conducted with various techniques. Matrix-Assisted Laser Desorption/Ionization Fourier Transform Mass Spectrometry (MALDI FTMS), was performed on an IonSpec FTMS mass spectrometer. Samples are irradiated with a nitrogen laser (Laser Science Inc.) operated at 337nm and the laser beam is attenuated by a variable attenuator and focused on the sample target. The ions are then differentiated according to their  $m/z$  using an ion cyclotron resonance mass analyzer. The electrospray ionization (ESI) mass spectrometry experiments were performed on an API 100 Perkin Elmer SCIEX single quadrupole mass spectrometer. Electrospray samples are typically introduced into the mass analyzer at a rate of 4.0  $\mu\text{l}/\text{minute}$ . The positive and negative ions, generated by charged droplet evaporation, enter the analyzer through an interface plate and a 100 mm orifice, while the declustering potential is maintained between 50 and 200V to control the collisional energy of the ions entering the mass analyzer. The emitter voltage is typically maintained at 4000V. The liquid chromatography (LC) electrospray ionization (ESI) mass spectrometry experiments are performed on an Hewlett-Packard (HP) 1100 MSD single quadrupole mass spectrometer. Electrospray samples are typically introduced into the mass analyzer at a rate of 100 to 1000  $\mu\text{l}/\text{minute}$ . The positive and negative ions, generated by charged droplet evaporation, enter the analyzer through a heated capillary plate, while the declustering potential is maintained between 100 and 300V to control the collisional energy of the ions entering the mass analyzer. The emitter voltage is typically maintained at 4000V.

Reserved phase preparative HPLC purification was performed on Gilson 321 system, using a C18 reversed phase preparative column (Metasil AQ 10  $\mu$ , C18, 120A 250 x 21.2 mm, MetaChem) in a gradient from 0.1%TFA/5% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  to 0.1%TFA/5% $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  over 20 minutes at a flow rate of 20ml/min.

In Examples, the compounds may be obtained in the form of salts or free bases. When the compounds are obtained in the form of TFA salts, they were obtained by lyophilizing the HPLC fractions of the corresponding compounds. When the compounds are obtained in the form of the HCl salts, they were obtained by lyophilizing the HPLC fractions of the corresponding compounds in the presence of excess hydrochloric acid. When the compounds are obtained in the form of the free bases, they were obtained by concentrating ethyl acetate extracts of HPLC fractions, neutralized with  $\text{Na}_2\text{CO}_3$ .


Compounds in accordance with the invention may be prepared in manners analogous to those specifically described below, with the lettered example prefixes (i.e., A, B, C, D, E, F and G) designating general synthesis schemes.

5

**Method A**

10

The common starting material, 2-bromo-2',6'-difluoroacetophenone, was prepared as follows.

*2-Bromo-2',6'-difluoroacetophenone*

15

To a mechanically stirring solution of 2',6'-difluoroacetophenone (100.0 g, 640.0 mmol; Melford Laboratories, Ltd.) in ethyl acetate (1300 ml) was added freshly milled copper(II) bromide (300 g, 1.35 mol) and bromine (1.6 ml, 32 mmol). The mixture was heated at reflux for 2.25 hours and allowed to cool to room temperature. The resultant green mixture was filtered and the solids rinsed with ethyl acetate (4×100 ml). The filtrate was concentrated with a rotary evaporator at <40°C under reduced pressure, diluted with methyl t-butyl ether (MTBE; 650 ml), filtered through a pad of silica gel (230–400  $\mu$ ; 9.5 cm diam.× 4 cm. ht.), and solids rinsed with MTBE (5×200 ml). Concentration of the filtrate gave a pale green oil, which was purified by fractional vacuum distillation to give 117 g of pale yellow oil, bp 88–97°C (2.0 mm Hg) in 78% yield. The results

matched that previously described in PCT Patent Publication WO99/21845 (in Example C(79)) and was used without any further purification or characterization.

<sup>1</sup>H NMR: δ 7.48 (ddd, 1H, J=6.3, 8.5, 14.8 Hz), 7.01 (ddd, 2H, J=4.6, 5.8, 16.6 Hz), 4.37 (t, 2H, J=0.7 Hz).

5 Step 1. *4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic Acid Ethyl Ester (2)*

To a mechanically stirring mixture of 4-ethoxycarbonylphenyl isothiocyanate (**1**; 25.0 g, 121 mmol; TransWorld), cyanamide (5.32 g, 127 mmol), and acetonitrile (150 ml) at 2.6°C was added dropwise over 12 minutes DBU (1,8-diazabicyclo [5.4.0]-7-undecene; 19.8 ml, 133 mmol). After 2 hours at 0°C, a solution of 2-bromo-2',6'-difluoro-acetophenone (29.8 g, 127 mmol) in  
10 acetonitrile (25 ml) was added dropwise over 13 minutes. Upon warming to ambient temperature, the resultant mixture turned into a dark red solution, and was allowed to stir overnight. Water (175 ml) was added over 5 minutes, and the resultant yellow suspension stirred for one hour. The yellow solid was rinsed with acetonitrile/water (1/1; 3 × 75 ml) and dried under high vacuum at 45°C to give 45.39 g of yellow solid in 93% yield.

15 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.24 (bs, 1H), 8.30 (br, 1H), 8.06 (d, 2H, J=8.7 Hz), 7.82 (d, 2H, J=8.7 Hz), 7.66 (m, 1H), 7.30 (t, 2H, J=8.7 Hz), 4.38 (t, 3H, J=5.6 Hz), 1.38 (t, 3H, J=5.6 Hz).

Step 2. *4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic Acid (3)*

To a mechanically stirring solution of 4-{4-amino-5-[1-(2,6-difluoro-benzoyl)]-thiazol-2-ylamino}-benzoic acid ethyl ester (**2**, 45.7g, 113 mmol) in methanol (250 ml) was added 3N  
20 aqueous NaOH solution (378 ml). An exotherm resulted and the resultant yellow solution at 31°C was allowed to stir at room temperature overnight. The methanol was removed under reduced pressure and the resultant aqueous solution was washed with hexane (2×150 ml). The aqueous layer was cooled to 5°C, adjusted to pH4.0 with 6N HCl (~200 ml), and diluted with water (100 ml). The bright yellow solid was filtered off, washed with water (3×250 mL), and dried under  
25 vacuum at 45°C to provide 38.65 g of canary yellow solid in 91% yield.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.18 (bs, 1H), 8.20 (br, 1H), 7.90 (d, 2H, J=8.8Hz), 7.72 (d, 2H, J=8.8Hz), 7.60 (m, 1H), 7.24 (t, 2H, J=8.7Hz).

Anal. Calcd. for C<sub>17</sub>H<sub>11</sub>F<sub>2</sub>N<sub>3</sub>O<sub>3</sub>S • 1.25 H<sub>2</sub>O: C, 51.32%; H, 3.42%; N, 10.56%; S, 8.06%. Found: C, C, 51.32%; H, 3.42%; N, 10.56%; S, 8.06%.

30 Step 3. *4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzamide(4)*

General Method

To a solution of 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**; 1 eq), appropriate amine (1.2 eq) and DIEA (N,N-diisopropylethylamine; 6 eq) in DMF, PyBop (benzotriazol-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate; 1.1 eq) was added. The reaction mixture was stirred at room temperature for 3 hours.

- 5 DMF was removed under reduced pressure and a solution of the resultant residue in ethyl acetate was extracted with saturated aqueous NaHCO<sub>3</sub>, brine, dried with MgSO<sub>4</sub>, filtered and concentrated to give a crude product that was further purified by either silica gel chromatography or reversed phase preparative HPLC.

- 10 Reversed phase preparative HPLC purification was performed on Gilson 321 system, using a C18 reversed phase preparative column (Metasil AQ 10 □, C18, 120A 250 x 21.2 mm, MetaChem) with a gradient from 0.1%TFA/5%CH<sub>3</sub>CN/H<sub>2</sub>O to 0.1%TFA/5%H<sub>2</sub>O/CH<sub>3</sub>CN over 20 minutes at a flow rate of 20 ml/min.


The TFA salts were obtained by lyophilizing the HPLC fractions of the corresponding compounds.

- 15 The HCl salts were obtained by lyophilizing the HPLC fractions of the corresponding compounds in the presence of excess hydrochloric acid.

The free bases were obtained by concentrating ethyl acetate extracts of HPLC fractions, neutralized with Na<sub>2</sub>CO<sub>3</sub>.

- 20 The compounds of the following Examples A1-A62 were prepared in a manner similar to Step 3 in Method A from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid **3** and appropriate amines (R-NH<sub>2</sub>) with PyBop, HATU [O-(7-azabenzotriazol-1-yl)-N, N, N', N'-tetramethyluronium hexafluoro-phosphate] or other similar coupling reagents.

**Example A1:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-morpholin-4-yl-ethyl)-benzamide




25

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ7.88 (d, 2H, J=8.9 Hz), 7.76 (d, 2H, J=8.9 Hz), 7.54-7.43 (m, 1H), 7.08 (m, 2H), 3.84-3.75 (m, 4H), 3.62 (t, 2H, J=6.4 Hz), 2.98-2.78 (m, 6H).

HRFABMS: Calcd for C<sub>23</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>3</sub>S (M+Na<sup>+</sup>): 510.1387. Found: 510.1382.

Anal. Calcd. for  $C_{23}H_{23}F_2N_5O_3S \cdot 0.82 \text{ MeOH} \cdot 0.12 \text{ CH}_3\text{CN}$ : C, 55.71%; H, 5.18%; N, 13.83%; S, 6.18%. Found: C, 55.69%; H, 5.17%; N, 13.84%; S, 6.11%.

**Example A2:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-phenylamino-ethyl)-benzamide




5

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  11.06 (s, 1H), 8.50 (t, 1H,  $J=2.8$  Hz), 8.19 (bs, 2H), 7.86 (d, 2H,  $J=8.7$  Hz), 7.68 (d, 2H,  $J=8.7$  Hz), 7.58 (m, 1H), 7.22 (m, 2H), 7.08 (t, 2H,  $J=7.8$  Hz), 6.61 (d, 2H,  $J=7.8$  Hz), 6.52 (t, 1H,  $J=7.2$  Hz), 5.71 (br, 1H), 3.42 (m, 2H), 3.22 (m, 2H).

HRFABMS: Calcd for  $C_{25}H_{21}F_2N_5O_2S$  ( $M+H^+$ ): 494.1462. Found: 494.1444.


10 **Example A3:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-ethyl)-N-methyl-benzamide.



$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.74 (d, 2H,  $J=8.6$  Hz), 7.53-7.39 (m, 3H), 7.07 (t, 2H,  $J=8.7$  Hz), 3.68 (m, 1H), 3.46 (m, 1H), 3.05 (s, 3H), 2.64 (m, 1H), 2.52 (m, 1H), 2.37 (s, 3H), 2.09 (s, 3H).

15 HRFABMS: Calcd for  $C_{22}H_{23}F_2N_5O_2S$  ( $M+H^+$ ): 460.1619. Found: 460.1612.

**Example A4:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-ethyl)-benzamide Hydrochloride Salt




20  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.84 (d, 2H,  $J=8.8$  Hz), 7.73 (d, 2H,  $J=8.8$  Hz), 7.52-7.83 (m, 1H), 7.03 (t, 2H,  $J=7.9$  Hz), 3.75 (t, 2H,  $J=5.8$  Hz), 3.38 (t, 2H,  $J=5.9$ ), 2.98 (s, 6H).

FABMS ( $M+H^+$ ): 446; ( $M-H$ ): 444.



Anal. Calcd. for  $C_{21}H_{21}F_2N_5O_2S \cdot 0.7 H_2O \cdot 0.08 EtOAc \cdot 1.55 HCl$ : C, 49.09%; H, 4.75%; N, 13.43%; S, 6.15%. Found: C, 49.07%; H, 4.62%; N, 13.43%; S, 6.15%.

**Example A5:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(6-hydroxy-6-methyl-hept-2-yl)-benzamide



5


$^1H$  NMR ( $CD_3OD$ ):  $\delta$  7.82 (d, 2H,  $J=8.8$  Hz), 7.74 (d, 2H,  $J=8.8$  Hz), 7.50 (m, 1H), 7.02-7.13 (m, 2H), 4.14 (m, 1H), 1.68-1.37 (m, 6H), 1.23 (d, 3H,  $J=6.4$  Hz), 1.18 (s, 3H), 1.17 (s, 3H).

FABMS ( $M+H^+$ ): 503; ( $M-H^-$ ): 501.

Anal. Calcd. For  $C_{25}H_{28}F_2N_4O_3S \cdot 0.20 H_2O \cdot 0.24 EtOAc$ : C, 59.13%; H, 5.80%; N, 10.63%; S, 6.08%. Found: C, 59.13%; H, 5.82%; N, 10.77%; S, 6.06%.

10

**Example A6:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(5-methyl-furan-2-ylmethyl)-benzamide.




15  $^1H$  NMR ( $DMSO-d_6$ ):  $\delta$  11.06 (s, 1H), 8.83 (br, 1H), 8.22 (bs, 2H), 7.88 (d, 2H,  $J=8.7$  Hz), 7.67 (d, 2H,  $J=8.7$  Hz), 7.55 (m, 1H), 7.21 (t, 2H,  $J=7.5$  Hz), 6.13 (d, 1H,  $J=3.0$  Hz), 5.99 (d, 1H,  $J=3.0$  Hz), 4.39 (d, 2H,  $J=5.6$  Hz), 2.21 (s, 3H).

FABMS ( $M+H^+$ ): 469; ( $M-H^-$ ): 467.

Anal. Calcd. for  $C_{23}H_{18}F_2N_4O_3S \cdot 0.10 H_2O \cdot 0.12 EtOAc$ : C, 58.65%; H, 4.02%; N, 11.65%; S, 6.67%. Found: C, 58.66%; H, 4.02%; N, 11.85%; S, 6.68%.

20


**Example A7:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-isopropoxy-ethyl)-benzamide.



$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.06 (s, 1H), 8.41 (br, 1H), 8.22 (bs, 2H), 7.84 (d, 2H,  $J=8.7$  Hz), 7.64 (d, 2H,  $J=8.7$  Hz), 7.56 (m, 1H), 7.21 (t, 2H,  $J=8.7$  Hz), 3.56 (m, 1H), 3.48 (t, 2H,  $J=6.0$  Hz), 3.35 (m, 2H), 1.09 (d, 6H,  $J=6.1$  Hz).

5 FABMS ( $M+H^+$ ): 461; ( $M-H^-$ ): 459.

**Example A8:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(pyrrolidin-2S-yl-methyl)-benzamide.



The intermediate 2S-([1-(4-{4-amino-5-[1-(2,6-difluoro-phenyl)-methanoyl]-thiazol-2-ylamino}-phenyl)-methanoyl]-amino)-methyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester was prepared in a manner similar to Step 3 in Method A from 2S-aminomethyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester (AstaTech, Inc.).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 8.45 (br, 1H), 7.84 (d, 2H,  $J=8.6$  Hz), 7.39-7.24 (m, 3H), 6.95 (m, 2H), 4.20 (m, 1H), 3.54-3.31 (m, 4H), 2.04-1.65 (m, 2H)


15 The title compound was prepared by treating the above intermediate with 30 % TFA/ $\text{CH}_2\text{Cl}_2$  in 30 minutes, followed by HPLC purification.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.85 (d, 2H,  $J=9.0$  Hz), 7.74 (d, 2H,  $J=9.0$  Hz), 7.50 (m, 1H), 7.09 (t, 2H,  $J=8.0$  Hz), 3.55-3.38 (m, 3H), 3.12-2.90 (m, 2H), 2.07-1.77 (m, 3H), 1.59 (m, 1H).

FABMS ( $M+H^+$ ): 458; ( $M-H^-$ ): 456.

20 Anal. Calcd. for  $\text{C}_{22}\text{H}_{21}\text{F}_2\text{N}_5\text{O}_2\text{S} \cdot 0.50 \text{H}_2\text{O} \cdot 0.06 \text{TFA}$ : C, 56.13%; H, 4.70%; N, 14.80%; S, 6.77%. Found: C, 56.18%; H, 4.78%; N, 14.72%; S, 6.81%.

**Example A9:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-phenyl-benzamide.




$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  11.19 (s, 1H), 10.20 (s, 1H), 8.29 (bs, 2H), 8.04 (d, 2H,  $J=8.7$  Hz), 7.82 (m, 4H), 7.62 (m, 1H), 7.41 (t, 2H,  $J=7.8$  Hz), 7.28 (t, 2H,  $J=7.9$  Hz), 7.17 (t, 1H,  $J=7.8$  Hz).

FABMS ( $M+H^+$ ): 451; ( $M-H^-$ ): 449.

- 5     Anal. Calcd. for  $\text{C}_{23}\text{H}_{16}\text{F}_2\text{N}_4\text{O}_2\text{S}$ : C, 61.32%; H, 3.58%; N, 12.44%; S, 7.12%. Found: C, 61.30%; H, 3.66%; N, 12.35%; S, 7.07%.

**Example A10** : 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2S-ylamino]-N-(1-Acetyl-pyrrolidin-2S-ylmethyl)-benzamide.




- 10             The title compound was prepared by acylation of the compound of Example A8 with acetic anhydride in  $\text{CH}_2\text{Cl}_2$ , followed by preparative HPLC purification.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.97–7.71 ( $\mu$ , 4H), 7.47 (m, 1H), 7.08 (t, 2H,  $J=7.8$  Hz), 3.74 (m, 1H), 3.68–3.40 (m, 4H), 2.95–2.63 (m, 2H), 2.21 (s, 1H), 2.09 (s, 3H), 1.88 (m, 1H).

HRFABMS: Calcd. For  $\text{C}_{24}\text{H}_{23}\text{F}_2\text{N}_5\text{O}_3\text{S}$  ( $M+H^+$ ): 500.1568. Found: 500.1514.

- 15     **Example A11** : 4-[4-Amino-5-(2,6-difluoro-benzoyl)-2-ylamino]-N-(1-ethyl-pyrrolidin-2RS-ylmethyl)-benzamide.




- $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.79 (d, 2H,  $J=8.7$  Hz), 7.40 (d, 2H,  $J=8.7$  Hz), 7.33 (m, 1H), 6.94 (t, 2H,  $J=8.7$  Hz), 3.69 (m, 1H), 3.37–3.17 (m, 2H), 2.85 (m, 1H), 2.72 (m, 1H), 2.26 (m, 2H), 1.94–1.58 (m, 4H), 1.16 (t, 3H,  $J=7.1$  Hz).

FABMS ( $M+H^+$ ): 486; ( $M-H^-$ ): 484.

Anal. Calcd. for  $C_{24}H_{25}F_2N_5O_2S \cdot 0.2 H_2O \cdot 0.1 EtOAc$ : C, 58.85%; H, 5.19%; N, 14.06%; S, 6.44%.

Found: C, 58.84%; H, 5.30%; N, 14.09%; S, 6.38%.

**Example A12** : 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-2-methyl-propyl)-benzamide.



5

The starting material 2,N<sup>2</sup>,N<sup>2</sup>-trimethyl-propane-1,2-diamine was prepared according to the literature procedure (Yang et al, *Eur J. Med. Chem.* Vol 31, pp. 231-239, (1996)).


Coupling as in Step 3 for Method A and column chromatography with 15% MeOH/CHCl<sub>3</sub> provided a yellow solid in 25% yield.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 8.19 (bs, 1H), 7.86 (d, 2H, J=8.7 Hz), 7.66 (d, 2H, J=8.6 Hz), 7.61-7.48 (m, 1H), 7.21 (t, 2H, J=7.9 Hz), 2.30 (bs, 2H), 1.02 (s, 6H).

HRFABMS Calcd. for  $C_{23}H_{26}F_2N_5O_2S (M+H^+)$ : 474.1775. Found: 474.1784.

Anal. Calcd for  $C_{23}H_{25}F_2N_5O_2S \cdot 0.45 CHCl_3 \cdot 0.15 DMSO$ : C, 52.93%; H, 4.93%; N, 12.99%; S, 6.84%. Found: C, 52.89%; H, 5.32%; N, 12.61%; S, 6.89%.

**Example A13**: 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-benzyl-piperidin-4-yl)-benzamide.



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.05 (br, 1H), 8.32-8.10 (m, 3H), 7.86 (d, 2H, J=8.7 Hz), 7.65 (d, 2H, J=8.7 Hz), 7.57 (m, 1H), 7.39-7.18 (m, 7H), 3.78 (m, 1H), 3.49 (s, 2H), 2.85 (m, 2H), 2.13-1.98 (m, 2H), 1.80 (m, 2H), 1.68-1.53 (m, 2H).


20

FABMS (M+H<sup>+</sup>): 548; (M-H<sup>-</sup>): 546.

Anal. Calcd. for  $C_{29}H_{27}F_2N_5O_2S \cdot 0.25 H_2O \cdot 0.25 TFA$ : C, 61.02%; H, 4.82%; N, 12.06%; S, 5.52%. Found: C, 61.03%; H, 5.00%; N, 12.23%; S, 5.62%.

**Example A14**: 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-[2-(4-hydroxy-phenyl)-ethyl]-benzamide.

25




$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.04 (s, 1H), 9.12 (s, 1H), 8.40 (t, 1H,  $J=5.6$  Hz), 8.21 (bs, 2H), 7.81 (d, 2H,  $J=8.7$  Hz), 7.67 (d, 2H,  $J=8.7$  Hz), 7.55 (m, 1H), 7.21 (t, 2H,  $J=7.8$  Hz), 7.02 (d, 2H,  $J=8.4$  Hz), 6.68 (d, 2H,  $J=8.5$  Hz), 3.47-3.34 (m, 2H), 2.73 (t, 2H,  $J=7.4$  Hz).

5 FABMS ( $M+H^+$ ): 495, ( $M-H^-$ ): 493.

Anal. Calcd. for  $C_{25}H_{20}F_2N_4O_3S \cdot 0.3 H_2O \cdot 0.04 EtOAc$ : C, 60.02%; H, 4.19%; N, 11.13%; S, 6.37%. Found: C, 60.04%; H, 4.22%; N, 11.27%; S, 6.24%.

**Example A15:** 4-[4-Amino-5-(2,6-difluorobenzoyl)-thiazol-2-ylamino]-N-(2-piperidin-1-ylethyl)-benzamide.




$^1\text{H}$  NMR ( $CD_3OD$ ):  $\delta$  8.72 (d, 2H,  $J=8.7$  Hz), 7.73 (d, 2H,  $J=8.7$  Hz), 7.47 (m, 1H), 7.05 (t, 2H,  $J=7.8$  Hz), 3.54 (t, 2H,  $J=6.8$  Hz), 2.68-2.49 (m, 6H), 1.71-1.58 (m, 4H), 1.56-1.44 (m, 2H).

FABMS( $M+H^+$ ): 486; ( $M-H^-$ ): 484.

15 Anal. Calcd. for  $C_{24}H_{25}F_2N_5O_2S \cdot 0.4 H_2O \cdot 0.5 EtOAc$ : C, 58.17%; H, 5.60%; N, 13.05%; S, 5.97%. Found: C, 58.34%; H, 5.46%; N, 13.20%; S, 5.78%.

**Example A16:** 4-[4-Amino-5-(2,6-difluorobenzoyl)-thiazol-2-ylamino]-N-(4-dimethylaminophenyl)-benzamide.




$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.05 (s, 1H), 9.88 (s, 1H), 8.21 (bs, 2H), 7.94 (d, 2H,  $J=8.7$  Hz), 7.70 (d, 2H,  $J=8.7$  Hz), 7.55 (m, 3H), 7.21 (t, 2H,  $J=7.9$  Hz), 6.86 (d, 2H,  $J=9.1$  Hz), 2.88 (s, 6H).

FABMS ( $M+H^+$ ): 494; ( $M+Na^+$ ): 516.

Anal. Calcd. for  $C_{25}H_{22}F_2N_5O_2S \cdot 0.1 H_2O \cdot 0.05 EtOAc$ : C, 60.56%; H, 4.36%; N, 14.01%; S, 6.42%. Found: C, 60.57%; H, 4.54%; N, 14.00%; S, 6.64%.

**Example A17:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2, 3-dihydro-benzo[1, 4]dioxin-6-yl)-benzamide.




$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.10 (s, 1H), 9.95 (s, 1H), 8.21 (bs, 2H), 7.92 (d, 2H,  $J=8.8$  Hz), 7.72 (d, 2H,  $J=8.8$  Hz), 7.60 (m, 1H), 7.37 (d, 1H,  $J=2.4$  Hz), 7.26-7.13 (m, 3H), 6.81 (d, 2H,  $J=8.7$  Hz), 4.25-4.18 (m, 4H).

FABMS ( $M+H^+$ ): 509; ( $M-H^-$ ): 507.


Anal. Calcd. for  $C_{25}H_{18}F_2N_4O_4S \cdot 0.07 H_2O \cdot 0.05 EtOAc$ : C, 58.90%; H, 3.59%; N, 10.99%; S, 6.29%. Found: C, 58.90%; H, 3.70%; N, 10.88%; S, 6.21%.

**Example A18:** 4-[4-Amino-5-[1-(2,6-difluoro-phenyl)-methanoyl]-thiazol-2-ylamino]-N-(2-methyl-2-methylamino-propyl)-benzamide Acetic Acid Salt.



The starting material was prepared as follows:

2, $N^2$ -Dimethyl-propane-1,2-diamine.



20

Crude 2-methyl-2-methylamino-propionitrile (2.00 g, 20.4 mmol; Gabriel, *Chem. Ber.*, 47, 2922-2925 (1914) and  $^1\text{H}$  NMR matched Stork et al., *J. Am. Chem. Soc.*; 96; 1974; 5787-5791 (1974)) was added to a suspension of  $LiAlH_4$  (1.55 g, 20.38 mmol) in ether (40 mL) at  $0^\circ\text{C}$ . The resultant mixture was heated at reflux for 3 hours, then cooled to  $0^\circ\text{C}$ , and quenched with  $H_2O$  (4

mL) and 2N NaOH (3 mL). After a few minutes of stirring, the granular white solids were filtered off. The filtrate was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated *in vacuo* to afford 1.5 g of a colorless oil in 72% yield, which was used without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 2.47 (2H, s), 2.22 (3H, s), 0.94 (6H, s).


5 The title compound was prepared in a manner similar to Step 2 in Method A from 2,N<sup>2</sup>-dimethyl-propane-1,2-diamine, and purified via column chromatography with 2% HOAc/10%MeOH/ CHCl<sub>3</sub> as eluant.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 8.28 (bs, 1H), 8.15 (bs, 1H), 7.87 (d, 2H, J = 8.7 Hz), 7.62 (d, 2H, J = 8.5 Hz), 7.60-7.48 (m, 1H), 7.20 (t, 2H, J = 7.9 Hz), 3.32 (d, 2H, J = 6.0 Hz), 2.28 (s, 3H), 1.02 (s, 6H).

HRFABMS Calcd. for C<sub>22</sub>H<sub>24</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S (M+H<sup>+</sup>): 460.1619. Found: 460.1612.

Anal. Calcd. for C<sub>22</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•2 HOAc•0.4 CHCl<sub>3</sub>: C, 50.54%; H, 5.04%; N, 11.16%; S, 5.11%. Found: C, 50.31%; H, 5.20%; N, 10.88%; S, 5.03%.

**Example A19:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(5-methoxy-2-methyl-phenyl)-benzamide




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.11 (br, 1H), 9.70 (s, 1H), 8.21 (bs, 2H), 7.98 (d, 2H, J=8.8 Hz), 7.73 (d, 2H, J=8.8 Hz), 7.56 (m, 1H), 7.29-7.13 (m, 3H), 7.01 (d, 1H, J=2.7 Hz), 6.75 (dd, 1H, J=2.7, 5.6 Hz), 3.73 (s, 3H), 2.17 (s, 3H).

20 FABMS (M+H<sup>+</sup>): 495; (M-H): 493.

Anal. Calcd. for C<sub>25</sub>H<sub>20</sub>F<sub>2</sub>N<sub>4</sub>O<sub>3</sub>S•0.19 H<sub>2</sub>O•0.15 EtOAc: C, 60.15%; H, 4.26%; N, 10.96%; S, 6.27%. Found: C, 60.03%; H, 4.21%; N, 10.91%; S, 6.33.

**Example A20:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-1RS-methyl-ethyl)-benzamide Hydrochloride Salt




25

$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.02 (br, 1H), 8.18 (bs, 2H), 8.03 (d, 1H,  $J=8.1$  Hz), 7.82 (d, 2H,  $J=8.8$  Hz), 7.68 (d, 2H,  $J=8.8$  Hz), 7.55 (m, 1H), 7.26-7.14 (t, 2H,  $J=8.7$  Hz), 4.16 (m, 1H), 2.42 (m, 1H), 2.28 (m, 1H), 2.21 (s, 6H), 1.14 (d, 3H,  $J=4.5$  Hz).

FABMS ( $M+H^+$ ): 460; ( $M-H^-$ ): 458.

- 5     Anal. Calcd. for  $C_{22}H_{23}F_2N_5O_2S \cdot 0.80 \text{ HCl} \cdot 0.4 \text{ EtOAc}$ : C, 54.10%; H, 5.19%; N, 13.37%; S, 6.12%. Found: C, 53.97%; H, 5.10%; N, 13.49%; S, 6.16%.

**Example A21:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(4-morpholin-4-yl-phenyl)-benzamide.




- 10      $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.10 (s, 1H), 9.94 (s, 1H), 8.21 (bs, 2H), 7.95 (d, 2H,  $J=8.7$  Hz), 7.71 (d, 2H,  $J=8.7$  Hz), 7.56 (m, 3H), 7.22 (t, 2H,  $J=7.8$  Hz), 6.94 (d, 2H,  $J=9.1$  Hz), 3.84 (t, 2H,  $J=4.7$  Hz), 3.08 (t, 2H,  $J=4.7$  Hz).

FABMS ( $M+H^+$ ): 536; ( $M-H^-$ ): 534.

- 15     Anal. Calcd. for  $C_{27}H_{23}F_2N_5O_3S \cdot 1.25 \text{ H}_2\text{O}$ : C, 58.11%; H, 4.61%; N, 12.55%; S, 5.75%. Found: C, 58.11%; H, 4.48%; N, 12.58%; S, 5.79%.

**Example A22:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-indan-2-yl-benzamide.




- 20      $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.05 (s, 1H), 8.54 (d, 1H,  $J=7.0$  Hz), 8.20 (bs, 2H), 7.87 (d, 2H,  $J=8.7$  Hz), 7.64 (d, 2H,  $J=8.7$  Hz), 7.55 (m, 1H), 7.28-7.12 (m, 6H), 4.68 (m, 1H), 3.24-3.19 (m, 2H), 3.01-2.89 (m, 2H).

FABMS( $M+H^+$ ): 548; ( $M-H^-$ ): 546.

Anal. Calcd. for  $C_{26}H_{20}F_2N_4O_2S \cdot 1.00 \text{ CHCl}_3$ : C, 53.17%; H, 3.47%; N, 9.19%; S, 5.26%. Found: C, 52.95%; H, 3.38%; N, 9.45%; S, 5.54%.



**Example A23:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-diisopropylamino-ethyl)-benzamide.




<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 8.83 (d, 2H, J=8.9 Hz), 7.76 (d, 2H, J=8.9 Hz), 7.48 (m, 1H), 7.08 (t, 2H, J=7.1 Hz), 3.42 (t, 2H, J=7.1), 3.18 (m, 2H), 2.85 (m, 2H), 1.12 (d, 6H, J=6.5 Hz).

FABMS (M+H<sup>+</sup>): 515; (M-H<sup>-</sup>): 513.

Anal. Calcd. For C<sub>25</sub>H<sub>29</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•0.12 H<sub>2</sub>O: C, 59.60%; H, 5.85%; N, 13.90%; S, 6.37%. Found: C, 59.60%; H, 5.82%; N, 13.80%; S, 6.34%.

**Example A24:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-[4-(2-hydroxy-ethyl)-phenyl]-benzamide.




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.12 (s, 1H), 10.15 (s, 1H), 8.21 (bs, 2H), 7.94 (d, 2H, J=8.7 Hz), 7.84 (d, 2H, J=8.7 Hz), 7.64 (d, 2H, J=8.7 Hz), 7.55 (m, 1H), 7.30-7.12 (m, 4H), 4.60 (t, 1H, J=5.2 Hz), 3.40 (dd, 2H, J=5.2, 7.1 Hz), 2.71 (t, 2H, J=7.1 Hz).

FABMS (M+H<sup>+</sup>): 495; (M-H<sup>-</sup>): 493.

Anal. Calcd. For C<sub>25</sub>H<sub>20</sub>F<sub>2</sub>N<sub>4</sub>O<sub>3</sub>S•0.25 H<sub>2</sub>O: C, 60.17%; H, 4.14%; N, 11.23%; S, 6.43%. Found: C, 60.26%; H, 4.22%; N, 11.14%; S, 6.36%.


**Example A25:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-[4-(acetyl-methyl-amino)-phenyl]-benzamide.



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.12 (s, 1H), 10.25 (s, 1H), 8.21 (bs, 2H), 7.96 (d, 2H, J=8.8 Hz), 7.82 (d, 2H, J=8.7 Hz), 7.75 (d, 2H, J=8.9 Hz), 7.56 (m, 1H), 7.31 (d, 2H, J=8.8 Hz), 7.21 (t, 2H, J=7.8 Hz), 3.16 (s, 3H), 1.88 (s, 3H).

LCMS (M+H<sup>+</sup>): 522

- 5 **Example A26:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-thiophen-2-yl-ethyl)-benzamide.




- <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 11.05 (s, 1H), 8.54 (br, 1H), 8.21 (bs, 2H), 7.84 (d, 2H, J=8.8 Hz), 7.67 (d, 2H, J=8.8 Hz), 7.55 (m, 1H), 7.34 (dd, 1H, J=1.2, 5.4 Hz), 7.22 (t, 2H, J=7.8 Hz), 6.98-6.90 (m, 2H), 3.21 (m, 2H), 3.08 (t, 2H, J=7.2 Hz).

FABMS (M+H<sup>+</sup>): 485; (M-H<sup>-</sup>): 483.

Anal. Calcd. For C<sub>23</sub>H<sub>18</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub>S<sub>2</sub>•0.18 H<sub>2</sub>O: C, 59.63%; H, 3.79%; N, 11.49%; S, 13.15%. Found: C, 59.80%; H, 3.96%; N, 11.39%; S, 13.09%.


- 15 **Example A27:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(pyridin-3-yl)-benzamide



<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 8.90 (s, 1H), 8.28 (m, 2H), 8.04 (d, 2H, J=8.8Hz), 7.82 (d, 2H, J=8.8 Hz), 7.44 (m, 2H), 7.75 (d, 2H, J=8.9 Hz), 7.08 (t, 2H, J=8.7 Hz).

- Anal. Calcd. For C<sub>22</sub>H<sub>15</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•0.5 H<sub>2</sub>O: C, 57.38%; H, 3.50%; N, 15.21%; S, 6.96%. Found: C, 57.49%; H, 3.75%; N, 15.09%; S, 6.82%.

**Example A28:** 4-[[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(4-cyanomethyl-benzyl)-benzamide.




<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.10 (br, 1H), 10.20 (s, 1H), 8.21 (bs, 2H), 7.98 (d, 2H, J=8.8 Hz), 7.86-7.71 (m, 4H), 7.57 (m, 1H), 7.35 (d, 2H, J=8.6 Hz), 7.22 (t, 2H, J=7.9 Hz), 3.18 (d, 2H, J=5.2 Hz), 2.09 (s, 2H).

5 FABMS (M+H<sup>+</sup>): 490; (M-H): 488.

Anal. Calcd. For C<sub>23</sub>H<sub>18</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub>S<sub>2</sub>•0.20 H<sub>2</sub>O•0.20 EtOAc: C, 60.67%; H, 3.75%; N, 13.71%; S, 6.28%. Found: C, 60.50%; H, 3.89%; N, 13.72%; S, 6.29%.

**Example A29:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-pyridin-4-yl-ethyl)-benzamide.



10


<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.08 (br, 1H), 8.52-8.45 (m, 3H), 8.24 (bs, 2H), 7.84 (d, 2H, J=8.7 Hz), 7.69 (d, 2H, J=8.7 Hz), 7.57 (m, 1H), 7.31-7.18 (m, 4H), 3.55 (m, 2H), 2.92 (t, 2H, J=7.1 Hz).

FABMS (M+H<sup>+</sup>): 480; (M-H): 478.

Anal. Calcd. For C<sub>23</sub>H<sub>18</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub>S<sub>2</sub>•0.70 H<sub>2</sub>O•0.14 EtOAc: C, 58.70%; H, 4.37%; N, 13.94%; S, 6.38%. Found: C, 58.77%; H, 4.15%; N, 14.10%, S, 6.29%.

15

**Example A30:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-pyrrolidin-3R-yl)-benzamide.



The starting material, 1-methyl-pyrrolidin-3R-ylamine trifluoroacetic acid salt, was  
20 prepared as follows:

A mixture of 3R-(*t*-Boc-amino)pyrrolidine (0.38 g, 2.0 mmol; TCT) and paraformaldehyde (0.06 g, 2.2 mmol) in MeOH (15 ml) was hydrogenated at 60 psi with 10% Pd/C for 16 hours. Catalyst was filtered off and the filtrate concentrated. A solution of the resultant residue in ethyl acetate was washed with sat. NaHCO<sub>3</sub>, dried over MgSO<sub>4</sub>, filtered, and concentrated to give 0.27 g of (1-methyl-pyrrolidin-3R-yl)-carbamic acid *tert*-butyl ester as a clear oil in 67% yield, which was used without further purification.

<sup>1</sup>H NMR: δ 4.80 (m, 1H), 4.10 (m, 1H), 2.74 (m, 1H), 2.50 (m, 2H), 2.30 (s, 3H), 2.22 (m, 1H), 1.40 (s, 9H).

(1-Methyl-pyrrolidin-3R-yl)-carbamic acid *tert*-butyl ester (0.20 g, 1.0 mmol) in 30% TFA/CH<sub>2</sub>Cl<sub>2</sub> (10 ml) was stirred at room temperature for 30 minutes. The solvent was removed and the residue was dried under vacuum to give 1-methyl-pyrrolidin-3R-ylamine trifluoroacetic acid salt as a clear oil in 100% yield which was used directly in the next step.


The title compound was prepared in a manner similar to Step 3 in Method A.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.05 (br, 1H), 8.35 (d, 1H, J=7.0 Hz), 8.20 (br, 2H), 7.86 (d, 2H, J=8.7 Hz), 8.64 (d, 2H, J=8.7 Hz), 7.56 (m, 1H), 7.19 (t, 2H, J=7.9 Hz), 4.38 (m, 1H), 2.71 (m, 1H), 2.58 (m, 1H), 2.48-2.37 (m, 2H), 2.27 (s, 3H), 2.15 (m, 1H), 1.76 (m, 1H).

FABMS (M+H<sup>+</sup>): 458; (M-H<sup>-</sup>): 456.

Anal. Calcd. For C<sub>22</sub>H<sub>21</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S: C, 57.76%; H, 4.63%; N, 15.31%; S, 7.01%. Found: C, 57.76%; H, 4.79%; N, 15.05%; S, 6.90%.

**Example A31:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-pyrrolidin-2S-yl-methyl)-benzamide Dihydrochloride.



4-[4-Amino-5-(2, 6-difluoro-benzoyl-thiazol)-2-ylamino]-N-(1-methyl-pyrrolidin-2S-yl-methyl)-benzamide

To a solution of (1-methyl-pyrrolidin-2S-yl)-methylamine (3.04 g, 26.6 mmol; Sassaman et al., *Bioorg. Med. Chem.*, 6, 1759-1766 (1998)) in DMF (70 ml) was added 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**; 9.08 g, 24.2 mmol). The resultant yellow solution was cooled to 4.7°C, PyBOP (13.2 g, 25.4 mmol) and diisopropylethylamine (10.5 ml, 60.5 mmol) were sequentially added (slight exotherm

observed after each), and allowed to warm to room temperature overnight. Most of the DMF was removed in vacuo, and the resultant liquid diluted with ethyl acetate (1.2 L). The solution was washed with 5% aq Na<sub>2</sub>CO<sub>3</sub> (4 × 350 ml) and sat. NaCl (250 ml), dried over MgSO<sub>4</sub>, and concentrated to give a yellow solid, typically used without any further purification. An

5 analytically pure sample was available via preparative HPLC.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.04 (br, 1H), 8.34-8.05 (m, 3H), 7.84 (d, 2H, J=8.8 Hz), 7.66 (d, 2H, J=8.8 Hz), 7.56 (m, 1H), 7.22 (t, 2H, J=7.8 Hz), 3.46 (m, 1H), 3.15 (m, 1H), 2.94 (m, 1H), 2.38 (m, 1H), 2.35 (s, 3H), 2.17 (m, 1H), 1.84 (m, 1H), 1.72-1.51 (m, 3H).

FABMS (M+H<sup>+</sup>): 472. (M-H<sup>-</sup>): 470.

10 Anal. Calcd. For C<sub>23</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S: C, 58.00%; H, 5.13%; N, 14.16%; S, 6.48%. Found: C, 57.98%; H, 5.13%; N, 14.03%; S, 6.32%.


The title compound was prepared as follows. Through a solution of crude 4-[4-amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-pyrrolidin-2S-yl-methyl)-benzamide (theoretical 24.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (170 ml) at 0°C was passed HCl gas for a few minutes. The resultant suspension was allowed to warm and stir for 2 hours. The yellow solid was filtered off, rinsed with CH<sub>2</sub>Cl<sub>2</sub> (3 × 100 ml), and dried under vacuum to obtain 10.54 g in 80% yield from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**).

[α]<sub>D</sub> = -20.2° (MeOH).

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 7.93 (d, 2H, J=8.5 Hz), 7.72 (d, 2H, J=8.5 Hz), 7.54 (m, 1H), 7.20 (m, 2H), 3.82-3.43 (m, 4H), 3.08 (m, 1H), 2.48 (s, 3H), 2.11 (m, 1H), 1.96 (m, 1H).

Anal. Calcd for C<sub>23</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•2.25 HCl: C, 49.90; H, 4.60; N, 12.65; S, 5.79. Found: C, 49.75; H, 4.88; N, 12.50; S, 5.77.

**Example A32:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-pyrrolidin-3S-yl)-benzamide.




The starting material, 1-methyl-pyrrolidin-3S-ylamine trifluoroacetic acid salt, was prepared in a route similar to that for Example A30 from 3R-(t-Boc-amino)pyrrolidine.

The title compound was prepared in a manner similar to Step 3 in Method A.

Spectral data are identical to Example A30.

Anal. Calcd. For  $C_{22}H_{21}F_2N_5O_2S \cdot 0.90 H_2O \cdot 0.08 EtOAc \cdot 0.09 CHCl_3$ : C, 54.76%; H, 4.83%; N, 14.25%; S, 6.52%. Found: C, 54.69%; H, 4.68%; N, 14.12%; S, 6.41%.

**Example A33:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(piperidin-2R-ylmethyl)-benzamide Trifluoroacetate Salt.



The intermediate 2-({[1-(4-{4-Amino-5-[1-(2,6-difluoro-phenyl)-methanoyl]-thiazol-2-ylamino}-phenyl)-methanoyl]-amino}-methyl)-piperidine-1-carboxylic acid *tert*-butyl ester was prepared in a manner similar to Step 3 in Method A from 2-aminomethyl-piperidine-1-carboxylic acid *tert*-butyl ester.


The title compound was prepared by treating the above intermediate with 30% TFA/ $CH_2Cl_2$  for 30 minutes, followed by HPLC purification.

$^1H$  NMR ( $DMSO-d_6$ ):  $\delta$  11.04 (br, 1H), 8.15 (bs, 2H), 7.82 (m, 2H), 7.66 (d, 2H,  $J=8.7$  Hz), 7.54 (m, 1H), 7.43 (d, 2H,  $J=8.7$  Hz), 7.21 (t, 2H,  $J=7.8$  Hz), 3.68-3.21 (m, 5H), 1.82-1.25 (m, 6H).

FABMS ( $M+H^+$ ): 472; ( $M-H^-$ ): 470.

Anal. Calcd. For  $C_{23}H_{23}F_2N_5O_2S \cdot 0.60 H_2O \cdot 1.32 TFA$ : C, 48.66%; H, 4.06%; N, 11.07%; S, 4.83%. Found: C, 48.55%; H, 4.03%; N, 11.24%; S, 4.83%.

**Example A34:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-pyrrolidin-2R-ylmethyl)-benzamide




The starting material (1-methyl-pyrrolidin-2R-yl)-methylamine was prepared in analogous fashion to that for the (S) isomer (Sassaman et al. *Bioorg. Med. Chem.* Vol. 6, pp.1759-1766, (1998)).

The title compound was prepared in a manner similar to Step 3 in Method A.

Spectra data are identical to Example 31.

Anal. Calcd. For  $C_{23}H_{23}F_2N_5O_2S \cdot 0.52 H_2O \cdot 0.18 EtOAc$ : C, 57.35%; H, 5.17%; N, 14.10%; S, 6.45%. Found: C, 57.40%; H, 4.98%; N, 14.04%; S, 6.31%.

**Example A35:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-pyrrolidin-1-yl-ethyl)-benzamide.



5


$^1H$  NMR ( $CD_3OD$ ):  $\delta$  7.88 (s, 2H,  $J=8.7$  Hz), 7.74 (d, 2H,  $J=8.7$  Hz), 7.49 (m, 1H), 7.10 (t, 2H,  $J=7.8$  Hz), 3.59 (t, 2H,  $J=6.7$  Hz), 2.82 (t, 2H,  $J=6.7$  Hz), 2.73 (bs, 4H), 1.88 (bs, 4H).

FABMS ( $M+H^+$ ): 472; ( $M-H$ ): 470.

Anal. Calcd. For  $C_{23}H_{23}F_2N_5O_2S \cdot 0.60 H_2O \cdot 0.20 EtOAc \cdot 0.10 CHCl_3$ : C, 56.08%; H, 5.10%; N, 13.68%; S, 6.26%. Found: C, 56.11%; H, 4.98%; N, 13.63%; S, 6.34%.

10

**Example A36:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-1S-methyl-ethyl)-N-methyl-benzamide.



The starting material, (S)- $N^1, N^1, N^2$ -trimethyl-propane-1,2-diamine, was prepared as follows:

15

To a solution of N-(*tert*-butoxycarbonyl)-L-alanine (1.5 g, 8.0 mmol) in DMF (100 ml), were added PyBop (5.0g, 9.6 mmol), DIEA (5.0 ml, 28.0 mmol), and dimethyl-amine hydrochloride salt (0.98 g, 12 mmol). The mixture was stirred at room temperature for 3 hours. The solvent was evaporated under reduced pressure and a solution of the resultant residue in ethyl acetate was washed with sat.  $Na_2CO_3$  and brine, dried over  $MgSO_4$ , filtered, and concentrated to a syrup. Column chromatography (hexane/ethyl acetate=1/1) afforded 1.50 g of (1S-dimethylcarbamoyl-ethyl)-carbamic acid *tert*-butyl ester in 87% yield, which was used without further purification.

20

To a solution of 1.0 M  $LiAlH_4$  (5.6 mmol, 5.6 ml) in 50% THF/ $Et_2O$  (50 ml), (1S-dimethylcarbamoyl-ethyl)-carbamic acid *tert*-butyl ester (0.60 g, 2.8 mmol) in THF (12 ml)

25

was added dropwise over 30 minutes. The reaction mixture was stirred at room temperature for 60 minutes and then refluxed for 18 hours. The resulting opaque milky-white solution was cooled to room temperature and quenched with careful addition of sat.  $\text{Na}_2\text{SO}_4$  (1.0 ml). The mixture was filtered and the filtrate was carefully concentrated below  $50^\circ\text{C}$  under reduced pressure to give the desired amine as a clear oil in 56 % yield, which was used without further purification.


The title compound was prepared in a manner similar to Step 3 in Method A.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.74 (d, 2H,  $J=8.5$  Hz), 7.58-7.39 (m, 3H), 7.08 (t, 3H,  $J=7.8$  Hz), 2.91 (m, 3H), 2.80-2.52 (m, 2H), 2.38 (bs, 3H), 2.28 (m, 1H), 2.07 (bs, 3H), 1.23 (d, 3H,  $J=6.7$  Hz).

10 FABMS ( $\text{M}+\text{H}^+$ ): 474; ( $\text{M}-\text{H}^-$ ): 472.

Anal. Calcd. For  $\text{C}_{23}\text{H}_{25}\text{F}_2\text{N}_5\text{O}_2\text{S}\cdot 0.40 \text{ H}_2\text{O}\cdot 0.20 \text{ EtOAc}\cdot 0.50 \text{ CHCl}_3$ : C, 52.30%; H, 5.04%; N, 12.55%; S, 5.75%. Found: C, 52.24%; H, 5.26%; N, 12.53%; S, 5.73%.

**Example A37:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-[2-dimethylamino-1R-methyl-ethyl]-N-methyl-benzamide.



15


The starting material (*R*)- $N^1$ ,  $N^1$ ,  $N^2$ -trimethyl-propane-1,2-diamine was prepared with a route similar to that for Example 36 from 2*R*-*tert*-butoxycarbonylamino-propionic acid.

The title compound was prepared in a manner similar to Step 3 in Method A.

Spectral data are identical to Example A36.

20 Anal. Calcd. For  $\text{C}_{23}\text{H}_{25}\text{F}_2\text{N}_5\text{O}_2\text{S}\cdot 0.40 \text{ H}_2\text{O}\cdot 0.28 \text{ EtOAc}\cdot 0.39 \text{ CHCl}_3$ : C, 53.33%; H, 5.19%; N, 12.69%; S, 5.81%. Found: C, 53.32%; H, 5.19%; N, 12.69%; S, 5.81%.

**Example A38:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-ethyl-pyrrolidin-2S-yl-methyl)-benzamide.






Starting material, S-(-)-2-aminomethyl-1-ethyl-pyrrolidine, is available from TCT.

The title compound was prepared in a manner similar to Step 3 in Method A.

Spectra data are identical to Example A11.


Anal. Calcd. For  $C_{24}H_{25}F_2N_5O_2S \cdot 0.10 H_2O \cdot 0.25 EtOAc \cdot 0.17 CHCl_3$ : C, 57.07%; H, 5.21%; N, 13.22%; S, 6.05%. Found: C, 56.93%; H, 5.21%; N, 13.20%; S, 6.05%.

**Example A39:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino}-N-2-(cyclopropylmethyl-methyl-amino)-2-methyl-propyl]-benzamide



The starting materials were prepared as follows:


Step 1. 2-(Cyclopropylmethyl-methyl-amino)-2-methyl-propionitrile.



To a solution of 0.262 M of cyclopropylmethyl-methyl-amine in ether (39.0 mL, 10.1 mmol; Grotjahn, *J. Het. Chem.*, 20, 1031-1036 (1983)) at ambient temperature was added 2-hydroxy-2-methyl-propionitrile (922  $\mu$ L, 10.1 mmol) and 10% HCl (20 mL). After 12 hours, the mixture was brought to pH8 with 10% NaOH and extracted with ether (2  $\times$  25 mL). The combined organic layers were dried over  $Na_2SO_4$ , filtered, and carefully concentrated under aspirator pressure below 50°C to provide 1.49 g of a volatile colorless oil in 97% yield, which was used without further purification.

$^1H$  NMR:  $\delta$  2.30 (s, 3H), 1.35 (s, 6H).

Step 2. *N*<sup>2</sup>-Cyclopropylmethyl-2,*N*<sup>2</sup>-dimethyl-propane-1,2-diamine.



To a suspension of  $LiAlH_4$  (749 mg, 19.72 mmol) in ether (40 mL) at 0°C was added crude 2-(cyclopropylmethyl-methyl-amino)-2-methyl-propionitrile (1.50 g, 9.86 mmol). The resultant mixture was heated at reflux for 3 hours, then cooled to 0°C, and carefully quenched with  $H_2O$  (4 mL) and 2N NaOH (3 mL). After a few minutes of stirring, the granular white solids were filtered

off. The filtrate was dried over  $\text{Na}_2\text{SO}_4$ , filtered, and carefully concentrated under aspirator pressure below  $50^\circ\text{C}$  to afford 995 mg of a volatile colorless oil in 65% yield, which was used without further purification.

$^1\text{H}$  NMR:  $\delta$  2.45 (s, 2H), 2.19 (s, 3H), 2.12 (d, 2H,  $J=6.4$  Hz), 0.88 (s, 6H).


5

The title compound was prepared in a manner similar to Step 3 in Method A.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  7.80 (d, 2H,  $J=8.8$  Hz), 7.64 (d, 2H,  $J=8.7$  Hz), 7.60-7.50 (m, 1H), 7.20 (dd, 2H,  $J=7.7, 8.2$  Hz), 2.50 (s, 3H), 1.00 (s, 6H), 0.88-0.78 (m, 1H), 0.48-0.40 (m, 2H), 0.09 (d, 2H,  $J=4.2$  Hz).


10 Anal. Calcd. for  $\text{C}_{26}\text{H}_{29}\text{F}_2\text{N}_5\text{O}_2\text{S} \cdot 0.4 \text{ H}_2\text{O}$ : C, 59.96%; H, 5.77%; N, 13.45%; S, 6.16%. Found: C, 59.66%; H, 5.82%; N, 13.40%; S, 6.06%.

**Example A40:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-1R-methyl-ethyl)-benzamide Dihydrochloride



15 The starting material, (R)-N, N-dimethyl-propane-1, 2-diamine dihydrochloride salt was prepared as follows:


Step 1: (1R-Dimethylcarbamoyl-ethyl)-carbamic Acid Benzyl Ester



20 To a solution of N-benzyloxycarbonyl-D-alanine (11.1 g, 50.0 mmol), dimethylamine hydrochloride salt (8.10 g, 100 mmol) and DIEA (13.0 g, 100 mmol) in  $\text{CH}_3\text{CN}$  (200 ml) at  $0^\circ\text{C}$  was added dicyclohexyl-carbodiimide (10.3 g, 50.0 mmol). The mixture was stirred for 2 hours and filtered. The filtrate was concentrated and a solution of the resultant residue in ethyl acetate was washed with 0.1 N HCl, 0.1 N NaOH, dried over  $\text{MgSO}_4$ , filtered, and concentrated. Column  
25 chromatography (hexane/ethyl acetate=1/1) afforded 14.1 g of desired product in 98% yield, which

displayed an  $^1\text{H}$  NMR that matched literature (Isogai et al, *J. Chem. Soc. Perkin Trans. 1*, 1405-1411 (1984)) and used without further purification.


Step 2: (2-Dimethylamino-1R-methyl-ethyl)-carbamic Acid Benzyl Ester



- 5 To a solution of (1R-dimethylcarbamoyl-ethyl)-carbamic acid benzyl ester (14.0 g, 56.0 mmol) in THF (200 ml), was added dropwise over 30 minutes 1.0 M borane/THF (9.7 mmol, 9.7 ml). The reaction mixture was stirred at room temperature for 18 hours then quenched with conc. HCl (5ml). Solvent was evaporated under reduced pressure. The residue was dissolved in water and extracted with ethyl ether. The aqueous layer was separated, neutralized with 1M NaOH and  
10 extracted with ethyl acetate three times. The organic layers were combined, dried over  $\text{MgSO}_4$ , filtered and concentrated to give 6.2 g of clear oil in 50% yield, which was used without further purification.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.34 (m, 5H), 5.12 (m, 2H), 3.72 (m, 1H), 2.32 (m, 1H), 2.24 (s, 6H), 2.16 (m, 1H), 1.20 (d, 3H,  $J=5.0$  Hz).


- 15 Step 3: (R)-N, N-Dimethyl-propane-1, 2-diamine Dihydrochloride Salt



- A solution of above intermediate (6.0 g, 25 mmol) in 50% MeOH/ $\text{H}_2\text{O}$  (250 ml), conc. HCl (2 ml), and 10% Pd/C (0.5 g) was shaken under hydrogen at 30 psi for 1 hour. The catalyst was filtered off and the filtrate was lyophilized to give 5.4 g of white hygroscopic amorphous solid in  
20 90% yield, which was used without further purification.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  3.82 (m, 1H), 3.46-3.28 (m, 2H), 2.90 (s, 3H), 1.38 (d, 3H,  $J=5.0$  Hz).

Step 4: 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-1R-methyl-ethyl)-benzamide



- 25 To a mixture of (R)-N,N-dimethylpropane-1,2-diamine dihydrochloride salt (8.42 g, 48.1 mmol) and 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**; 16.4 g, 43.7 mmol) in DMF (125 ml) at  $2.5^\circ\text{C}$  was added PyBOP (23.9 g, 45.9 mmol). Diisopropylethyl-amine

(26.6 ml, 153 mmol) was added dropwise over 12 minutes. The resultant bright yellow-orange mixture stirred at 0°C for 30 minutes, then allowed to warm to room temperature overnight. The DMF was removed under reduced pressure, and the resultant oil partitioned with ethyl acetate (600 ml) and 5% aq Na<sub>2</sub>CO<sub>3</sub> (300 ml). The organic layer was separated, washed with 5% aq Na<sub>2</sub>CO<sub>3</sub> (2 × 300 ml) and sat. NaCl (250 ml), dried over MgSO<sub>4</sub>, and concentrated to give a bright yellow solid, which was used without any further purification.

<sup>1</sup>H NMR spectral data are identical to that for Example A20.

HRFABMS Calcd. For C<sub>22</sub>H<sub>24</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S (M+H<sup>+</sup>): 460.1613. Found: 460.1628.

Anal. Calcd. For C<sub>23</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S: C, 58.00%; H, 5.13%; N, 14.16%; S, 6.48%. Found: C, 57.98%; H, 5.13%; N, 14.03%; S, 6.32%.


The title compound was prepared as follows. Through a solution of crude 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-pyrrolidin-2S-yl-methyl)-benzamide (theoretical 43.7 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (375 ml) at 0°C was passed HCl gas for a few minutes. The resultant suspension was allowed to warm and stir overnight. The yellow solid was filtered off, rinsed with CH<sub>2</sub>Cl<sub>2</sub> (3 × 150 ml), and dried under vacuum. The solid was suspended in CH<sub>2</sub>Cl<sub>2</sub> (150 ml) and MeOH (0.75 ml), stirred 24 hours, filtered off, rinsed with CH<sub>2</sub>Cl<sub>2</sub> (2 × 100 ml), and dried under vacuum to obtain 17.92 g of bright yellow solid in 77% yield from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.90 (d, 2H, J=8.8 Hz), 7.78 (d, 2H, J=8.8 Hz), 7.52 (m, 1H), 7.08 (t, 2H, J=7.9 Hz), 4.62 (m, 1H), 3.04 (s, 3H), 2.95 (s, 3H), 1.38 (d, 2H, J=6.8 Hz).

Anal. Calcd. For C<sub>22</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•1.88 HCl: C, 50.04%; H, 4.75%; N, 13.26%; S, 6.07%. Found: C, 50.04%; H, 5.93%; N, 13.21%; S, 6.05%.

Chiral HPLC: Retention time 13.9 min. with Chiralcel OD-R 250 x 4.6 mm column, 0 to 30% acetonitrile/0.5 M NaClO<sub>4</sub> gradient over 20 min, flow rate 0.8 mL/min.

**Example A41:** 4-[4-Amino-5-(2,6-difluoro-phenyl)-methanoyl]-thiazol-2-ylamino}-N-(2-dimethylamino-1S-methyl-ethyl)-benzamide.



The starting material (S)-N, N-dimethyl-propane-1, 2-diamine dihydrochloride salt was prepared in a route similar to that in Example A40 from 2S-benzyloxycarbonylamino-propionic acid.


The title compound was prepared in a manner similar to Step 3 in Method A.

5 Spectral data are identical to Example A40.

HRFABMS Calcd. for  $C_{22}H_{23}F_2N_5O_2SNa$  ( $M+Na^+$ ): 482.1433. Found: 482.1436.


Chiral HPLC: Retention time 16.6 min. with Chiralcel OD-R 250 x 4.6 cm column, 0 to 30% acetonitrile/0.5 M  $NaClO_4$  gradient over 20 min, flow rate 0.8 mL/min.

10 **Example A42:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2S-dimethylamino-propyl)-benzamide



The starting material was prepared as follows:


Step 1. (S)-2-Dimethylamino-propionamide Hydrochloride Salt



15 According to a procedure from Dallavalle et al, *Helv. Chim. Acta*; Vol. 72, pp. 1479-1486 (1989), (S)-2-amino-propionamide hydrochloride (1.0 g, 8.0 mmol; Advanced ChemTech) and formaldehyde (37% aq. soln., 662  $\mu$ L, 8.8 mmol) in MeOH (15 mL) were stirred in the presence of 10% Pd/C (320 mg) at 40°C under hydrogen for 5 hours. Catalyst was filtered off, and the filtrate was acidified to pH 3 with 1N HCl in MeOH. The solvent was  
20 removed and the resultant slurry was crystallized from MeOH/acetone to give a white solid, which was used without further purification.

$^1H$  NMR ( $DMSO-d_6$ ):  $\delta$  3.90 (m, 1H), 2.77 (d, 3H,  $J=4.9$  Hz), 2.74 (d, 3H,  $J=4.9$  Hz), 1.44 (d, 3H,  $J=7.0$  Hz).

Step 2. (S)- $N^2,N^2$ -Dimethyl-propane-1,2-diamine Dihydrochloride Salt



25

To the above intermediate in THF (10 ml) at 0°C under argon, was added dropwise 1M LiAlH<sub>4</sub> in THF (16 ml). The mixture was stirred at room temperature for one hour, then refluxed for 5 hours, cooled, and quenched with sat. Na<sub>2</sub>SO<sub>4</sub> (1 ml). The mixture was diluted with ethyl ether, and stirred another 10 minutes, filtered, acidified with 4N HCl/dioxane, and  
 5 evaporated to dryness to afford 0.70 g of white solid in 61% yield from (S)-2-amino-propionamide hydrochloride, which was used without further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 3.80 (m, 1H), 3.55 (dd, 2H, J=5.4, 13.6 Hz), 3.17 (dd, 1H, J=7.0, 13.6 Hz), 1.49 (d, 3H, J=6.8 Hz).


The title compound was prepared in a manner similar to Step 3 in Method A.

10 <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.88 (d, 2H, J=8.8 Hz), 7.75 (d, 2H, J=8.8 Hz), 7.49 (m, 1H), 7.08 (t, 2H, J=7.8 Hz), 3.51 (dd, 1H, J=6.5, 13.6 Hz), 3.39 (dd, 1H, J=6.5, 13.6 Hz), 2.89 (sextet, 1H, J=6.6 Hz), 2.35 (s, 6H), 1.08 (d, 3H, J=6.0 Hz).

<sup>13</sup>C NMR (DMSO-d<sub>6</sub>): δ 11.3, 40.3, 42.1, 57.8, 96.3, 112.1, 118.3, 119.2, 128.3, 129.4, 131.6, 141.7, 164.6, 165.3, 167.2, 172.8.


15 Anal. Calcd. For C<sub>22</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S: C, 57.50%; H, 5.05%; N, 15.24%; S, 6.98%. Found: C, 57.50%; H, 5.06%; N, 15.23%; S, 6.93%.

**Example A43:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-piperidin-2S-yl-methyl)-benzamide



20 The starting material was prepared as follows:

(1-Methyl-piperidin-2S-yl)-methylamine



Anhydrous ammonia gas was bubbled through a solution of (S)-(-)-1-(*tert*-butoxycarbonyl)-2-piperidine carboxylic acid (500 mg, 2.18 mmol) in DMF (50 ml) for 15  
 25 minutes. PyBop (1.25 g, 2.40 mmol) was added. After 5 hours, solvent was evaporated under reduced pressure. A solution of the resultant residue in ethyl acetate was washed with 5% citric acid (50 ml x 3), 1 N NaOH (50 ml x 3), dried with MgSO<sub>4</sub>, filtered, and concentrated to give 450

mg of (S)-2-carbamoyl-piperidine-1-carboxylic acid *tert*-butyl ester as a clear oil in 95% yield, which was used without further purification.

- (1-Methyl-piperidin-2S-yl)-methylamine was prepared in a manner similar to the production step of (S)-N<sup>1</sup>, N<sup>1</sup>, N<sup>2</sup>-trimethyl-propane-1,2-diamine in Example A36 from (S)-2-carbamoyl-piperidine-1-carboxylic acid *tert*-butyl ester in 56% yield.


<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ3.58-3.49 (m, 1H), 3.48-3.36 (m, 1H), 3.13-3.02 (m, 1H), 2.81 (s, 3H), 2.16-1.60 (m, 7H).

The title compound was prepared in a manner similar to Step 3 in Method A.

- <sup>1</sup>H NMR (acetone-d<sub>6</sub>): δ7.71 (m, 2H), 7.54 (m, 3H), 7.11 (m, 2H), 4.06 (m, 1H), 3.81-3.52 (m, 3H), 3.18 (s, 3H), 2.97 (m, 2H), 2.00-1.61 (m, 5H).

HRMALDIMS Calcd for C<sub>24</sub>H<sub>25</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S (M+H<sup>+</sup>): 486.1770. Found: 486.1761

**Example A44:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2R-dimethylamino-propyl)-benzamide




- The starting material (R)- N<sup>2</sup>,N<sup>2</sup>-dimethyl-propane-1,2-diamine hydrochloride was prepared with a route similar to that for Example A42, from (L)-2-amino-propionamide hydrochloride.

The title compound was prepared in a manner similar to Step 3 in Method A.

Spectral data are identical to Example A42.


- Anal. Calcd. For C<sub>22</sub>H<sub>23</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S: C, 57.50%; H, 5.05%; N, 15.24%; S, 6.98%. Found: C, 57.72%; H, 5.68%; N, 15.31%; S, 7.12%.

**Example A45:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-[2-(2,5-dihydro-pyrrol-1-yl)-ethyl]-benzamide Dihydrochloride Salt



- The starting material was prepared as follows:


Step 1. [2-(2,5-Dihydro-pyrrol-1-yl)-2-oxo-ethyl]-carbamic Acid *tert*-Butyl Ester



To a solution of N-*tert*-butoxycarbonyl glycine (1.4 g, 5.0 mmol), DIEA (2.6 ml, 15 mmol) and 2,5-dihydro-1*H*-pyrrole (0.51 g, 7.5 mmol) in DMF (15 ml), was added PyBop (2.7 g, 5.25 mmol). The reaction solution was stirred at room temperature for 3 hours. Solvent was removed under reduced pressure and a solution of resultant residue in ethyl acetate was extracted with sat. NaHCO<sub>3</sub>, brine, dried with MgSO<sub>4</sub>, filtered and concentrated to a syrup. Chromatography on silica (hexane/ethyl acetate=1/2) afforded 1.05 g of desired product in 93% yield, which was used without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 5.88-5.72 (m, 2H), 4.24-4.10 (m, 4H), 3.82 (s, 2H), 1.4 (s, 9H).


Step 2. 2-Amino-1-(2,5-dihydro-pyrrol-1-yl)-ethanone



A solution of above intermediate (1.0 g, 4.6 mmol) in 30% TFA/CH<sub>2</sub>Cl<sub>2</sub> (20 ml) was stirred at room temperature for 30 min. Solvent was removed and a solution of resultant residue in ethyl acetate was extracted with sat. NaHCO<sub>3</sub>, dried with MgSO<sub>4</sub>, filtered and concentrated to give 0.5 g of product as a clear oil in 95% yield, which was used without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 5.80 (m, 2H), 4.20 (m, 4H), 3.45 (s, 2H).

Step 3. 2-(2,5-Dihydro-pyrrol-1-yl)-ethylamine Dihydrochloride Salt



The desired amine was obtained after LiAlH<sub>4</sub> reduction of the above intermediate as described in Step 2 in Example A42 in 50% yield and used as crude.

The title compound was prepared in a manner similar to Step 3 in Method A.


<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.92 (d, 2H, J=8.8 Hz), 7.78 (d, 2H, J=8.8 Hz), 7.50 (m, 1H), 7.08 (t, 2H, J=15.9 Hz), 5.98 (s, 2H), 4.49 (d, 2H, J=12.4 Hz), 4.12 (d, 2H, J=12.4 Hz), 3.78 (t, 2H, J=6.1 Hz), 3.58 (t, 2H, J=5.6 Hz).

LCMS (M+H<sup>+</sup>): 470; (M-H<sup>-</sup>): 468.

Anal. Calcd. for C<sub>23</sub>H<sub>21</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•2.05 HCl•1.10 H<sub>2</sub>O: C, 48.97%; H, 4.51%; N, 12.42%; S, 5.68%. Found: C, 49.13%; H, 4.71%; N, 12.29%; S, 5.37%.



**Example A46:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1R-methyl-2-piperidin-1-yl-ethyl)-benzamide Trifluoroacetate Salt



The starting material 1R-methyl-2-piperidin-1-yl-ethylamine was prepared in a route similar to that for Example A45 from *N-tert*-butoxycarbonyl D-alanine and piperidine in 43% overall yield and used without further purification.

The title compound was prepared in a manner similar to Step 3 in Method A.


<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 10.08 (d, 2H, J=8.8 Hz), 7.68 (d, 2H, J=8.8 Hz), 7.48 (m, 1H), 6.96 (t, 2H, J=15.9 Hz), 4.54 (m, 1H), 3.86 (m, 1H), 3.38 (m, 1H), 3.18 (m, 2H), 2.90 (m, 2H), 1.95-1.40 (m, 6H), 1.27 (d, 2H, J=6.8 Hz).

MALDIMS (M+H<sup>+</sup>): 500.

Anal. Calcd. for C<sub>25</sub>H<sub>27</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•1.50 TFA•0.60 H<sub>2</sub>O: C, 49.35%; H, 4.39%; N, 10.28%; S, 4.71%.

Found: C, 49.29%; H, 4.42%; N, 10.38%; S, 4.63%.

**Example A47:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1R-dimethylaminomethyl-2-methyl-propyl)-benzamide.



The starting material (R)-3,N<sup>1</sup>,N<sup>1</sup>-trimethyl-butane-1,2-diamine was prepared in a route similar to that in Example A45 from *N-tert*-butoxycarbonyl D-valine and dimethylamine and used without further purification.


The title compound was prepared in a manner similar to Step 3 in Method A.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.10 (br, 1H), 8.22 (br, 2H), 8.01-7.89 (m, 3H), 7.72 (d, 2H, J=8.6 Hz), 7.62 (m, 1H), 7.26 (t, 2H, J=7.8 Hz), 4.11 (m, 1H), 2.50 (m, 1H), 2.34 (br, 6H), 1.88 (m, 1H), 0.99-0.88 (m, 7H).

FABMS (MH<sup>+</sup>): 488; (M-H<sup>-</sup>): 486.

Anal. Calcd. For  $C_{24}H_{27}F_2N_5O_2S \cdot 1.08 H_2O \cdot 0.35 EtOAc$ : C, 56.72%; H, 5.99%; N, 13.02%; S, 5.96%. Found: C, 56.66%; H, 5.77%; N, 13.03%; S, 5.93%.

**Example A48:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(5,5-dimethyl-pyrrolidin-2RS-yl-methyl)-benzamide Hydrochloride Salt



5

The starting material 5-aminomethyl-2,2-dimethyl-pyrrolidin-1-ol was prepared according to the literature procedure (Zhong et al., *Bioorg. Med. Chem.* Vol. 6, pp.2405-2419, (1998)) from 5,5-dimethyl-1-pyrroline *N*-oxide.

The intermediate 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-hydroxy-5,5-dimethyl-pyrrolidin-2-ylmethyl)-benzamide was prepared in a manner similar to Step 3 in Method A from the above amine.

$^1H$  NMR ( $CDCl_3$ ):  $\delta$  8.14 (d, 2H,  $J=8.9$  Hz), 7.95 (d, 2H,  $J=8.9$ ), 7.68 (m, 1H), 7.25 (t, 2H,  $J=9.0$  Hz), 4.20 (m, 2H), 3.85 (m, 1H), 2.56-2.08 (m, 4H), 1.68 (s, 3H), 1.58 (s, 3H).


The title compound was prepared as follows:

A solution of above intermediate in MeOH was hydrogenated on 10% Pd/C at 30 psi for 3 hours. The catalyst was filtered off and the filtrate was concentrated. Preparative HPLC purification afforded a white solid in 25% yield.

$^1H$  NMR ( $CD_3OD$ ):  $\delta$  7.84 (d, 2H,  $J=8.8$  Hz), 7.73 (d, 2H,  $J=8.8$  Hz), 7.42 (m, 1H), 6.98 (t, 2H,  $J=15.9$  Hz), 3.92 (m, 1H), 3.62 (d, 2H,  $J=5.5$  Hz), 2.22 (m, 1H), 1.94-1.81 (m, 3H), 1.40 (s, 6H).

LC-MS ( $M+H^+$ ): 486; ( $M-H$ ): 484.

**Example A49:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(thiazol-2-yl-methyl)-benzamide Hydrochloride



The starting material, C-thiazol-2-yl-methylamine, was prepared as follows.

According to a procedure from Kuo et al, *Chem. Pharm. Bull.*, 39, 181-183 (1991), to a solution of 2-thiazolecarboxaldehyde (1.52 g, 13.4 mmol) in ethanol (16 ml) and  $H_2O$  (3.3 ml) were added hydroxylamine hydrochloride (1.40 g, 20.1 mmol) and NaOH (1.61 g, 40.3 mmol).


The mixture was heated at reflux for 0.5 h, allowed to cool to ambient temperature, and acidified to pH4 with 2N HCl. The aqueous layer was extracted with ether (100 ml × 2). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to give a white solid, which was placed in a mixture of ethanol (30 ml) and conc. aq. NH<sub>4</sub>OH (67 ml). Zn dust  
 5 (6.58 g, 101 mmol) and ammonium acetate (0.826 g, 10.7 mmol) were added. The mixture was heated at reflux for 0.5 hour, allowed to cool to ambient temperature, and filtered. The filtrate was diluted with H<sub>2</sub>O (25 ml) and extracted with 10% MeOH/CHCl<sub>3</sub> (50 ml). The organic layer was separated, dried over MgSO<sub>4</sub>, and concentrated to afford 998 mg of yellow oil in 65% yield, which displayed an <sup>1</sup>H NMR that matched previously reported (Dondoni et al., *Synthesis*, 641-646 (1996)) and was used without further purification.  
 10

The title compound was prepared in a manner similar to Step 3 in Method A.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.96 (s, 1H), 9.10 (t, 1H, J=5.9 Hz), 8.05 (br, 2H), 7.72 (d, 2H, J=8.8 Hz), 7.55 (d, 1H, J=3.3 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.44 (d, 1H, J=3.3 Hz), 7.37 (m, 1H), 7.03 (dd, 2H, J=7.8, 8.0 Hz), 4.56 (d, 2H, J=5.9 Hz).


15 Anal. Calcd. for C<sub>21</sub>H<sub>15</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S<sub>2</sub>•1.4 HCl•1.0 H<sub>2</sub>O: C, 46.66%; H, 3.43%; N, 12.96%; S, 11.86%. Found: C, 46.77%; H, 3.51%; N, 13.02%; S, 11.81%.

**Example A50:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-1H-imidazol-2-ylmethyl)-benzamide Hydrochloride Salt



20 The starting material was prepared as follows:

*C-(1-Methyl-1H-imidazol-2-yl)-methylamine.*



According to a procedure from Kruse et al, *J. Med. Chem.* Vol. 33, pp. 781-789 (1990), hydroxylamine hydrochloride (3.78 g, 54.5 mmol) and sodium acetate (9.38 g, 114 mmol) were  
 25 added to a solution of 1-methyl-2-imidazolecarboxaldehyde (2.00 g, 18.2 mmol) in H<sub>2</sub>O (90 ml). After one hour, the resultant mixture was filtered. The white solid was washed with a small amount of water and dried under vacuum to give 1.01 g of a colorless solid in 44% yield. This presumed 1-methyl-1H-imidazole-2-carboxaldehyde (E/Z)-oxime was submitted to the

conditions described for thiazol-2-yl-methylamine in Example A49 to give 643 mg of yellow oil in 72% yield, which was used without any further purification.


$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  6.94 (d, 1H,  $J=1.2$  Hz), 6.82 (d, 1H,  $J=1.2$  Hz), 3.91 (s, 2H), 3.64 (s, 3H).

The title compound was prepared in a manner similar to Step 3 in Method A.

- 5  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  14.22 (br, 1H), 11.20 (s, 1H), 9.21 (t, 1H,  $J=5.2$  Hz), 8.11 (br, 2H), 7.82 (d, 2H,  $J=8.8$  Hz), 7.64 (d, 2H,  $J=8.8$  Hz), 7.55 (d, 1H,  $J=1.9$  Hz), 7.49 (d, 1H,  $J=1.9$  Hz), 7.45 (m, 1H), 7.11 (dd, 2H,  $J=7.7, 8.2$  Hz), 4.64 (d, 2H,  $J=5.2$  Hz).

Anal. Calcd. for  $\text{C}_{22}\text{H}_{18}\text{F}_2\text{N}_6\text{O}_2\text{S} \cdot 1.6 \text{ HCl} \cdot 1.1 \text{ H}_2\text{O}$ : C, 48.34%; H, 4.02%; N, 15.37%; S, 5.87%.  
Found: C, 48.23%; H, 4.29%; N, 15.27%; S, 5.86%.

- 10 **Example A51:** 4-[4-Amino-5-2,6-difluoro-benzoyl]-thiazol-2-ylamino]-N-[2-(cis-2,6-dimethyl-piperidin-1-yl)-ethyl]-benzamide



- The starting material 1-(2-aminoethyl)-2,6-dimethylpiperidine was prepared in a route similar to that for  $\text{N}^2$ -cyclopropylmethyl-2, $\text{N}^2$ -dimethylpropane-1,2-diamine in Example A39 from glycolonitrile and 2,6-dimethylpiperidine.


$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 2.76-2.65 (m, 4H), 2.56-2.41 (m, 2H), 1.77-1.48 (m, 5H), 1.38-1.19 (m, 3H), 1.14 (d, 6H,  $J=6.3$  Hz).

The title compound was prepared in a manner similar to Step 3 in Method A.

- 15  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  10.85 (br, 1H), 8.3 (m, 3H), 7.65 (d, 2H,  $J=8.8$  Hz), 7.49 (d, 2H,  $J=8.8$  Hz), 7.40 (m, 1H), 7.05 (t, 2H,  $J=7.7$  Hz), 3.10 (m, 2H), 2.6-2.4 (m, 4H), 1.5-1.2 (m, 6H), 0.95 (d, 6H,  $J=6.1$  Hz).

Anal. Calcd for  $\text{C}_{26}\text{H}_{29}\text{F}_2\text{N}_5\text{O}_2\text{S} \cdot 1.0 \text{ H}_2\text{O}$ : C, H, N.

**Example A52:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-methyl-2-piperidin-1-yl-propyl)-benzamide



The starting material 1-(2-amino-1,1-dimethylethyl)piperidine was prepared in a route similar to that in Example 39 from 2-hydroxy-2-methyl-propionitrile and piperidine.


$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  2.56 (s, 2H), 2.52-2.41 (m, 4H), 1.87 (br, 2H), 1.61-1.49 (m, 4H), 1.46-1.35 (m, 2H), 0.97 (s, 6H).

The title compound was prepared in manner similar to Step 3 in Method A.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  8.18 (br, 2H), 7.8 (d, 2H,  $J=8.7$  Hz), 7.67 (d, 2H,  $J=8.7$  Hz), 7.56 (m, 1H), 7.20 (t, 1H,  $J=7.8$  Hz), 3.45 (m, 2H), 1.7-1.4 (m, 10H), 1.00 (s, 6H).

Anal. Calcd For  $\text{C}_{26}\text{H}_{29}\text{F}_2\text{N}_5\text{O}_2\text{S} \cdot 0.5 \text{H}_2\text{O}$ : C, H, N.

**Example A53:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-[2-(2,2,6,6-tetramethyl-piperidin-1-yl)-ethyl]-benzamide



The starting material 1-(2-aminoethyl)-2,2,6,6-tetramethylpiperidine was prepared in a route similar to that of Example A39 from 2,2,6,6-tetramethylpiperidine and glycolonitrile.


$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  2.46-2.26 (m, 4H), 1.52-1.40 (m, 2H), 1.35-1.25 (m, 4H), 0.97 (s, 12H).

The title compound was prepared in a manner similar to Step 3 in Method A.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  8.4 (m, 3H), 7.85 (d, 2H,  $J=8.7$  Hz), 7.65 (d, 2H,  $J=8.7$  Hz), 7.59 (m, 1H), 7.22 (m, 2H), 3.2 (m, 2H), 3.16 (m, 1H), 1.7 (m, 1H), 1.4-1.2 (m, 6H), 1.06 (s, 12H)


Anal. Calcd For  $\text{C}_{28}\text{H}_{33}\text{F}_2\text{N}_5\text{O}_2\text{S} \cdot 1.0 \text{H}_2\text{O}$ : C, H, N.

**Example A54:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(4R-hydroxy-1-methyl-pyrrolidin-2S-yl-methyl)-benzamide



The starting material was prepared as follows:

Step 1. *[4S,2S]-2-Carbamoyl-4-hydroxy-pyrrolidine-1-carboxylic Acid Benzyl Ester*




- 5        4S-Hydroxy-pyrrolidine-1,2S-dicarboxylic acid 1-benzyl ester (Bachem, 5.00 g, 18.8 mmol) in THF (25 ml) was cooled to 0°C. Pyridine (892 mg, 11.3 mmol),  $\text{NH}_4\text{HCO}_3$  (1.93 g, 24.4 mmol) and di-*tert*-butyl dicarbonate (5.3 g, 24.4 mmol) were added. The reaction was allowed to reach 25°C over 1 h and was stirred an additional 24 h. The mixture was diluted with EtOAc, washed with saturated aqueous  $\text{NaHCO}_3$ , 0.4 N  $\text{NaHSO}_4$ , brine, dried over  $\text{Na}_2\text{SO}_4$  and
- 10    concentrated to a syrup. Chromatography on silica (3%-12%  $\text{MeOH-CH}_2\text{Cl}_2$ ) afforded 1.79 g of amide as a white foam in 38% yield, which was used without further purification.

$R_f=0.22$  (10%  $\text{MeOH-CH}_2\text{Cl}_2$ ).

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.32 (s, 5H), 6.72, 5.97 (two bs, 1H), 5.53 (two bs, 1H), 5.13 (m, 2H), 4.45 (bs, 2H), 3.58 (bs, 2H), 2.38 (bm, 2H), 2.11 (bs, 1H).

- 15    LCMS ( $\text{M}+\text{H}^+$ ): 265; ( $\text{M}+\text{Na}^+$ ): 287.

Step 2. *[5R,3S]-5-Aminomethyl-1-methyl-pyrrolidin-3-ol*



- To the above amide (1.7g, 6.43 mmol) in THF (30 ml) at 0°C was added lithium aluminum hydride (1.22 g, 32.2 mmol) in 15 mL THF. The mixture was refluxed for 12 hours. The reaction
- 20    was cooled to 0°C and treated with solid  $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$  until gas evolution ceased. The whole was filtered and concentrated to yield 1.2 g of desired amine as a clear oil which was used directly in the next step.


The title compound was prepared in a manner similar to Step 3 in Method A.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ -acetone- $d_6$ ):  $\delta$  7.58 (d, 2H,  $J=8.6$  Hz), 7.38 (d, 2H,  $J=8.6$  Hz), 7.17 (m, 1H), 6.74 (t, 3H,  $J=8.1$  Hz), 4.14 (m, 1H), 3.82 (m, 1H), 3.54 (m, 1H), 3.20 (dd, 1H,  $J=9.9$  Hz, 13.1 Hz), 3.07 (bd, 1 H,  $J=13.1$  Hz), 2.65 (m, 1H), 2.19 (s, 3H), 1.72 (m, 2H).

LCMS ( $\text{M}+\text{H}^+$ ): 488.

5     Anal. Calcd For  $\text{C}_{23}\text{H}_{23}\text{F}_2\text{N}_5\text{O}_3\text{S}$ : C, H, N.

**Example A55:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-[2-(imidazol-1-yl)-ethyl]-benzamide Dihydrochloride Salt



The starting material 2-imidazol-1-yl-ethylamine was prepared according to the  
10     literature procedure (Hay et al., *J. Med. Chem.*, 37; pp. 381-391 (1994)).


The title compound was prepared in a manner similar to Step 3 in Method A.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  14.48 (bs, 1H), 11.40 (bs, 1H), 9.15 (s, 1H), 8.69 (t, 1H,  $J=5.4$  Hz),  
8.18 (bs, 2H), 7.80 (d, 2H,  $J=8.8$  Hz), 7.75 (t, 1H,  $J=1.5$  Hz), 7.69 (d, 2H,  $J=8.8$  Hz), 7.63 (t,  
1H,  $J=1.5$  Hz), 7.52 (m, 1H), 7.17 (dd, 2H,  $J=7.7$ , 8.1 Hz), 4.37 (t, 2H,  $J=5.4$  Hz), 3.67 (q, 2H,  
15      $J=5.4$  Hz).

LCMS: ( $\text{M}+\text{H}^+$ ): 469.

Anal. Calcd. for  $\text{C}_{22}\text{H}_{18}\text{F}_2\text{N}_6\text{O}_2\text{S}\cdot 2.5 \text{ HCl}\cdot 1.0 \text{ H}_2\text{O}$ : C, 45.74%; H, 3.93%; N, 14.55%; S, 5.55%.  
Found: C, 45.45%; H, 4.16%; N, 14.55%; S, 5.57%.

**Example A56:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2,2,5,5-tetramethyl-  
20     pyrrolidin-3R,S-ylmethyl)-benzamide Dihydrochloride Salt



The intermediate 4-[4-amino-5-[1-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-hydroxy-2,2,5,5-tetramethyl-pyrrolidin-3-ylmethyl)-benzamide] was prepared in a manner similar to Step 3 in Method A, from 3R,S-aminomethyl-2,2,5,5-tetramethyl-1-pyrrolidinyloxy.

The title compound was prepared in a manner similar to that for Example A48 in 30% yield after HPLC purification.


$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.88 (d, 2H,  $J=8.8$  Hz), 7.88 (d, 2H,  $J=8.8$  Hz), 7.58 (m, 1H), 7.13 (t, 2H,  $J=8.7$  Hz), 2.67 (m, 1H), 2.22-2.00 (m, 2H), 1.58 (s, 6H), 1.47 (d, 6H,  $J=9.4$  Hz), 1.20 (m, 2H).

LCMS ( $\text{M}+\text{H}^+$ ): 514.

Anal. Calcd. For  $\text{C}_{26}\text{H}_{29}\text{F}_2\text{N}_5\text{O}_2\text{S}\cdot 2\text{HCl}\cdot 1.60\text{H}_2\text{O}$ : C, 50.75%; H, 5.60%; N, 11.38%; S, 5.21%.


Found: C, 50.75%; H, 5.74%; N, 11.35%; S, 5.21%.

**Example A57:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-2-(cis-3,5-dimethyl-piperazin-1-yl)-ethyl]-benzamide



The starting materials were prepared as follows:

Step 1. *cis*-3,5-Dimethyl-piperazine-1-carbonitrile




To a solution of *cis*-3,5-dimethyl-piperazine (1.5 g, 13.13 mmol) in  $\text{CH}_3\text{CN}$  (10 ml) were added  $\text{K}_2\text{CO}_3$  (8.70 g, 63.02 mmol) and chloroacetonitrile (0.90 ml, 14.44 mmol). The mixture



was stirred for 12 h, filtered, and concentrated to afford the desired product as a brown solid in 78% yield, which was used without further purification.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.08 (3H, s), 3.00-2.88 (2H, m), 1.94 (2H, dd,  $J = 10.4, 10.5$  Hz), 1.08 (3H, s), 1.06 (3H, s).

5 Step 2. 2-(*cis*-3,5-Dimethyl-piperazin-1-yl)-ethylamine




The title amine was prepared according to the procedure described in Step 2 in Example A39 and used as crude.

The title compound was prepared in a manner similar to Step 3 in Method A.

10  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.83 (d, 2H,  $J = 8.9$  Hz), 7.72 (d, 2H,  $J = 8.7$  Hz), 7.52-7.42 (m, 1H), 7.06 (dd, 2H,  $J = 7.9, 8.0$  Hz), 3.54 (dd, 2H,  $J = 6.6, 6.8$  Hz), 2.61 (dd, 2H,  $J = 6.4, 7.2$  Hz), 1.76 (dd, 2H,  $J = 10.8, 11.2$  Hz), 1.18 (dd, 2H,  $J = 6.9, 7.1$  Hz), 1.08 (s, 3H), 1.06 (s, 3H).

Anal. Calcd. for  $\text{C}_{25}\text{H}_{28}\text{F}_2\text{N}_6\text{O}_2\text{S} \cdot 1.5 \text{ H}_2\text{O}$ : C, 55.44%; H, 5.77%; N, 15.52%; S, 5.92%. Found: C, 55.31%; H, 5.47%; N, 15.35%; S, 5.96%.

15 **Example A58:** 4-[5-(1-Adamantan-1-yl-methanoyl)-4-amino-thiazol-2-ylamino]-N-(2-dimethylamino-ethyl)-benzamide




The title compound was prepared from 1-adamantyl bromomethyl ketone in a route similar to that of Method A.

20  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  11.00 (bs, 1H), 8.30 (bs, 1H), 8.10 (br, 2H), 7.88 (d, 2H,  $J=6.5$  Hz), 7.74 (d, 2H,  $J=6.5$  Hz), 2.40 (m, 2H), 2.20 (s, 6H), 2.05 (m, 3H), 1.92 (m, 6H), 1.70 (m, 6H).


ESIMS ( $\text{M}+\text{H}^+$ ): 468

**Example A59:** 4-{4-Amino-5-[1-(4-methyl-pyridin-3-yl)methanoyl]-thiazol-2-ylamino}-N-(1-methyl-pyrrolidin-2S-ylmethyl)-benzamide Hydrochloride Salt.



The starting materials were prepared as follows:

Step 1. 3-(1-Ethoxy-vinyl)-4-methyl-pyridine



- 5            3-Bromo-4-methyl-pyridine (2.46 g, 14.3 mmol; Emka Chemie), tributyl-(1-ethoxyvinyl)-stannane (6.20 g, 17.2 mmol), and tetrakis(triphenylphosphine)-palladium(0) (1.32 g, 1.14 mmol) stirred in dry toluene (50 ml) at reflux under argon for 4 hours. The mixture was concentrated *in vacuo* and purified by column chromatography (25% EtOAc/hexanes) to give 2.12 g of a clear oil in 91% yield, which was used without any further purification.
- 10        <sup>1</sup>H NMR: δ 8.50 (s, 1H), 8.40 (d, 1H, J=5.1 Hz), 7.08 (s, 1H, J=5.1 Hz), 4.38 (d, 1H, J=2.4 Hz), 4.24 (d, 1H, J=2.4 Hz), 3.91 (q, 2H, J=7.2 Hz), 2.37 (s, 3H), 1.38 (t, 3H, J=7.2 Hz).


Step 2. 1-(4-Methyl-pyridin-3-yl)-ethanone.



- 15            3-(1-Ethoxy-vinyl)-4-methyl-pyridine stirred in a mixture of toluene (150 ml), water (30 ml), and concentrated HCl (30 ml) for 2 hours. The solution was concentrated to ~40 ml *in vacuo*, made basic with sat. NaHCO<sub>3</sub>, and extracted with EtOAc. Organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo*. Purification by column chromatography (75% EtOAc/hexanes) gave 1.7 g of a clear oil in 96% yield, which was used without any further purification.

- 20        <sup>1</sup>H NMR: δ 8.95 (s, 1H), 8.54 (d, 1H, J=5.1 Hz), 7.19 (d, 1H, J=5.1 Hz), 2.64 (s, 3H), 2.56 (s, 3H).

Step 3. 2-Bromo-1-(4-methyl-pyridin-3-yl)-ethanone Hydrobromide Salt




Bromine (0.66 ml, 12.7 mmol) in acetic acid (2.5 ml) was added slowly to a stirred solution of 1-(4-methyl-pyridin-3-yl)-ethanone (1.68 g, 12.4 mmol) in acetic acid (2.5 ml) and 48% aqueous HBr (2.5 ml) at 0°C. The reaction mixture was stirred for 16 hours at room temperature and then diluted with ether (50 ml) to form a precipitate. The white solid was collected by filtration and washed with ether and acetone to give 2.59 g of white solid in 71% yield, which was used without any further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 9.24 (s, 1H), 8.83 (d, 1H, J=5.7 Hz), 7.86 (d, 1H, J=5.7 Hz), 5.02 (s, 2H), 2.58 (s, 3H).

Step 4. 4-{4-Amino-5-[1-(4-methyl-pyridin-3-yl)-methanoyl]-thiazol-2-ylamino}-benzoic acid

10 Ethyl Ester




The title compound was prepared in a manner similar to that for Step 1 in Method A from ethoxycarbonylphenyl isothiocyanate (700 mg, 3.40 mmol) and 2-bromo-1-(4-methyl-pyridin-3-yl)-ethanone hydrobromide (1.0 g, 3.4 mmol) and purified by silica gel chromatography (7:2:1

15 EtOAc/THF/hexanes) to provide 967 mg of a yellow solid in 75% yield.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.11 (s, 1H), 8.47-8.50 (m, 2H), 8.20 (br s, 2H), 7.93 (d, 2H, J=8.7 Hz), 7.73 (d, 2H, J=8.7 Hz), 7.32 (d, 1H, J=5.1 Hz), 4.27 (q, 2H, J=7.2 Hz), 2.30 (s, 3H), 1.30 (t, 3H, J=7.2 Hz).

Anal. (C<sub>19</sub>H<sub>18</sub>N<sub>4</sub>O<sub>3</sub>S) C, H, N, S.

20 Step 5. 4-{4-Amino-5-[1-(4-methyl-pyridin-3-yl)-methanoyl]-thiazol-2-ylamino}-benzoic Acid




4-{4-Amino-5-[1-(4-methyl-pyridin-3-yl)-methanoyl]-thiazol-2-ylamino}-benzoic acid ethyl ester was hydrolyzed by refluxing for 5 hours with lithium hydroxide in water/THF (1/1) to give the title compound in a quantitative yield, which was used without any further purification.

25 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.37 (br, 1H), 8.47 (bd, 2H, J=5.1 Hz), 8.19 (br, 2H), 7.80 (d, 2H, J=8.7 Hz), 7.74 (d, 2H, J=8.7 Hz), 7.32 (d, 1H, J=5.1 Hz), 2.29 (s, 3H).

ESIMS (M+H<sup>+</sup>): 355.

Step 6. 4-{4-Amino-5-[1-(4-methyl-pyridin-3-yl)methanoyl]-thiazol-2-ylamino}-N-(1-methyl-pyrrolidin-2S-ylmethyl)-benzamide



The title compound was prepared in a manner similar to that of Step 3 in Method A from 4-{4-amino-5-[1-(4-methyl-pyridin-3-yl)-methanoyl]-thiazol-2-ylamino}-benzoic acid (420 mg, 1.19 mmol) and (1-methyl-pyrrolidin-2S-yl)-methylamine (from Example A31; 150 mg, 1.32 mmol) as a bright yellow solid (320 mg) in 51% yield, which was used without any further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.01 (br s, 1H), 8.45-8.48 (m, 2H), 8.24 (t, 1H, J=6.0 Hz), 8.17 (br s, 2H), 7.80 (d, 2H, J=8.7 Hz), 7.61 (d, 2H, J=8.7 Hz), 7.31 (d, 1H, J=5.1 Hz), 3.38-3.46 (m, 1H), 3.07-3.14 (m, 1H), 2.87-2.93 (m, 1H), 2.28-2.34 (m, 7H), 2.10 (q, 1H, J=8.4 Hz), 1.78-1.82 (m, 1H), 1.52-1.62 (m, 3H).


ESIMS (M+H<sup>+</sup>): 451.

For the title compound, a solution of 4-{4-amino-5-[1-(4-methyl-pyridin-3-yl)methanoyl]-thiazol-2-ylamino}-N-(1-methyl-pyrrolidin-2S-ylmethyl)-benzamide was dissolved in 1 N HCl, washed with EtOAc, and concentrated *in vacuo* to give a yellow solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.88 (s, 1H), 10.79 (s, 1H), 8.98-9.04 (m, 2H), 8.83 (d, 1H, J=6.0 Hz), 8.40 (bs, 2H), 8.03 (s, 1H), 7.99 (d, 2H, J=8.7 Hz), 7.82 (d, 2H, J=8.7 Hz), 3.50-3.82 (m, 4H), 3.00-3.07 (m, 1H), 2.84 (d, 3H, J=5.1 Hz), 2.54 (s, 3H), 1.77-2.12 (m, 4H), 124-1.31 (m, 1H).

Anal. For C<sub>23</sub>H<sub>26</sub>N<sub>6</sub>O<sub>2</sub>S • 2.0 HCl • 2.0 H<sub>2</sub>O: C, H, N, S.

**Example A60:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2RS-dimethylamino-1-methyl-ethyl)-2-methoxy-benzamide




The title compound was prepared starting from 4-amino-2-methoxy-benzoic acid methyl ester and 2-bromo-2',6'-difluoro-acetophenone in a route similar to that of Method A.

$^1\text{H}$  NMR:  $\delta$  7.92 (d, 1H,  $J=8.8$  Hz), 7.28 (m, 1H), 7.12 (s, 1H), 6.88 (t, 2H,  $J=8.7$  Hz), 6.72 (d, 1H,  $J=8.8$  Hz), 4.24 (m, 1H), 3.86 (s, 3H), 2.50 (m, 1H), 2.28 (s, 6H), 1.22 (d, 3H,  $J=6.5$  Hz).

HRFABMS Calcd for  $\text{C}_{23}\text{H}_{25}\text{F}_2\text{N}_5\text{O}_3\text{S}$  ( $\text{M}+\text{H}^+$ ): 490.1724. Found: 490.1722.

5 **Example A61:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-1,1-dimethyl-ethyl)-benzamide Dihydrochloride



The title compound was prepared in a manner analogous to Step 3 of Method A, from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**) and 2, $N^1$ , $N^1$ -trimethylpropane-1,2-diamine (Tsuji, et al *Chem. Pharm. Bull.* Vol. 12, pp. 946-950 (1964)). Radial chromatography with 0.5% (58%  $\text{NH}_4\text{OH}$ )/ 5% MeOH/  $\text{CHCl}_3$  gave a yellow oil, which was placed in  $\text{CHCl}_3$ , treated with 4M HCl in dioxane (2.2equiv), and azeotroped from  $\text{CHCl}_3$  in succession to provide 100 mg of yellow powder in 23% yield, mp 275-290°C (decomp).


15  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  8.16 (bs, 1H), 7.91 (s, 1H), 7.90 (d, 2H,  $J=6.8$  Hz), 7.77 (d, 2H,  $J=6.8$  Hz), 7.51 (ddd, 1H,  $J=6.4, 8.4, 15.1$  Hz), 7.08 (dd, 2H,  $J=7.6, 8.3$  Hz), 3.66 (s, 2H), 3.00 (s, 6H), 1.58 (s, 6H).

FTIR (KBr): 1602, 1610, 1543, 1524, 1464, 1426  $\text{cm}^{-1}$ .

HRESIMS. Calcd for  $\text{C}_{23}\text{H}_{26}\text{F}_2\text{N}_5\text{O}_2\text{S}$  ( $\text{M}+\text{H}^+$ ): 474.1775. Found: 474.1793.


20 Anal. Calcd. for  $\text{C}_{23}\text{H}_{25}\text{F}_2\text{N}_5\text{O}_2\text{S} \cdot 2.0 \text{ HCl} \cdot 0.3 \text{ CHCl}_3 \cdot 0.9 \text{ H}_2\text{O}$ : C, 46.76%; H, 4.90%; N, 11.70%; Cl, 17.18%; S, 5.36%. Found: C, 46.77%; H, 4.66%; N, 11.34%; Cl, 17.19%; S, 5.28%.

25 **Example A62:** 4-{4-Amino-5-[1-(3-methyl-thiophen-2-yl)-methanoyl]-thiazol-2-ylamino}-N-(2-dimethylamino-1R-methyl-ethyl)-benzamide Trifluoroacetic Acid Salt



Starting materials were prepared as follows:


Step 1. 4-{4-Amino-5-[1-(3-methyl-thiophen-2-yl)-methanoyl]-thiazol-2-ylamino}-benzoic Acid Ethyl Ester



Prepared in a manner analogous to Step 1 of Method A. To 4-ethoxycarbonyl-phenylisothiocyanate (0.930 g, 4.48 mmol) in CH<sub>3</sub>CN (5 ml) were sequentially added  
 5 cyanamide (0.207 g, 4.92 mmol) and a solution of potassium t-butoxide (0.552 g, 4.92 mmol) in t-butanol (5 ml). The resultant mixture stirred for 0.5 h, then 2-bromoacetyl-3-methyl-thiophene (1.00 g, 4.56 mmol; PCT Patent Publication WO 99/21845) was added. The mixture stirred for 3 h, was diluted with 10% MeOH/CHCl<sub>3</sub> (100 ml), and washed  
 10 with H<sub>2</sub>O (25 ml X 2). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to a brown solid, which recrystallized from hot EtOAc to give 1.1 g of a yellow solid in 65% yield, which was used without any further purification.

<sup>1</sup>H NMR: δ 8.18 (2H, d, J=8.8 Hz), 7.56 (2H, d, J=8.8 Hz), 7.42 (1H, d, J=5.0 Hz), 7.04 (1H, d, J=5.0 Hz), 4.49 (2H, q, J=7.1 Hz), 2.60 (3H, s), 1.50 (3H, t, J=7.1 Hz).

15 Step 2. 4-{4-Amino-5-[1-(3-methyl-thiophen-2-yl)-methanoyl]-thiazol-2-ylamino}-benzoic Acid



Prepared in a manner analogous to Step 2 of Method A. To a suspension of 4-{4-amino-5-[1-(3-methyl-thiophen-2-yl)-methanoyl]-thiazol-2-ylamino}-benzoic acid ethyl  
 20 ester (1.00 g, 2.58 mmol) in MeOH (6 ml) was added 3N NaOH (8.60 ml, 25.8 mmol). The resultant solution stirred at ambient temperature for 18 h. The methanol was removed in vacuo and the aqueous layer adjusted to pH 3 with 10% HCl to give 0.80 g of a yellow solid in 90% yield, which was used without further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.02 (1H, s), 8.02 (1H, bs), 7.90 (2H, d, J=8.7 Hz), 7.74 (2H, d, J=8.7 Hz), 7.61 (1H, d, J=4.9 Hz), 7.00 (1H, d, J=4.9 Hz), 2.4 (3H, s).


The title compound was prepared in a manner similar to Step 3 in Method A, from 4-{4-amino-5-[1-(3-methyl-thiophen-2-yl)-methanoyl]-thiazol-2-ylamino}-benzoic acid and (R)-

N,N-dimethyl-propane-1,2-diamine dihydrochloride salt (from Step 3 of Example A40) to provide a yellow foam in 62% yield.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.88 (2H, d,  $J=8.7$  Hz), 7.76 (2H, d,  $J=8.7$  Hz), 7.42 (1H, d,  $J=5.1$  Hz), 6.98 (1H, d,  $J=4.8$  Hz), 2.56 (6H, s), 2.42 (3H, s), 1.28 (3H, d,  $J=6.6$  Hz).


- 5 Anal. Calcd. for  $\text{C}_{21}\text{H}_{25}\text{N}_5\text{O}_2\text{S}_2 \cdot 0.6 \text{CF}_3\text{CO}_2\text{H}$ : C, 52.08%; H, 5.04%; N, 13.68%; S, 12.53%. Found: C, 52.28%; H, 5.22%; N, 13.34%; S, 12.29%.

**Example A63:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-imidazol-4-yl-methyl)-benzamide Trifluoroacetic Acid Salt



10

Step 1. 1-Methyl-1H-imidazole-4-carboxylic acid amide




Prepared according to a procedure described by Piotrovskii, et al, *Chem.*

- 15 *Heterocycl. Compd. (Eng. Transl.)*, 26, 4, 407-409 (1990).

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  7.68 (s, 1H), 7.64 (s, 1H), 7.23 (bs, 1H), 7.04 (bs, 1H), 3.28 (s, 3H).

Step 2. C-1-Methyl-1H-imidazol-4-yl)-methylamine



- 20 To a solution of 1-methyl-1H-imidazole-4-carboxylic acid amide (0.80 g, 6.4 mmol) in THF (15 ml) at  $0^\circ\text{C}$  was added  $\text{LiAlH}_4$  (486 mg, 12.8 mmol). The mixture was refluxed for 24 h, cooled to  $0^\circ\text{C}$ , then carefully quenched with sat  $\text{NaHCO}_3$  (2 ml), diluted with ether (80 ml), and filtered through a pad of Celite. The filtrate was dried over  $\text{Na}_2\text{SO}_4$  and concentrated to give 476 mg of yellow oil in 67% yield and was used without any further purification.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.35 (s, 1H), 6.73 (s, 1H), 3.78 (s, 2H), 3.64 (s, 3H).


The title compound was prepared in a manner analogous to Step 3 of Method A, from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**) and C-1-methyl-1H-imidazol-4-yl)-methylamine. Purified via preparative HPLC.

5  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  11.07 (s, 1H), 8.99 (t, 1H,  $J=5.3$  Hz), 8.94 (s, 1H), 8.19 (bs, 2H), 7.88 (d, 2H,  $J=8.7$  Hz), 7.69 (d, 2H,  $J=8.7$  Hz), 7.56 (s, 1H), 7.54 (m, 1H), 7.21 (dd, 2H,  $J=8.0, 8.0$  Hz), 4.47 (d, 2H,  $J=5.3$  Hz), 3.83 (s, 3H).


ESMS ( $\text{M}+\text{H}^+$ ): 469.

Anal. Calcd. for  $\text{C}_{22}\text{H}_{18}\text{F}_2\text{N}_6\text{O}_2\text{S} \cdot 2.3 \text{ TFA} \cdot 1.0 \text{ H}_2\text{O}$ : C, 42.67%; H, 3.00%; N, 11.22%; S, 4.28%. Found: C, 42.40%; H, 3.14%; N, 11.17%; S, 4.30%.

**Example A64:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methylimidazol-5-yl-methyl)-benzamide.



15 Step 1. C-(3-Methyl-3H-imidazol-4-yl)-methylamine



Prepared in a manner analogous to Step 2 of Example A63, from 1-methylimidazole-5-carboxamide (Apollo Scientific, Ltd) to give a yellow oil in 86% yield, which was used without any further purification.

20  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.39 (s, 1H), 6.89 (s, 1H), 3.85 (d, 2H,  $J=0.6$  Hz), 3.66 (s, 3H).

The title compound was prepared in a manner analogous to Step 3 of Method A, from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**) and C-(3-methyl-3H-imidazol-4-yl)-methylamine. Purified via preparative HPLC.


25  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  11.00 (s, 1H), 8.93 (s, 1H), 8.85 (t, 1H,  $J=5.8$  Hz), 8.11 (bs, 2H), 7.79 (d, 2H,  $J=8.7$  Hz), 7.60 (d, 2H,  $J=8.7$  Hz), 7.51 (s, 1H), 7.47 (m, 1H), 7.14 (dd, 2H,  $J=7.8, 8.0$  Hz), 4.48 (d, 2H,  $J=5.5$  Hz), 3.80 (s, 3H).

ESMS ( $\text{M}+\text{H}^+$ ): 469.




Anal. Calcd. for  $C_{22}H_{18}F_2N_6O_2S \cdot 1.5 \text{ TFA} \cdot 1.0 \text{ H}_2\text{O}$ : C, 45.67%; H, 3.30%; N, 12.78%; S, 4.88%. Found: C, 45.66%; H, 3.44%; N, 12.96%; S, 4.89%.

**Example A65:** 2R-{4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoyl}-amino}-propionic Acid Trifluoroacetic Acid Salt



The starting material was prepared as follows:

*2R-{4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoyl}amino-propionic Acid tert-Butyl Ester*



10

Prepared in a manner similar to Step 3 in Method A from D-alanine tert-butyl ester hydrochloride (Novabiochem). Column chromatography with 10% MeOH/ $\text{CHCl}_3$  provided a yellow powder in 98% yield, which was used without any further purification.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.86 (d, 2H,  $J=8.9$  Hz), 7.74 (d, 2H,  $J=8.8$  Hz), 7.52-7.42 (m, 1H), 7.06 (dd, 2H,  $J=7.5, 7.5$  Hz), 2.98 (s, 3H), 1.48 (s, 9H).

15

The title Example was prepared as follows. A solution of 2R-{4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoyl}amino-propionic acid tert-butyl ester (350 mg, 0.696 mmol) in TFA (2 ml) stirred for 0.5 h. Ether was added to the solution until a yellow precipitate formed, which was filtered and dried under high vacuum to afford 150 mg of yellow powder in 49% yield.

20

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.89 (d, 2H,  $J=8.8$  Hz), 7.76 (d, 2H,  $J=8.7$  Hz), 7.58-7.46 (1H, m), 7.06 (dd, 2H,  $J=7.7, 8.1$  Hz), 1.54 (d, 3H,  $J=7.3$  Hz).


LC-MS ( $\text{M}+\text{H}^+$ ): 447; ( $\text{M}-\text{H}$ ): 445.

Anal. Calcd. for  $C_{20}H_{16}F_2N_4O_4S \cdot 1\text{H}_2\text{O} \cdot 0.5\text{TFA}$ : C, 48.37%; H, 3.58%; N, 10.74%; S,

6.15%. Found: C, 48.68%; H, 3.58%; N, 10.45%; S, 5.94%.


25

**Example A66:** 4-{4-Amino-5-(2-fluoro-benzoyl)-thiazol-2-ylamino}-N-dimethylamino-1R-(methyl-ethyl)-benzamide Trifluoroacetic Acid Salt




Starting materials were prepared as follows:

Step 1. *2-Bromo-2'-fluoro-acetophenone*.



5 The above intermediate was prepared in a manner similar to that for 2-bromo-2',6'-difluoro-acetophenone as described in Method A and used without any further purification. The  $^1\text{H}$  NMR spectrum matched that described previously in C. Giuseppe et. al. *J. Med. Chem.*, 20, 3763-3772 (1998).

Step 2. *4-[4-Amino-5-(2-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic Acid Ethyl Ester*




10

As described in Step 1 in Method A, 4-ethoxycarbonylphenyl isothiocyanate and 2-bromo-2'-fluoro-acetophenone provided a yellow solid in 99% yield, which was used without any further purification.

15  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  11.09 (s, 1H), 8.15 (br, 2H), 7.94 (d, 2H,  $J=8.6$  Hz), 7.75 (d, 2H,  $J=8.6$  Hz), 7.55 (d, 1H,  $J=7.2$  Hz), 7.50 (d, 1H,  $J=7.0$  Hz), 4.30 (q, 2H,  $J=7.1$  Hz), 1.31 (t, 3H,  $J=7.1$  Hz).

ESIMS ( $\text{M}+\text{H}^+$ ): 386.

Step 3. *4-[4-Amino-5-(2-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic Acid*



Prepared as described in Step 2 of Method A from 4-[4-amino-5-(2-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid ethyl ester to give a yellow solid in 90% yield, which was used without any further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 12.88 (br, 1H), 11.25 (s, 1H), 8.32 (br, 2H), 8.09 (d, 2H, J=8.7 Hz), 7.89 (d, 2H, J=8.7 Hz), 7.72 (d, 1H, J=7.1 Hz), 7.66 (d, 1H, J=6.7 Hz).  
ESIMS (M+H<sup>+</sup>): 358.


The title compound was prepared in a manner analogous to Step 3 in Method A from 4-[4-amino-5-(2-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid and (R)-N,N-dimethyl-propane-1,2-diamine dihydrochloride salt (from Step 3 of Example A40).

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.81 (s, 1H), 8.87 (br, 1H), 8.12 (d, 1H, J=8.6 Hz), 7.92 (br, 2H), 7.65 (d, 2H, J=8.7 Hz), 7.45 (d, 2H, J=8.7 Hz), 4.23 (sextet, 1H, J=6.3 Hz), 2.61 (d, 3H, J=4.7 Hz), 2.56 (d, 3H, J=4.7 Hz), 0.96 (d, 3H, J=6.7 Hz).

ESIMS (M+H<sup>+</sup>): 442.

Anal. Calcd. for C<sub>22</sub>H<sub>24</sub>FN<sub>5</sub>O<sub>2</sub>S • 0.9 H<sub>2</sub>O • 1.9 TFA: C, 45.95%; H, 4.14%; N, 10.38%; S, 4.75%. Found: C, 45.76%; H, 4.10%; N, 10.67%; S, 4.95%.

**Example A67:** 4-{4-Amino-5-(2-fluoro-benzoyl)-thiazol-2-ylamino}-N-(1-methyl-pyrrolidin-2S-ylmethyl)-benzamide Trifluoroacetic Acid Salt




The title compound was prepared in a manner analogous to Step 3 in Method A, from 4-[4-amino-5-(2-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (from Step 3 in Example A66) and (1-methyl-pyrrolidin-2S-yl)-methylamine (Sassaman, et al., *Bioorg. Med. Chem.*, 6, 1759-1766 (1998)), and purified via preparative HPLC.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.10 (s, 1H), 9.40 (br, 1H), 8.78 (t, 1H, J=5.6 Hz), 8.21 (br, 2H), 7.95 (d, 2H, J=8.7 Hz), 7.79 (d, 2H, J=8.7 Hz), 3.18 (m, 1H), 3.03 (d, 3H, J=4.7 Hz).  
ESIMS (M+H<sup>+</sup>): 454.


Anal. Calcd. for C<sub>23</sub>H<sub>24</sub>FN<sub>5</sub>O<sub>2</sub>S • 1.6 H<sub>2</sub>O • 0.2 CH<sub>3</sub>CN • 2.0 TFA: C, 45.80%; H, 4.18%; N, 10.14%; S, 4.46%. Found: C, 45.50%; H, 4.03%; N, 10.48%; S, 4.80%.

**Example A68:** 4-{4-Amino-5-(2,6-difluoro-4-methyl-benzoyl)-thiazol-2-ylamino}-N-(1-methyl-pyrrolidin-2S-ylmethyl)-benzamide



The starting materials were prepared as follows:


Step 1. (4-Bromo-2,6-difluoro-phenyl)trimethylsilane.



5 To diisopropylamine (1.73 ml, 12.4 mmol) in THF (30 ml) at  $-78^{\circ}\text{C}$  was added slowly n-butyllithium (7.73 ml of 1.6 M in hex). The mixture stirred at  $0^{\circ}\text{C}$  for 20 min and then was recooled to  $-100^{\circ}\text{C}$  with a liquid nitrogen/ether slush bath, whereupon 1-bromo-3,5-difluorobenzene (2.17 g, 11.2 mmol) was added at such a rate that the temperature never exceeded  $-90^{\circ}\text{C}$ . After 2h at  $-100^{\circ}\text{C}$ , chlorotrimethylsilane (1.86 ml, 14.6 mmol)  
 10 was added dropwise at such a rate that the temperature kept below  $-85^{\circ}\text{C}$ . The resultant mixture was allowed to warm to ambient temperature overnight, then quenched with water (2 ml), and extracted with ether. The separated organic layer was washed with brine and carefully concentrated below  $30^{\circ}\text{C}$  under reduced pressure on a rotary evaporator to give  
 15 2.97 g (100%) of a colorless oil, which was used in the next step without any further purification.

$^1\text{H}$  NMR:  $\delta$  7.00 (ddd, 2H, ,  $J=2.6, 2.6, 7.9$  Hz), 0.36 (dd, 9H,  $J=1.4, 1.4$  Hz).


Step 2. (2,6-Difluoro-4-methyl-phenyl)-trimethyl-silane



To a solution of (4-bromo-2,6-difluoro-phenyl)trimethylsilane (2.52 g, 9.50 mmol)  
 20 in ether (25 ml) at  $-60^{\circ}\text{C}$  was added n-butyllithium (7.1 ml of 1.6 M in THF) and the cooling bath removed. After a half-hour, the temperature rose to  $0^{\circ}\text{C}$ , recooled to  $-60^{\circ}\text{C}$ , and iodomethane (0.89 ml, 14.3 mmol) was added. The mixture was allowed to warm over one hour, quenched with water, and extracted with ether. The organic layer was washed  
 with water and brine, dried over dry  $\text{MgSO}_4$ , and concentrated to give a yellow oil, which  
 25 was used immediately without any further purification.

$^1\text{H}$  NMR:  $\delta$  6.61 (d, 2H,  $J=8.1$  Hz), 2.31(s, 3H), 0.34 (t, 9H,  $J=1.3$  Hz).


## Step 3. 2,6-Difluoro-4-methyl-acetophenone



According to a procedure described by Bennetau, et al., *Tetrahedron*, 49, 10843-10845 (1993), a mixture of  $\text{AlCl}_3$  (1.58 g, 11.9 mmol) in  $\text{CH}_2\text{Cl}_2$  (17 ml) was cooled to 0°C, and acetyl chloride (0.84 ml, 12 mmol) was added. The resultant suspension stirred at 0°C for 15 min, and a solution of (2,6-difluoro-4-methyl-phenyl)-trimethylsilane (9.50 mmol) in  $\text{CH}_2\text{Cl}_2$  (15 ml) was added dropwise. The mixture was allowed to warm over one hour, recooled to 0°C, quenched with 1N HCl, and extracted with ether. The organic layer was washed with 1N HCl and brine, dried over  $\text{MgSO}_4$ , and concentrated to give 1.56 g of a brown oil in 97% yield for two steps from 4-bromo-2,6-difluoro-phenyl)trimethylsilane, and was used without any further purification.

$^1\text{H}$  NMR:  $\delta$  6.76 (d, 2H,  $J=9.3$  Hz), 2.56 (t 3H,  $J=1.9$  Hz), 2.37 (s, 3H).


## Step 4. 2-Bromo-2',6'-difluoro-4'-methyl-acetophenone.



Prepared in a manner similar to that for 2-bromo-2',6'-difluoro-acetophenone in Method A, and used without any further purification.

$^1\text{H}$  NMR:  $\delta$  6.81 (d, 2H,  $J=9.4$  Hz), 4.35 (s 2H), 2.40 (s, 3H).

## Step 5. 4-[4-Amino-5-(2,6-difluoro-4-methyl-benzoyl)-thiazol-2-ylamino]-benzoic Acid



As described in Step 1 in Method A, 4-ethoxycarbonyl-phenylisothiocyanate and 2-bromo-2',6'-difluoro-4'-methyl-acetophenone provided a yellow solid in 92% crude yield, which displayed a MS consistent for desired 4-[4-amino-5-(2,6-difluoro-4-methyl-benzoyl)-thiazol-2-ylamino]-benzoic acid ethyl ester (ESIMS ( $\text{M}+\text{H}^+$ ): 418) and was treated as described in Step 2 in Method A to furnish a yellow solid in 88% yield, which was used without any further purification.

$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  12.77 (br, 1H), 11.17 (s, 1H), 8.20 (br, 2H), 7.92 (d, 2H,  $J=8.7$  Hz), 7.71 (d, 2H,  $J=8.7$  Hz), 7.04 (d, 2H,  $J=8.6$  Hz), 2.37 (s, 3H).

ESIMS ( $M+H^+$ ): 390.

The title compound was prepared in a manner analogous to Step 3 in Method A from 4-[4-amino-5-(2,6-difluoro-4-methyl-benzoyl)-thiazol-2-ylamino]-benzoic acid and (1-methyl-pyrrolidin-2S-yl)-methylamine (Sassaman, et al., *Bioorg. Med. Chem.*, 6, 1759-1766 (1998)).


$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  10.99 (br, 1H), 8.32 (t, 1H,  $J=5.9$  Hz), 8.20 (br, 2H), 7.83 (d, 2H,  $J=8.7$  Hz), 7.65 (d, 2H,  $J=8.7$  Hz), 7.04 (d, 2H,  $J=8.5$  Hz), 3.17 (m, 1H), 2.99 (m, 1H), 2.37 (s, 3H), 2.35 (s, 3H).

ESIMS ( $M+H^+$ ): 486.

Anal. Calcd. for  $C_{24}H_{25}F_2N_5O_2S \cdot 1.3$  MeOH: C, 57.64%; H, 5.77%; N, 13.28%; S, 6.08%.


Found: C, 57.59%; H, 5.44%; N, 12.90%; S, 6.22%.

- 15 **Example A69:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino-N-[1S-(1-methyl-pyrrolidin-2S-yl)-ethyl]-benzamide



The starting materials were prepared as follows:

Step 1. 2S-(1S-Amino-ethyl)-pyrrolidine-1-carboxylic Acid tert-Butyl Ester




To a solution of 2S-[1S-(benzyl-hydroxyamino)-ethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester (0.500 g, 1.56 mmol; Merino, et al., *Tetrahedron: Asymmetry*, 10, 1861-1865 (1999); Merino, et al., *Tetrahedron: Asymmetry*, 10, 1867-1871 (1999)) in MeOH (10 ml) was added 20% palladium hydroxide on activated charcoal (0.2 g). The resultant mixture stirred under a hydrogen balloon at ambient temperature for 2 days. The

catalyst was filtered off through a plug of Celite and rinsed with MeOH. The filtrate was concentrated *in vacuo* to afford 0.33 g of a colorless oil in 99% yield, which was used without further purification.

<sup>1</sup>H NMR: δ 3.71 (bs, 1H), 3.49 (s, 2H), 3.32-3.24 (m, 1H), 3.00 (m, 1H, J=6.6 Hz), 1.46 (s, 9H), 1.05 (d, 3H, J=6.5 Hz).

Step 2. (1*S*-Methyl-pyrrolidin-2*S*-yl)-ethylamine Hydrochloride



Prepared in a manner similar to Step 2 of Example A39: 2*S*-(1*S*-amino-ethyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (0.760 g, 3.54 mmol) was reduced with LiAlH<sub>4</sub> and the crude filtrate treated with 4*N* HCl in dioxane (1 ml). The resultant solution was concentrated *in vacuo* to provide 0.70 g of a gummy oil in 99% yield, which was used without further purification.

<sup>1</sup>H NMR: δ 3.94 (ddd, 1H, J=2.5, 7.1, 9.5 Hz), 3.74-3.68 (m, 1H), 3.42-3.32 (m, 1H), 2.79 (s, 3H), 2.28-2.18 (m, 1H), 1.42 (d, 3H, J=6.6 Hz).


The title Example was prepared in a manner similar to that for Step 3 in Method A from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (3) and (1*S*-methyl-pyrrolidin-2*S*-yl)-ethylamine (Sassaman, et al., *Bioorg. Med. Chem.*, 6, 1759-1766 (1998)).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.73 (d, 2H, J=8.7 Hz), 7.57 (d, 2H, J=8.5 Hz), 7.52-7.44 (m, 1H), 7.06 (dd, 2H, J=7.5, 8.3 Hz), 4.50-4.44 (m, 1H), 2.48 (s, 3H), 1.10 (d, 3H, J=6.3 Hz).

LC-MS (M+H<sup>+</sup>): 486; (M-H<sup>-</sup>): 484.

Anal. Calcd. for C<sub>24</sub>H<sub>25</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S•1.3H<sub>2</sub>O•0.2CHCl<sub>3</sub>: C, 54.55%; H, 5.03%; N, 13.14%; S, 6.02%. Found: C, 54.71%; H, 5.03%; N, 13.08%; S, 5.89%.

**Example A70:** 4-{4-Amino-5-[1-(3-methyl-thiophen-2-yl)-methanoyl]-thiazol-2-ylamino}-*N*-(1-methyl-pyrrolidin-2*S*-ylmethyl)-benzamide




The title Example was prepared in a manner similar to Step 3 in Method A from 4-{4-amino-5-[1-(3-methyl-thiophen-2-yl)-methanoyl]-thiazol-2-ylamino}-benzoic acid (Example A62 Step 2) and (R)-N,N-dimethyl-propane-1,2-diamine dihydrochloride salt (from Example A33).

- 5 <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.85 (d, 2H, J=8.8 Hz), 7.76 (d, 2H, J=8.8 Hz), 7.43 (d, 1H, J=5.0 Hz), 6.94 (d, 1H, J=5.0 Hz), 3.68 (1H, dd, J=4.3, 13.9 Hz), 2.40 (s, 3H).

Anal. Calcd. for C<sub>22</sub>H<sub>25</sub>N<sub>5</sub>O<sub>2</sub>S<sub>2</sub>•0.35Hexane•0.3CHCl<sub>3</sub>: C, 56.19%; H, 5.84%; N, 13.43%; S, 12.30%. Found: C, 56.54%; H, 5.84%; N, 13.79%; S, 11.92%.


LC-MS (M+H<sup>+</sup>): 456; (M-H): 454.

- 10 **Example A71:** 4-[4-Amino-5-(2-chloro-6-fluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-pyrrolidin-2S-ylmethyl)-benzamide



The starting materials were prepared as follows:

Step 1. 2-Bromo-1-(2-chloro-6-fluoro-phenyl)-ethanone




15

Prepared in a manner similar to that for 2-bromo-2',6'-difluoro-acetophenone in Method A, and used without any further purification.

<sup>1</sup>HNMR: δ 7.45-7.32 (m, 1H), 7.12 (d, 1H, J=8.8 Hz), 7.07 (dd, 1H, J=4.2, 8.7 Hz), 4.38 (s, 2H).

- 20 **Step 2. 4-[4-Amino-5-(2-chloro-6-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic Acid Ethyl Ester**




Prepared in a manner analogous to Step 1 of Method A and used without any further purification.



<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 8.00 (d, 2H, J=8.8 Hz), 7.77 (d, 2H, J=8.8 Hz), 7.50-7.42 (m, 1H), 7.34 (d, 1H, J=8.1 Hz), 7.18 (dd, 1H, J=8.3, 8.7 Hz), 4.36 (q, 1H, J=7.1 Hz), 1.40 (t, 3H, J=14.2 Hz).

Step 3. 4-[4-Amino-5-(2-chloro-6-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic Acid



5

Prepared in a manner similar to Step 2 of Method A and used without any further purification.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 8.06 (d, 2H, J=8.9 Hz), 7.82 (d, 2H, J=8.9 Hz), 7.52-7.44 (m, 1H), 7.38 (d, 1H, J=8.1 Hz), 7.24 (dd, J=8.5, 8.5 Hz).


10 The title Example was prepared in a manner similar to step 3 in Method A, from 4-[4-amino-5-(2-chloro-6-fluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid and (1-methylpyrrolidin-2S-yl)-methylamine (Sassaman, et al., *Bioorg. Med. Chem.*, 6, 1759-1766 (1998)).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.87 (d, 2H, J=8.8 Hz), 7.76 (d, 2H, J=8.8 Hz), 7.52-7.43 (m, 1H),  
15 7.35 (d, 1H, J=8.1 Hz), 7.21 (dd, 1H, J=7.9, 8.5 Hz), 3.71 (dd, 1H, J=4.0, 13.5 Hz), 3.18-3.08 (m, 1H), 2.65-2.56 (m, 1H), 2.50 (s, 3H), 2.36 (dd, 1H, J=9.0, 18.1 Hz), 2.08-1.98 (m, 1H).


LC-MS (M+H<sup>+</sup>): 488; (M-H): 486.

Anal. Calcd. for C<sub>23</sub>H<sub>23</sub>ClFN<sub>5</sub>O<sub>2</sub>S • 0.8H<sub>2</sub>O • 0.06CH<sub>2</sub>Cl<sub>2</sub>: C, 54.58%; H, 4.91%; N,  
20 13.80%; S, 6.32%; Cl, 7.82%. Found: C, 54.82%; H, 4.97%; N, 13.48%; S, 6.07%; Cl, 8.11%.

**Example A72:** 4-{5-(2-Acetylamino-benzoyl)-4-amino-thiazol-2-ylamino}-N-(1-methylpyrrolidin-2S-ylmethyl)-benzamide




25 Step 1. N-(2-Acetyl-phenyl)-acetamide




Acetic anhydride (10 ml) was added to a solution of 2'-aminoacetophenone (2.00 ml, 16.2 mmol) in acetic acid (10 ml) and heated at 70°C for a half-hour. Allowed to cool and quenched with ice water (150 ml). The resultant white solid was filtered, washed with ice-  
 5 water, dried under high vacuum to give 2.75 g of solid which displayed an NMR spectrum identical to that described in Adam, et. al, *J. Org. Chem*, 59, 2733-2739 (1994) and was used without any further purification.

Step 2. *N*-[2-(2-Bromo-acetyl)-phenyl]-acetamide



10 Made in a manner similar to that for 2-bromo-2',6'-difluoro-acetophenone in Method A, to provide an NMR which matched that described in Alkhathlan et. al, *Heterocycles*, 48, 641-656 (1998) and used without any further purification.

Step 3. 4-[5-(2-Acetyl-amino-benzoyl)-4-amino-thiazol-2-ylamino]-benzoic Acid Ethyl Ester




15 Prepared as described in Step 1 in Method A. 4-Ethoxycarbonylphenyl isothiocyanate and *N*-[2-(2-bromo-acetyl)-phenyl]-acetamide provided a yellow solid in 100% yield, which was used without any further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.12 (s, 1H), 9.89 (s, 1H), 8.26 (br, 2H), 8.01 (d, 1H, J=8.2 Hz), 7.99 (d, 2H, J=8.8 Hz), 7.77 (d, 2H, J=8.8 Hz), 7.58 (dd, 1H, J=1.2, 7.6 Hz), 7.49 (dt, 1H, J=1.2, 8.2  
 20 Hz), 7.18 (t, 1H, J=7.6 Hz), 4.30 (q, 2H, J=7.1 Hz), 2.03 (s, 3H), 1.32 (t, 3H, J=7.1 Hz).

ESIMS (M-H<sup>+</sup>): 423

## Step 4. 4-[5-(2-Acetylamino-benzoyl)-4-amino-thiazol-2-ylamino]-benzoic Acid



Prepared as described in Step 2 in Method A in 74% yield.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 12.66 (br, 1H), 11.05 (s, 1H), 8.23 (br, 2H), 7.98 (d, 1H, J=8.2 Hz),  
 5 7.89 (d, 2H, J=8.6 Hz), 7.71 (d, 2H, J=8.6 Hz), 7.54 (d, 1H, J=7.6 Hz), 7.40 (t, 1H, J=7.6 Hz),  
 7.19 (t, 1H, J=7.6 Hz), 1.99 (s, 3H).

ESIMS (M-H<sup>+</sup>): 395.


The title compound was prepared in a manner analogous to Step 3 in Method A.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.02 (br, 1H), 9.97 (s, 1H), 8.31 (s, 1H), 8.29 (br, 2H), 8.08 (d, 1H,  
 10 J=8.7 Hz), 7.89 (d, 2H, J=8.4 Hz), 7.74 (d, 2H, J=8.4 Hz), 7.63 (d, 1H, J=7.5 Hz), 7.48 (t, 1H,  
 J=7.5 Hz), 7.22 (t, 1H, J=7.5 Hz), 3.49 (m, 1H), 3.19 (m, 1H), 3.00 (m, 1H), 2.41 (m, 1H), 2.37  
 (s, 3H), 2.20 (m, 1H), 2.09 (s, 3H), 1.89 (m, 1H).


ESIMS (M+H<sup>+</sup>): 529.

Anal. Calcd. for C<sub>25</sub>H<sub>28</sub>N<sub>6</sub>O<sub>3</sub>S • 0.5 H<sub>2</sub>O: C, 58.52%; H, 6.23%; N, 15.75%; S, 6.01%. Found:  
 15 C, 58.35%; H, 5.89%; N, 15.79%; S, 6.03%.

**Example A73:** 4-{4-Amino-5-(2-methanesulfonyl-benzoyl)-thiazol-2-ylamino}-N-(1-methyl-pyrrolidin-2S-ylmethyl)-benzamide Trifluoroacetic Acid Salt



## Step 1. 1-(2-Methanesulfonyl-phenyl)-ethanone




Prepared as described in Binder et al, *Arch. Pharm. (Weinheim Ger.)*, 313, 587–602 (1980), and used without any further purification.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  8.09 (dd, 1H,  $J=1.2, 7.8$  Hz), 7.68 (m, 1H) 7.62 (m, 1H) 7.45 (dd, 1H,  $J=1.3, 7.4$  Hz), 3.24 (s, 3H), 2.65 (s, 3H).

5 ESIMS ( $\text{M}+\text{H}^+$ ): 199.


Step 2. 2-Bromo-1-(2-methanesulfonyl-phenyl)-ethanone



Prepared in a manner similar to that for 2-bromo-2',6'-difluoro-acetophenone in Method A and used without any further purification.

10  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  8.07 (dd, 1H,  $J=1.4, 7.7$  Hz), 7.72 (m, 1H) 7.68 (m, 1H) 7.53 (dd, 1H,  $J=1.4, 7.1$  Hz), 4.50 (s, 2H), 3.17 (s, 3H).

Step 3. 4-[4-Amino-5-(2-methanesulfonyl-benzoyl)-thiazol-2-ylamino]-benzoic Acid Ethyl Ester




15 Prepared as described for Step 1 in Method A, from 4-ethoxycarbonylphenyl-isothiocyanate and 2-bromo-1-(2-methanesulfonyl-phenyl)-ethanone to provide a yellow solid in 95% yield, which was used without any further purification.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  11.09 (s, 1H), 8.08 (br, 2H), 8.01 (d, 1H,  $J=7.0$  Hz), 7.93 (d, 2H,  $J=8.7$  Hz), 7.81 (dd, 1H,  $J=7.0, 7.3$  Hz), 7.73 (d, 2H,  $J=8.7$  Hz), 7.66 (dd 1H,  $J=7.0, 7.3$  Hz), 4.29 (q, 2H,  $J=7.1$  Hz), 3.37 (s, 3H), 1.31 (t, 3H,  $J=7.1$  Hz).

20

ESIMS ( $\text{M}-\text{H}^+$ ): 444.

Step 4. 4-[4-Amino-5-(2-methanesulfonyl-benzoyl)-thiazol-2-ylamino]-benzoic Acid



Prepared as described in Step 2 of Method A to afford a yellow solid in 87% yield, which was used without any further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 12.52 (br, 1H), 11.09 (s, 1H), 8.09 (br, 2H), 8.01 (d, 1H, J=7.4 Hz), 7.91 (d, 2H, J=8.7Hz), 7.81 (dd, 1H, J=7.3, 7.4 Hz), 7.73 (dd, 1H, J=7.3, 7.4 Hz), 7.71 (d, 2H, J=7.3, 7.4 Hz), 7.64 (d, 1H, J=7.3 Hz), 3.37 (s, 3H).

ESIMS (M-H<sup>+</sup>): 416.


The title compound was prepared in a manner analogous to Step 3 in Method A, from 4-[4-amino-5-(2-methanesulfonyl-benzoyl)-thiazol-2-ylamino]-benzoic acid and (1-methyl-pyrrolidin-2S-yl)-methylamine (Sassaman, et al., *Bioorg. Med. Chem.*, 6, 1759-1766 (1998)), and purified via preparative HPLC.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.03 (s, 1H), 9.36 (br, 1H), 8.71 (dd, 1H, J=5.3, 5.7 Hz), 8.07 (br, 2H), 8.01 (d, 1H, J=7.6 Hz), 7.87 (d, 2H, J=8.7Hz), 7.81 (dd, 1H, J=7.2, 7.6 Hz), 7.74 (d, 1H, J=7.2, 7.6 Hz), 7.69 (d, 2H, J=8.7 Hz), 7.64 (d, 1H, J=7.2 Hz), 3.37 (s, 3H), 3.10 (m, 1H), 2.94 (d, 2H, J=4.7), 2.15 (m, 1H), 1.19 (m, 3H).

ESIMS (M-H<sup>+</sup>): 512.


Anal. Calcd. for C<sub>24</sub>H<sub>27</sub>N<sub>5</sub>O<sub>4</sub>S<sub>2</sub> • 1.0 H<sub>2</sub>O • 1.2 TFA: C, 46.15%; H, 4.43%; N, 10.12%; S, 9.26%. Found: C, 46.31%; H, 4.41%; N, 10.39%; S, 9.48%.

**Example A74:** 4-{4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino}-N-(1,2-dimethyl-pyrrolidin-2S-ylmethyl)-benzamide Trifluoroacetic Acid Salt



The starting materials were prepared as follows:


Step 1. 2S-Methyl-pyrrolidine-1,2-dicarboxylic Acid 1-tert-Butyl Ester



To a solution of 2S-methyl-pyrrolidine-2-carboxylic acid hydrobromide (1.50 g, 11.6 mmol; Bachem) in a mixture of H<sub>2</sub>O (15 ml) and dioxane (15 ml) was added Et<sub>3</sub>N (3.6 ml, 26 mmol) and di-tert-butyl dicarbonate (5.57 g, 25.5 mmol). The resultant solution stirred for 5 h, diluted with H<sub>2</sub>O (50 ml), washed with Et<sub>2</sub>O (50 ml), acidified to pH 2 with

10% HCl, and extracted with 10% MeOH/CHCl<sub>3</sub> (2 X 100 ml). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated in vacuo to afford 1.1 g of white powder in 64% yield, which displayed an <sup>1</sup>H NMR that matched previously reported (Khalil, et al., *Tetrahedron Lett.*, 37, 3441-3444 (1996)) and was used without any further purification.


*2S-Carbamoyl-2-methyl-pyrrolidine-1-carboxylic Acid tert-Butyl Ester*



To a solution of 2S-methyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester (1.00 g, 6.74 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) was sequentially added 1,1'-carbonyldiimidazole (1.20 g, 7.42 mmol) and N-hydroxysuccinimide (0.930 g, 8.08 mmol). The resultant solution was stirred for 4 h, concentrated *in vacuo*, diluted with dioxane, and treated with 58% NH<sub>4</sub>OH (5 ml). After 2 days, diluted with EtOAc (100 ml). The organic layer was washed with H<sub>2</sub>O (50 ml) and sat. NaHCO<sub>3</sub> (50 ml), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to afford a white solid in 27% yield, which was used without further purification.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 3.72-3.62 (m, 1H), 3.58-3.52 (m, 1H), 2.32-2.23 (m, 1H), 1.60 (s, 3H), 1.53 (s, 9H).

*C-(1,2S-Dimethyl-pyrrolidin-2-yl)-methylamine*



Prepared in a manner similar to Step 2 in Example A39 and used without any further purification.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 2.78 (s, 2H), 2.50 (s, 3H), 1.25 (s, 3H).

The title compound was prepared in a manner similar to Step 3 in Method A, from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (3) and C-(1,2S-dimethyl-pyrrolidin-2-yl)-methylamine and purified via preparative HPLC.


<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.68 (d, 2H, J=8.7 Hz), 7.45 (dd, 2H, J=1.9, 8.7 Hz), 6.97 (dd, 2H, J=7.7, 8.1 Hz), 4.00 (d, 0.7H, J=15.1 Hz), 3.48 (d, 1.3H, J=15.2 Hz), 2.98 (s, 2.1H), 2.68 (s, 0.9H), 1.54 (s, 0.8H), 1.40 (s, 2.2H).

LC-MS (M+H<sup>+</sup>): 486; (M-H<sup>-</sup>): 484.

Anal. Calcd. for  $C_{24}H_{25}F_2N_5O_2S \cdot 1.7TFA$ : C, 48.44%; H, 3.96%; N, 10.31%; S, 4.72%.


Found: C, 48.57%; H, 4.11%; N, 10.39%; S, 4.82%.

**Example A75:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(1-methyl-azetidin-3-ylmethyl)-benzamide



The starting materials were prepared as follows:


**Step 1. Azetidine-1,3-dicarboxylic Acid Mono tert-Butyl Ester**



Prepared in a manner analogous to Step 1 in Example A74 for 2S-methyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester, from azetidine-3-carboxylic acid and used without further purification.

$^1H$  NMR ( $CD_3OD$ ):  $\delta$  4.18-4.00 (m, 4H), 3.42-3.38 (m, 1H), 1.48 (s, 9H).

**Step 2. 3-Carbamoyl-azetidine-1-carboxylic Acid tert-Butyl Ester**



Prepared in a manner analogous to Step 2 in Example A74 for 2S-carbamoyl-2-methyl-pyrrolidine-1-carboxylic acid tert-butyl ester, from azetidine-1,3-dicarboxylic acid mono tert-butyl ester and used without further purification.

$^1H$  NMR ( $CD_3OD$ ):  $\delta$  4.09-3.92 (m, 4H), 1.45 (s, 9H).


The title compound was prepared from 3-carbamoyl-azetidine-1-carboxylic acid tert-butyl ester after 1) reduction in a manner similar to similar to Step 2 in Example A39 and 2) subsequent coupling with 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (3) in a manner similar to Step 3 in Method A.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.82 (d, 2H,  $J=8.8$  Hz), 7.74 (d, 2H,  $J=8.8$  Hz), 7.52-7.42 (m, 1H), 7.06 (dd, 2H,  $J=7.7, 8.1$  Hz), 3.52 (dd, 4H,  $J=6.8, 12.7$  Hz), 3.10 (dd, 2H,  $J=7.1, 8.1$  Hz), 2.84-2.74 (m, 1H), 2.36 (s, 1H).

LC-MS ( $\text{M}+\text{H}^+$ ): 458.


- 5     Anal. Calcd. for  $\text{C}_{22}\text{H}_{21}\text{F}_2\text{N}_5\text{O}_2\text{S}\cdot 1.0\text{H}_2\text{O}\cdot 0.3\text{CHCl}_3$ : C, 52.38%; H, 4.59%; N, 13.70%; S, 6.27%. Found: C, 52.74%; H, 4.85%; N, 13.58%; S, 6.06%.

**Example A76:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-methyl-2-aza-bicyclo[2.2.1]hept-3-endo-ylmethyl)-benzamide



- 10     The starting materials were prepared as follows:


Step 1. 3-endo-2-(tert-Butoxycarbonyl)-2-azabicyclo[2.2.1]hept-5-ene-3-carboxylic Acid Ethyl Ester.



- 15     Prepared in a manner similar to Step 1 in Example A74, from 3-endo-2-azabicyclo[2.2.1]hept-5-ene-3-carboxylic acid ethyl ester (Hursthouse, et al, *J. Chem. Soc. Perkin Trans. 1*, 2419-2425 (1995)), and used without further purification.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  6.58 (bs, 1H), 6.10 (bs, 1H), 4.90 (bs, 1H), 4.32 (bs, 1H), 4.20-4.08 (q, 2H,  $J=9.0$  Hz), 3.48 (bs, 1H), 1.45 (s, 9H), 1.25 (t, 3H,  $J=9.0$  Hz).

Step 2. 3-endo-2-(tert-Butoxycarbonyl)-2-azabicyclo[2.2.1]heptane-3-carboxylic Acid.



20

According to the conditions described in Hursthouse, et al, *J. Chem. Soc. Perkin Trans. 1*, 2419-2425 (1995), to a solution of 3-endo-2-(tert-butoxycarbonyl)-2-azabicyclo[2.2.1]hept-5-ene-3-carboxylic acid ethyl ester (3.74 g, 14.0 mmol) in MeOH (13.8 ml) at  $0^\circ\text{C}$  was added 2.5 N NaOH (5.9 ml). Allowed to warm to ambient




temperature. After 2 days, the MeOH was removed under reduced pressure, the concentrate was washed with EtOAc (2 x 20 ml; discarded), adjusted to pH3 with 10% citric acid, and extracted with EtOAc (2 x 30 ml). The acidic extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated in vacuo to give 2.04 g of a viscous oil in 61% yield, which was used without any further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 6.42 (bs, 2H), 4.88 (bs, 1H), 4.38 (d, 1H, J= 3.0 Hz), 3.62 (bd, 1H), 1.72 (dd, 2H, J=9.0, 9.0 Hz), 1.52 (s, 9H).

Anal. Calcd. for C<sub>12</sub>H<sub>17</sub>NO<sub>4</sub>: C, 60.24%; H, 7.16%; N, 5.85. Found: C, 59.80%; H, 7.22%; N, 5.76.


10 Step 3. 3-endo-2-(tert-Butoxycarbonyl)-2-azabicyclo[2.2.1]heptane-3-carboxylic Acid



According to the conditions carefully defined in Alonso, et al, *J. Org. Chem.*, 64, 2276-2280 (1999), a mixture of 3-endo-2-(tert-butoxycarbonyl)-2-azabicyclo[2.2.1]hept-5-ene-3-carboxylic acid (1.00 g, 4.20 mmol) in ethanol (100 ml) and 10% Pd/C (100 mg) stirred under a balloon of hydrogen for 3.5 h at ambient temperature. The catalyst was filtered off and filtrate concentrated in vacuo to give 988 mg of a solid in 99% yield, which was used without any further purification.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 4.36-4.26 (m, 1H), 4.18 (d, 1H, J=3.7 Hz), 2.80 (bs, 1H), 1.84-1.50 (m, 6H), 1.47 (s, 3H), 1.42 (s, 6H).

20 Step 4. 3-endo-Carbamoyl-2-aza-bicyclo[2.2.1]heptane-2-carboxylic Acid tert-Butyl Ester



Prepared in manner analogous to Step 2 in Example A74 for 2S-carbamoyl 2-methyl-pyrrolidine-1-carboxylic acid tert-butyl ester, from 2-aza-bicyclo[2.2.1]heptane-2,3-endo-dicarboxylic acid 2-tert-butyl ester. Used without any further purification.

25 <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 4.38 (bs, 1H), 4.10 (d, 1H, J=3.8 Hz), 2.80 (bs, 1H), 1.45 (s, 9H).


The title compound was prepared in a manner similar to that of Example A75, originating from 3-endo-carbamoyl-2-aza-bicyclo[2.2.1]heptane-2-carboxylic acid tert-butyl ester.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.76 (d, 2H,  $J=8.5$  Hz), 7.54 (d, 2H,  $J=8.7$  Hz), 7.08 (dd, 2H,  $J=7.6$ , 8.1 Hz), 2.55 (s, 3H).

LC-MS ( $\text{M}+\text{H}^+$ ): 498; ( $\text{M}-\text{H}$ ): 496.

Anal. Calcd. for  $\text{C}_{25}\text{H}_{25}\text{F}_2\text{N}_5\text{O}_2\text{S}\cdot 1.3\text{H}_2\text{O}\cdot 0.1\text{CHCl}_3$ : C, 56.57%; H, 5.24%; N, 13.14%; S, 6.02%. Found: C, 56.97%; H, 5.31%; N, 13.07%; S, 5.65%.

**Example A77:** 4-[[4-Amino-5-(2,6-difluorobenzoyl)-1,3-thiazol-2-yl]amino]-N-[[1-(dimethylamino)cyclopentyl]methyl]benzamide



- 10 The title compound was prepared in a manner similar to Step 3 in Method A; from coupling 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (3) and 1-(aminomethyl)-N,N-dimethylcyclopentanamine (Yang et al., *Eur. J. Med. Chem.* **31**, 231-239 (1996)) to afford a yellow amorphous powder in 62% yield.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  10.71 (bs, 1H), 8.01 (bs, 2H), 7.94 (bs, 1H), 7.79 (d, 2H,  $J=8.7$  Hz),

- 15 7.64 (d, 2H,  $J=8.7$  Hz), 7.54 (ddd, 1H,  $J=6.4$ , 8.4, 15.1 Hz), 7.20 (t, 2H,  $J=7.7$  Hz), 3.38 (d, 2H,  $J=5.8$  Hz), 2.26 (s, 6H), 1.57 (bs, 8H).


LCESIMS ( $\text{M}+\text{H}^+$ ): 500.15.

Anal. Calcd. for  $\text{C}_{25}\text{H}_{27}\text{F}_2\text{N}_5\text{O}_2\text{S}\cdot 0.41\text{H}_2\text{O}$ : C, 59.23%; H, 5.53%; N, 13.81%; S, 6.33%.

Found: C, 59.50%; H, 5.37%; N, 13.41%; S, 5.96%.

20

**Example A78:** 4-[[4-Amino-5-(2,6-difluorobenzoyl)-1,3-thiazol-2-yl]amino]-N-[[1-(dimethylamino)cyclobutyl]methyl]benzamide



The title compound was prepared in a manner similar to Step 3 in Method A; from coupling 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**3**) and 1-(aminomethyl)-N,N-dimethylcyclobutanamine (Yang et al., *Eur. J. Med. Chem.* **31**, 231-239 (1996)) to afford a yellow amorphous powder in 61% yield.


<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.05 (bs, 1H), 8.17 (bs, 1H), 7.73 (s, 3H), 7.82 (d, 2H, J=8.7 Hz), 7.64 (d, 2H, J=8.7 Hz), 7.54 (ddd, 1H, J=6.6, 8.3, 15.1 Hz), 7.20 (t, 2H, J=8.1 Hz), 3.49 (d, 2H, J=6.0 Hz), 2.19 (s, 6H), 1.85-1.95 (m, 4H), 1.62 (quintet, 2H, J=7.3 Hz).

LCESIMS (M+H<sup>+</sup>): 486.20.

Anal. Calcd. for C<sub>23</sub>H<sub>25</sub>F<sub>2</sub>N<sub>3</sub>O<sub>2</sub>S • 0.3 H<sub>2</sub>O: C, 58.71%; H, 5.26%; N, 14.26%; S, 6.53%.

Found: C, 58.99%; H, 5.41%; N, 14.20%; S, 6.49%.

#### Method B



#### Step 1. 4-Isothiocyanato-benzoic Acid tert-Butyl Ester (**6**)

tert-Butyl 4-aminobenzoate (2.39 g, 12.3 mmol; Fluka) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 ml), and cooled to 0°C. Thiophosgene (1.87 ml, 24.7 mmol) was added dropwise over 15 minutes. The resultant solution was allowed to warm to room temperature and stirred for 3 hours. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (200 ml), washed with saturated aqueous NaHCO<sub>3</sub> solution, brine, and dried over MgSO<sub>4</sub>, filtered and concentrated to a syrup. Chromatography on silica (hexane/ethyl acetate = 3/1) afforded 2.52 g of desired product as a yellow solid in 86 % yield, which was used without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.98 (d, 2H, J=8.8 Hz), 7.22 (d, 2H, J=8.8 Hz), 1.60 (s, 9H).

IR (KBr): 2226 cm<sup>-1</sup>.

Step 2. 4-[4-Amino-5-(2,6-dichlorobenzoyl)-thiazol-2-ylamino]-benzoic Acid *tert*-Butyl Ester (**7**)

The title compound was prepared from 4-isothiocyanato-benzoic acid *tert*-butyl ester (**6**) and 2-bromo-2',6'-dichloro-acetophenone in a manner similar to Step 1 in Method A and used without further purification.

- 5 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.18 (bs, 1H), 8.20 (br, 1H), 7.90 (d, 2H, J=8.8 Hz), 7.72 (d, 2H, J=8.8 Hz), 7.54 (m, 1H), 7.20 (t, 2H, J=8.7 Hz), 1.52 (s, 9H).


Step 3. 4-[4-Amino-5-(2,6-dichloro-benzoyl)-thiazol-2-ylamino]-benzoic Acid (**8**)

- To a solution of 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid *tert*-butyl ester (**7**, 1.25 g, 2.9 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (14 ml), trifluoroacetic acid (6 ml) was added. The  
10 reaction solution was stirred for 1 hour. Solvent was evaporated and a solution of resultant residue in ethyl acetate was washed with brine, dried with MgSO<sub>4</sub>, filtered and concentrated to afford 0.98 g of desired product as a light yellow solid in 90% yield, which was used without further purification.

- <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.18 (bs, 1H), 8.20 (br, 1H), 7.90 (d, 2H, J=8.8 Hz), 7.72 (d, 2H, J=8.8  
15 Hz), 7.54 (m, 1H), 7.20 (t, 2H, J=8.7 Hz).


Step 4. The following Examples B1 to B5 were prepared from 4-[4-amino-5-(2,6-dichloro-benzoyl)-thiazol-2-ylamino]-benzoic acid (**8**) and corresponding amines (R-NH<sub>2</sub>) in a manner similar to Step 3 in Method A.

- 20 **Example B1:** 4-[4-Amino-5-(2,6-dichloro-benzoyl)-thiazol-2-ylamino]-N-carbamoylmethyl-benzamide.



- <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.02 (s, 1H), 8.58 (t, 1H, J=5.9 Hz), 8.18 (bs, 2H), 7.88 (d, 2H, J=8.8 Hz),  
7.67 (d, 2H, J=8.8 Hz), 7.58-7.45 (m, 3H), 7.35 (br, 1H), 7.02 (br, 1H), 3.80 (d, 2H, J=5.9 Hz).  
25 HRFABMS: Calcd for C<sub>19</sub>H<sub>15</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>3</sub>Na (M+Na<sup>+</sup>): 486.0170. Found: 486.0183.


**Example B2:** 4-[4-Amino-5-(2,6-dichloro-benzoyl)-thiazol-2-ylamino]-N-(2-hydroxy-ethyl)-benzamide.



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.02 (s, 1H), 8.35 (t, 1H, J=5.5 Hz), 8.20 (bs, 2H), 7.88 (d, 2H, J=8.7 Hz), 7.67 (d, 2H, J=8.7 Hz), 7.59-7.42 (m, 3H), 4.71 (t, 1H, J=5.6 Hz), 3.56-3.45 (m, 2H), 3.34-3.25 (m, 2H).

5 HRFABMS: Calcd for C<sub>19</sub>H<sub>16</sub>Cl<sub>2</sub>N<sub>4</sub>O<sub>3</sub>SNa (M+Na<sup>+</sup>): 473.0218. Found: 473.0229.


**Example B3:** 4-[4-Amino-5-(2,6-dichloro-benzoyl)-thiazol-2-ylamino]-N-(2,3-dihydroxy-propyl)-benzamide



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.95 (s, 1H), 8.25 (t, 1H, J=5.5 Hz), 8.18 (bs, 2H), 7.82 (d, 2H, J=8.7 Hz), 7.58 (d, 2H, J=8.7 Hz), 7.52-7.38 (m, 3H), 4.78 (d, 1H, J=6.0 Hz), 4.49 (t, 1H, J=6.0 Hz), 3.58 (m, 1H), 3.39-3.21 (m, 3H), 3.14 (m, 2H).

HRFABMS: Calcd for C<sub>20</sub>H<sub>18</sub>Cl<sub>2</sub>N<sub>4</sub>O<sub>4</sub>SNa (M+Na<sup>+</sup>): 503.0324. Found: 503.0336.

**Example B4:** 4-[4-Amino-5-(2,6-dichloro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-ethyl)-benzamide.




15

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.00 (s, 1H), 8.31 (br, 1H), 8.18 (bs, 1H), 7.82 (d, 2H, J=8.8 Hz), 7.65 (d, 2H, J=8.8 Hz), 7.59-7.43 (m, 3H), 3.35 (t, 2H, J=6.9 Hz), 2.44 (t, 2H, J=6.9 Hz), 2.20 (s, 6H).

HRFABMS: Calcd for C<sub>21</sub>H<sub>21</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>2</sub>SNa (M+Na<sup>+</sup>): 500.0691. Found: 500.0671.

**Example B5:** 4-[4-Amino-5-(2,6-dichloro-benzoyl)-thiazol-2-ylamino]-N-[2-(2-hydroxy-ethoxy)-ethyl]-benzamide.


20



$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  11.20 (s, 1H), 8.60 (br, 1H), 8.35 (bs, 2H), 8.04 (d, 2H,  $J=8.8$  Hz), 7.85 (d, 2H,  $J=8.8$  Hz), 7.75-7.64 (m, 3H), 4.79 (t, 1H,  $J=2.7$  Hz), 3.77-3.57 (m, 8H).

HRFABMS: Calcd for  $\text{C}_{21}\text{H}_{21}\text{Cl}_2\text{N}_5\text{O}_2\text{SNa}$  ( $\text{M}+\text{Na}^+$ ): 517.0480. Found: 517.0488.

## 5 Method C



### Step 1. 2-Chloro-4-isothiocyanato-benzoic Acid (**11**)


The title compound was prepared from 4-amino-2-chlorobenzoic acid in a manner similar to that of Step 1 in Method B, and used without further purification.

$^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  7.62 (s, 1H,  $J=8.4$  Hz), 7.47 (d, 1H,  $J=2.0$  Hz), 7.23 (dd, 1H,  $J=2.0$ , 8.4 Hz).

### Step 2. 4-[4-Amino-5-(2,6-difluorobenzoyl)-thiazol-2-ylamino]-2-chlorobenzoic Acid (**12**)

The title compound was prepared from 2-chloro-4-isothiocyanato-benzoic acid (**11**) and 2-bromo-2',6'-difluoroacetophenone in a manner similar to Step 1 in Method A to give a yellow solid in 45% yield.

Step 3. **Example C1:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-2-chloro-N-(2-dimethylamino-ethyl)-benzamide.




The title compound was prepared from 4-{4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino}-2-chloro-benzoic acid (**12**) and N, N-dimethyl-ethylene-diamine in a manner similar to Step 3 in Method A.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.99 (d, 1H, J=1.9 Hz), 7.58-7.43 (m, 3H), 7.06 (t, 2H, J=7.8 Hz), 3.53 (t, 2H, J=6.9 Hz), 2.66 (t, 2H, J=6.8 Hz), 2.39 (s, 6H).

FABMS (MH<sup>+</sup>): 480; (M-H<sup>-</sup>): 478.

#### Method D



#### Step 1. 2-Hydroxy-4-isothiocyanato-benzoic Acid Phenyl Ester (**15**)

The title compound was prepared from 4-amino-2-hydroxy-benzoic acid phenyl ester (**14**) in a manner similar to Step 1 in Method B, and used without further purification.


<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 10.64 (s, 1H), 8.04 (s, 1H, J=8.5Hz), 7.46 (m, 2H), 7.32 (m, 1H), 7.20 (m, 2H), 6.84 (m, 2H).

#### Step 2. 4-{4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino}-2-hydroxy-benzoic Acid Phenyl Ester (**16**)

The title compound was prepared from 2-hydroxy-4-isothiocyanato-benzoic acid phenyl ester (**15**) and 2-bromo-2',6'-difluoro-acetophenone in a manner similar to Step 1 in Method A, and used without further purification.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 8.04 (s, 1H, J=8.5 Hz), 7.46 (m, 2H), 7.32 (m, 1H), 7.25 (m, 2H), 7.10 (m, 3H), 6.76 (m, 2H).

Step 3. **Example D1:** 4-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-2-hydroxy-N-(2-phenylamino-ethyl)-benzamide




A solution of 4-[4-amino-5-(2,6-difluorobenzoyl)-thiazol-2-ylamino]-2-hydroxy-benzoic acid phenyl ester (**16**) and N-phenyl-ethylene-diamine (5 eq) in DMF was heated at 70°C for 1 hour. DMF was removed under reduced pressure and the solution of resultant residue in ethyl acetate was washed with saturated aqueous NaHCO<sub>3</sub> solution, brine, dried with MgSO<sub>4</sub>, filtered and concentrated. The product was purified by HPLC.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 12.94 (s, 1H), 10.99 (s, 1H), 8.81 (br, 1H), 8.22 (bs, 2H), 7.85 (d, 2H, J=8.8 Hz), 7.57 (m, 1H), 7.31 (s, 1H), 7.21 (t, 2H, J=7.9 Hz), 7.09 (t, 2H, J=7.8 Hz), 6.98 (d, 1H, J=8.5 Hz), 6.62 (d, 2H, J=7.8 Hz), 6.53 (t, 1H, J=3.5 Hz), 5.73 (br, 1H), 3.45 (m, 2H), 3.22 (m, 2H).

HRFABMS: Calcd. For C<sub>25</sub>H<sub>21</sub>F<sub>2</sub>N<sub>3</sub>O<sub>3</sub>S (M+H<sup>+</sup>): 510.1411. Found: 510.1422.

**Example D2:** 4-[4-Amino-5-(2, 6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-dimethylamino-1RS-methyl-ethyl)-2-hydroxy-benzamide




The title compound was prepared from 4-[4-amino-5-[1-(2,6-difluoro-phenyl)-methanoyl]-thiazol-2-ylamino]-2-hydroxy-benzoic acid phenyl ester (**16**) and N<sup>1</sup>N<sup>1</sup>-dimethyl-propane-1,2-diamine in a manner similar to Step 3 in Method D for Example D1.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.77(d, 1H, J=8.7 Hz), 7.47 (m, 1H), 7.31 (d, 1H, J=2.1 Hz), 7.09 (t, 2H, J=10.9 Hz), 6.97 (dd, 1H, J=2.1, 8.7 Hz), 4.36 (m, 1H), 2.81 (m, 1H), 2.50 (m, 1H), 2.42 (s, 6H), 1.25 (d, 3H, J=8.7 Hz).

HRFABMS: Calcd. For C<sub>25</sub>H<sub>21</sub>F<sub>2</sub>N<sub>3</sub>O<sub>3</sub>S (M+H<sup>+</sup>): 476.1568. Found: 476.1564.



## Method E

Step 1. 2-(2-Dimethylamino-ethyl)-5-nitro-isoindole-1,3-dione (**19**)

A reaction solution of 4-nitro-phthalic anhydride (**18**; 0.96 g, 5 mmol) and N, N-dimethylethylenediamine (0.5 g, 5.5 mmol) in toluene (50 ml) was refluxed for four hours. The reaction solution was diluted with ethyl acetate and washed with 0.1 N NaOH, brine, dried with MgSO<sub>4</sub>, filtered and concentrated to give 0.7 g of desired product in 53% yield, which was used without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.66 (d, 1H, J=1.6 Hz), 8.58 (dd, 1H, J=1.6, 6.7 Hz), 8.02 (d, 1H, J=6.7 Hz), 3.88 (t, 2H, J=7.2 Hz), 2.64 (t, 2H, J=7.2 Hz), 2.28 (s, 6H).

Step 2. 5-Amino-2-(2-dimethylamino-ethyl)-isoindole-1,3-dione (**20**)


A solution of 2-(2-dimethylamino-ethyl)-5-nitro-isoindole-1,3-dione (**19**; 0.70 g, 2.6 mmol) in methanol (150 ml) and concentrated HCl (2 ml) was hydrogenated on 10% Pd/C (0.5 g) at 20 psi for 2 hours. The catalyst was filtered off and the filtrate was concentrated to give the desired product as a hydrochloride salt, which was used without further purification.

Step 3. 2-(2-Dimethylamino-ethyl)-5-isothiocyanato-isoindole-1,3-dione (**21**)

The title compound was prepared from 5-amino-2-(2-dimethylamino-ethyl)-isoindole-1,3-dione (**20**) in a manner similar to Step 1 in Method B, and used without further purification.

Step 4. This step is carried out in a manner similar to Step 1 in Method A.

**Example E1:** 5-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-2-(2-dimethylamino-ethyl)-isoindole-1,3-dione




The title compound was prepared from 2-(2-dimethylamino-ethyl)-5-isothiocyanato-isoindole-1,3-dione (**21**) and 2-bromo-2',6'-difluoro-acetophenone in a manner similar to Step 1 in Method A.

- 5 <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.86 (s, 1H), 7.68 (q, 2H, J=5.0 Hz), 7.34 (m, 1H), 6.90 (t, 2H, J=8.7 Hz), 3.88 (t, 2H, J=5.2 Hz), 2.64 (t, 2H, J=5.2 Hz), 2.28 (s, 6H).

HRFABMS: Calcd for C<sub>23</sub>H<sub>19</sub>F<sub>2</sub>N<sub>5</sub>O<sub>3</sub>S (M+H<sup>+</sup>): 472.1255. Found: 472.1244.

#### Method F



#### Step 1. 3-Isothiocyanato-benzoic Acid tert-Butyl Ester (**24**)

- 10 The title compound was prepared from t-butyl-3-amino-benzoate (**23**) in a manner similar to Step 1 in Method B, and used without further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 8.08 (s, 1H), 7.94 (d, 1H, J=7.7 Hz), 7.60 (d, 1H, J=8.5 Hz), 7.52 (m, 1H), 1.55 (s, 9H).

#### Step 2. 3-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamin]-benzoic Acid tert-Butyl Ester (**25**)

- 15 The title compound was prepared from t-butyl-3-isothiocyanato-benzoate (**24**) and 2-bromo-2',6'-difluoro-acetophenone in a manner similar to Step 1 in Method A, and used without further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 11.02 (s, 1H), 8.15 (br, 2H), 8.08 (s, 1H), 7.94 (d, 1H, J=7.7 Hz), 7.60 (d, 1H, J=8.5 Hz), 7.52 (m, 3H), 7.20 (t, 2H, J=8.7 Hz), 1.55 (s, 9H).


- 20 Step 3. 3-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic Acid (**26**)

The title compound was prepared from 3-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid tert-butyl ester (**25**) in a manner similar to Step 3 in Method B, and used without further purification.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 13.00 (br, 1H), 11.02 (s, 1H), 8.15 (br, 2H), 8.08 (s, 1H), 7.94 (d, 1H, J=7.7 Hz), 7.60 (d, 1H, J=8.5 Hz), 7.52(m, 3H), 7.20 (t, 2H, J=8.7 Hz).

Step 4. This step was carried out in a manner similar to Step 3 in Method A.

**Example F1:** 3-[4-Amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-N-(2-methylamino-ethyl)-benzamide



The title compound was prepared from 3-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid **26** and N<sup>1</sup>, N<sup>1</sup>-dimethyl-ethylenediamine in a manner similar to Step 3 in Method A.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.98 (s, 1H), 8.48 (br, 1H), 8.21 (bs, 2H), 7.92-7.81 (m, 2H), 7.58-7.42 (m, 3H), 7.21 (t, 2H, J=7.9 Hz), 3.43 (m, 2H), 2.65 (m, 2H), 2.38 (s, 6H).

HRFABMS: Calcd for C<sub>21</sub>H<sub>21</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S (M+H<sup>+</sup>): 446.1462. Found: 446.1473.

#### Examples G1 - G396

All compounds from Example G1 to G396, formulae of which are shown in Table 2 below, were combinatorially synthesized by the general method described as Step 3 in Method A from 4-[4-amino-5-(2,6-difluoro-benzoyl)-thiazol-2-ylamino]-benzoic acid **3** and corresponding amines (R-NH<sub>2</sub>), except a stock solution of (**3**) in 5% DIEA/DMF was distributed into 96 deep-well plates such that each well contained 10 μmol of material. Then, 10 μmol of corresponding amine and 10 μmol of HATU [O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate] in DMF were added into individual wells of each plate. The reaction mixture was shaken at room temperature for 16 hours. The reaction solvent was removed and the resultant combinatorial compounds were submitted for bioassays without further purification.

**BIOCHEMICAL AND BIOLOGICAL EVALUATION:**

Cyclin-dependent kinase activity was measured by quantifying the enzyme-catalyzed, time-dependent incorporation of radioactive phosphate from [ $^{32}$ P]ATP or [ $^{33}$ P]ATP into a protein substrate. Unless noted otherwise, assays were performed in 96-well plates in a total volume of 50  $\mu$ L, in the presence of 10 mM HEPES (N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid]) (pH 7.4), 10 mM  $\text{MgCl}_2$ , 25  $\mu$ M adenosine triphosphate (ATP), 1 mg/mL ovalbumin, 5  $\mu$ g/mL leupeptin, 1 mM dithiothreitol, 10 mM  $\beta$ -glycerophosphate, 0.1 mM sodium vanadate, 1 mM sodium fluoride, 2.5 mM ethylene glycol-bis( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), 2% (v/v) dimethylsulfoxide, and 0.03 - 0.4  $\mu$ Ci [ $^{32/33}$ P]ATP per reaction. Reactions were initiated with enzyme, incubated at 30°C, and terminated after 20 minutes by the addition of ethylenediaminetetraacetic acid (EDTA) to 250 mM. The phosphorylated substrate was then captured on a nitrocellulose or phosphocellulose membrane using a 96-well filtration manifold, and unincorporated radioactivity was removed by repeated washing with 0.85% phosphoric acid. Radioactivity was quantified by exposing the dried membranes to a phosphorimager.

Compounds from combinatorial libraries were screened from 96-well plates for % inhibition of CDK activity at 30 nM theoretical compound concentration. Inhibition was measured relative to control wells that contained all reaction components including 2% (v/v) DMSO but no compound, after subtraction of background radioactivity measured in the absence of enzyme. Apparent  $K_i$  values of discrete compounds were measured by assaying enzyme activity in the presence of different inhibitor compound concentrations and subtracting the background radioactivity measured in the absence of enzyme. The kinetic parameters ( $k_{\text{cat}}$ ,  $K_m$  for ATP) were measured for each enzyme under the usual assay conditions by determining the dependence of initial rates on ATP concentration. Inhibition data were fit to an equation for competitive inhibition using Kaleidagraph (Synergy Software), or were fit to an equation for competitive tight-binding inhibition using the software KineTic (BioKin, Ltd.).

**Inhibition of CDK4/Cyclin D Retinoblastoma Kinase Activity:**

A complex of human CDK4 and genetically truncated (1-264) cyclin D3 was purified using traditional biochemical chromatographic techniques from insect cells that had been co-infected with the corresponding baculovirus expression vectors (see e.g., Meijer and Kim, "*Chemical Inhibitors of Cyclin-Dependent Kinases*," Methods in Enzymol., 283 (1997), pp. 113-128.). The enzyme complex (5 nM) was assayed with 0.3-0.5  $\mu$ g of purified recombinant retinoblastoma protein fragment (Rb) as a substrate. The engineered Rb fragment (residues 386-928 of the native retinoblastoma protein; 62.3 kDa) contains the majority of the phosphorylation sites found in the

native 106-kDa protein, as well as a tag of six histidine residues for ease of purification.

Phosphorylated Rb substrate was captured by microfiltration on a nitrocellulose membrane and quantified using a phosphorimager as described above. For measurement of tight-binding inhibitors, the assay duration was extended to 60 minutes, during which the time-dependence of product formation was linear and initial rate conditions were met.  $K_i$  values were measured as described above and shown in Table 1. Percent inhibition was calculated as described above and shown in Table 2.

#### Inhibition of CDK2/Cyclin A Retinoblastoma Kinase Activity:

CDK2 was purified using published methodology (Rosenblatt et al., "Purification and Crystallization of Human Cyclin-dependent Kinase 2," J. Mol. Biol., 230, 1993, pp. 1317-1319) from insect cells that had been infected with a baculovirus expression vector. Cyclin A was purified from E. coli cells expressing full-length recombinant cyclin A, and a truncated cyclin A construct was generated by limited proteolysis and purified as described previously (Jeffrey et al., "Mechanism of CDK Activation Revealed by the Structure of a Cyclin A-CDK2 Complex," Nature, 376 (1995), pp. 313-320). A complex of CDK2 and proteolyzed cyclin A was prepared and purified by gel filtration. The substrate for this assay was the same Rb substrate fragment used for the CDK4 assays, and the methodology of the CDK2/ delta cyclin A and the CDK4/ delta cyclin D3 assays was essentially the same, except that CDK2 was present at 10 nM or 19 nM. The duration of the assay was 60 or 75 minutes, during which the time-dependence of product formation was linear and initial rate conditions were met.  $K_i$  values were measured as described above and shown in Table 1. And, the percent inhibition was calculated as described above and shown in Table 2.

#### Inhibition of CDK1(cdc2)/Cyclin B Histone H1 Kinase Activity:

The complex of human CDK1 (cdc2) and cyclin B was purchased from New England Biolabs (Beverly, MA). Alternatively, a CDK1/glutathione-S-transferase-cyclin B1 complex was purified using glutathione affinity chromatography from insect cells that had been co-infected with the corresponding baculovirus expression vectors. The assay was executed as described above at 30°C using 2.5 units of cdc2/cyclin B, 10 µg Histone H1 protein, and 0.1-0.3 µCi [ $^{32/33}\text{P}$ ]ATP per assay. Phosphorylated histone substrate was captured by microfiltration on a phosphocellulose P81 membrane and quantified using a phosphorimager as described above.  $K_i$  values were measured using the described curve-fitting programs. The results are shown in Table 3.

Inhibition of phosphorylated FGF receptor and LCK Tyrosine Kinase Activity:

Cloning, expression and purification of the cytosolic domain of FGFR1 tyrosine kinase (amino acids 456-766) containing three amino acid substitutions (L457V, C488A, and C584S) were conducted as previously described (Mohammadi, M., Schlessinger, J., & Hubbard, S. R. Cell, 86, (1996) 577-587). This domain was expressed in Sf9 insect cells using a baculovirus expression vector, and protein was purified using conventional techniques. The LCK tyrosine kinase was expressed in insect cells as an N-terminal deletion starting from amino acid 223 to the end of the protein at amino acid 509. The N-terminus of the protein also had two amino acid substitutions, P223M and C 224D. Kinases were purified using conventional chromatographic methods.

Tyrosine kinase activity was measured using a coupled, continuous spectrophotometric assay, in which production of phosphorylated poly(Glu,Tyr; 4:1) substrate and ADP is coupled to the pyruvate kinase-catalyzed transfer of a phosphate from phosphoenolpyruvate to ADP, with generation of pyruvate and regeneration of ATP. Pyruvate production is in turn coupled to the lactate dehydrogenase-catalyzed reduction of pyruvate to form lactate, with concomitant conversion of NADH to NAD<sup>+</sup>. Loss of NADH is monitored by measuring absorbance at 340 nm (see e.g., Technikova-Dobrova et al., "Spectrophotometric Determination of Functional Characteristics of Protein Kinases with Coupled Enzymatic Assay," FEBS Letters, 292 (1991), pp. 69-72). Enzyme activity was measured in the presence of 200 mM HEPES (pH 7.5), 2 mM phosphoenolpyruvate, 0.3 mM NADH, 20 mM MgCl<sub>2</sub>, 100  $\mu$ M ATP, 5 mM DTT, 5.1 or 25 mM poly (Glu,Tyr) 4:1 for P-FGF or P-LCK assays, respectively, and 15 units/mL each of pyruvate kinase and lactate dehydrogenase. Phosphorylated FGF receptor kinase was present at 100 nM and phosphorylated LCK kinase was present at 50 nM. Assays were performed under initial rate conditions at 37°C, and rates were corrected for any background rate measured in the absence of enzyme. Percent inhibition was calculated relative to control enzyme assayed in the presence of 2% (v/v) DMSO. The results are shown in Table 1.

Inhibition of Cell Growth: Assessment of Cytotoxicity:










Inhibition of cell growth was measured using the tetrazolium salt assay, which is based on the ability of viable cells to reduce 3-(4,5-dimethylthiazol-2-yl)-2,5-[2H]-diphenyltetrazolium bromide (MTT) to formazan (Mossmann, *Journal of Immunological Methods*, 65, (1983), pp. 55-58). The water-insoluble purple formazan product was then detected spectrophotometrically. The HCT-116 cell line was used as a representative cancer cell line and grown in 96-well plates. Cells were plated in McCoy's 5A Medium at a volume of 135  $\mu$ L/well. Plates were incubated for four hours before addition of inhibitor compounds. Different concentrations of inhibitor compounds were added in 0.5% (v/v) dimethylsulfoxide (15  $\mu$ L/well), and cells were incubated at 37°C (5%

CO<sub>2</sub>) for three to five days. At the end of the incubation, MTT was added to a final concentration of 0.2 mg/mL, and cells were incubated for 4 hours more at 37°C. After centrifugation of the plates and removal of medium, the absorbance of the formazan (solubilized in dimethylsulfoxide) was measured at 540 nm. The concentration of inhibitor compound causing 50%(IC<sub>50</sub>) or 90%(IC<sub>90</sub>) inhibition of growth was determined from the linear portion of a semi-log plot of inhibitor concentration versus percent inhibition. All results were compared to control cells treated only with 0.5% (v/v) dimethylsulfoxide. The IC<sub>50</sub> and IC<sub>90</sub> of the A – F compounds are shown in Table 1. Percent inhibitions at 0.25μM or 0.1μM of G compounds were calculated and shown in Table 2.












For the compounds shown in Tables 1 and 2, the group of -N(H)- and methyl (-CH<sub>3</sub>) of the formulae are sometimes shown as “-N-” and “—” for the simplicity, respectively, and the compounds in the form of salts are shown in their free base forms. For both Tables 1 and 2, “NT” indicates not tested. In Table 2, the “ ” refers to the point of attachment of the Formula (I) attached to the group R.

5












**Table 1**

Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at $1\mu\text{M}$	at $1\mu\text{M}$
A1		0.14	0.027	0.75	1.5	NT	NT
A2		0.0032	0.008	0.14	0.61	8	4
A3		0.17	0.021	0.75	2.1	7	5
A4		0.23	0.016	0.19	0.23	NT	NT
A5		0.072	0.066	1.2	3.1	5	3
A6		0.0075	0.015	0.22	0.6	NT	NT
A7		0.061	0.026	0.41	1.1	NT	NT
A8		0.054	0.005	0.74	1.7	NT	NT
A9		0.027	0.012	0.14	0.4	5	5














Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at $1\mu\text{M}$	at $1\mu\text{M}$
A10		0.29	0.049	1.6	3.5	NT	NT
A11		0.082	0.014	0.17	0.31	9	6
A12		0.091	0.02	0.13	0.28	NT	NT
A13		0.079	0.009	0.21	0.43	NT	NT
A14		0.06	0.014	0.042	0.11	0	0
A15		0.15	0.007	0.09	0.2	0	0
A16		0.068	0.025	0.17	0.3	NT	NT
A17		0.051	0.026	0.1	0.21	0	0
A18		0.19	0.006	0.92	2.1	NT	NT
A19		0.064	0.105	0.11	0.21	NT	NT
A20		0.162	0.032	0.076	0.18	0	0











Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at $1\mu\text{M}$	at $1\mu\text{M}$
A21		0.105	0.105	0.1	0.21	0	0
A22		0.006	0.005	0.17	0.42	NT	NT
A23		0.13	0.008	0.1	0.21	NT	NT
A24		0.1	0.094	0.16	0.24	NT	NT
A25		0.062	0.013	0.094	0.21	0	0
A26		0.079	0.02	0.17	0.38	NT	NT
A27		0.066	0.041	NT	NT	NT	NT
A28		0.025	0.076	0.06	0.17	0	0
A29		0.052	0.026	0.22	0.53	NT	NT
A30		0.069	0.007	0.09	0.23	NT	NT
A31		0.049	0.007	0.03	0.07	0	0

Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at $1\mu\text{M}$	at $1\mu\text{M}$
A32		0.077	0.006	0.28	0.68	NT	NT
A33		0.091	0.01	2.2	4.8	NT	NT
A34		0.099	0.01	0.14	0.29	NT	NT
A35		0.054	0.006	0.12	0.26	NT	NT
A36		0.223	0.044	0.78	1.3	NT	NT
A37		0.111	0.053	0.3	0.73	NT	NT
A38		0.14	0.024	0.1	0.23	NT	NT
A39		0.16	0.011	0.17	0.36	NT	NT
A40		0.094	0.009	0.04	0.085	NT	NT
A41		0.23	0.035	0.18	0.31	NT	NT
A42		0.165	0.023	0.15	0.37	NT	NT

Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		μM	μM	μM	μM	at 1 μM	at 1 μM
A43		0.109	0.013	0.58	1.9	NT	NT
A44		0.12	0.01	0.13	0.31	NT	NT
A45		0.048	0.007	0.086	0.21	NT	NT
A46		0.22	0.015	0.086	0.21	NT	NT
A47		0.055	0.005	0.58	1.4	NT	NT
A48		0.055	0.005	0.58	1.4	NT	NT
A49		0.26	0.011	0.32	0.63	NT	NT
A50		0.02	0.015	0.17	0.3	NT	NT
A51		0.11	0.007	0.09	0.22	NT	NT
A52		0.17	0.009	0.13	0.27	NT	NT

Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at $1\mu\text{M}$	at $1\mu\text{M}$
A53		0.12	0.004	0.15	0.3	NT	NT
A54		0.052	0.009	0.6	1.9	NT	NT
A55		0.056	0.017	3.2	>5	NT	NT
A56		0.067	0.008	4.5	>5	NT	NT
A57		0.054	0.01	3.8	>5	NT	NT
A58		3.4	1	NT	NT	NT	NT
A59		0.83	0.097	2.9	>5	NT	NT
A60		0.091	0.031	0.2	0.5	NT	NT
A61		0.0375	0.232	0.22	0.52	NT	NT
A62		0.0028	0.120	0.044	0.11	NT	NT

Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at $1\mu\text{M}$	at $1\mu\text{M}$
A63		0.0153	0.018	1.6	3.8	NT	NT
A64		0.019	0.022	>5	>5	NT	NT
A65		0.16	0.057	>0.5	>0.5	NT	NT
A66		0.018	0.35	0.12	0.23	NT	NT
A67		0.013	0.088	0.077	0.17	NT	NT
A68		0.0085	0.045	0.075	0.19	NT	NT
A69		0.0146	0.234	2.1	4.8	NT	NT
A70		0.0061	0.128	0.023	0.051	NT	NT
A71		0.0056	0.082	0.034	0.075	NT	NT
A72		0% @ $1\mu\text{M}$	0% @ $1\mu\text{M}$	>0.5	>0.5	NT	NT
A73		1.2	36	>5	>5	NT	NT

Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at 1 $\mu\text{M}$	at 1 $\mu\text{M}$
A74		0.0098	0.38	1.9	4.1	NT	NT
A75		0.0073	0.051	1.9	5	NT	NT
A76		0.015	0.12	1.1	2.3	NT	NT
A77		0.084	0.0052	0.12	0.22	NT	NT
A78		0.11	0.0082	0.13	0.28	NT	NT
B1		0.065	0.017	18	25	NT	NT
B2		0.096	0.018	3.6	8.3	NT	NT
B3		0.066	0.008	18	25	NT	NT
B4		0.22	0.008	0.52	1.3	NT	NT
B5		0.15	0.059	NT	NT	NT	NT





































Example	STRUCTURE	CDK2	CDK4	HCT-116	HCT-116	P-FGF	P-LCK
		Ki	Ki	IC50	IC90	% Inhibition	% Inhibition
		$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	$\mu\text{M}$	at $1\mu\text{M}$	at $1\mu\text{M}$
C		0.17	0.027	0.21	0.52	NT	NT
D1		0.0094	0.017	1	3.7	NT	NT
D2		0.051	0.025	0.58	1.7	NT	NT
E		0.4	0.24	#N/A	#N/A	NT	NT
F		0.27	0.14	1.8	3.8	NT	NT



























Table 2













 (I)					
Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G1		-3	-10	56	NT
G2		-9	8	51	NT
G3		-1	4	50	NT
G4		-7	4	53	NT
G5		-23	-3	47	NT
G6		-1	10	47	NT
G7		4	5	80	NT













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G8		-1	-5	50	NT
G9		-1	2	46	NT
G10		-4	-2	51	NT
G11		-12	6	49	NT
G12		-14	10	48	NT
G13		-16	0	44	NT
G14		9	26	50	NT
G15		-12	4	46	NT
G16		-7	1	50	NT
G17		2	7	45	NT
G18		-10	4	41	NT
G19		-7	4	43	NT













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G20		10	9	50	NT
G21		3	14	46	NT
G22		2	-3	50	NT
G23		8	7	63	NT
G24		7	11	58	NT
G25		-16	10	52	NT
G26		3	13	53	NT
G27		-6	4	60	NT
G28		1	19	66	NT
G29		4	32	55	NT
G30		5	5	82	NT

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G31		4	4	75	NT
G32		8	27	55	NT
G33		-3	10	54	NT
G34		8	9	53	NT
G35		5	8	53	NT
G36		-4	6	62	NT
G37		11	25	65	NT
G38		3	7	65	NT
G39		-4	-1	65	NT
G40		4	18	61	NT
G41		11	10	50	NT
G42		10	7	54	NT

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G43		5	12	62	NT
G44		8	14	72	NT
G45		16	16	89	NT
G46		4	11	58	NT
G47		-15	6	53	NT
G48		1	3	50	NT
G49		7	4	57	NT
G50		9	10	61	NT
G51		9	-1	89	NT
G52		10	5	65	NT
G53		17	6	89	NT
G54		8	9	86	NT













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G55		22	16	57	NT
G56		24	10	59	NT
G57		17	19	51	NT
G58		19	17	53	NT
G59		25	3	53	NT
G60		12	18	66	NT
G61		19	19	58	NT
G62		7	22	50	NT
G63		4	7	54	NT
G64		5	17	57	NT
G65		12	17	58	NT
G66		26	14	60	NT













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G67		25	12	82	NT
G68		6	3	59	NT
G69		7	3	70	NT
G70		4	10	53	NT
G71		3	16	59	NT
G72		2	12	54	NT
G73		19	18	56	NT
G74		23	18	78	NT
G75		21	15	64	NT
G76		7	7	61	NT
G77		9	6	51	NT
G78		11	13	58	NT













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G79		2	10	51	NT
G80		3	7	36	NT
G81		15	19	58	NT
G82		24	15	48	NT
G83		16	7	64	NT
G84		9	3	63	NT
G85		7	6	56	NT
G86		10	0	73	39
G87		-40	13	78	54
G88		-23	6	49	24
G89		-20	-4	59	30
G90		-30	2	54	24



























Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G91		-19	9	68	34
G92		-7	10	98	85
G93		-4	11	58	26
G94		10	4	80	38
G95		12	18	63	32
G96		-10	5	56	20
G97		2	11	56	28
G98		-24	11	60	22
G99		-16	23	62	24
G100		13	22	89	27
G101		11	19	46	10
G102		-3	18	55	18













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G103		-2	5	59	21
G104		8	14	91	38
G105		3	9	67	24
G106		7	7	61	0
G107		-5	12	59	5
G108		-5	12	60	0
G109		6	22	54	0
G110		15	17	52	0
G111		15	20	52	0
G112		-11	13	57	6
G113		51	43	80	26
G114		-3	11	48	0













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G115		27	24	88	30
G116		7	24	46	0
G117		4	13	55	6
G118		11	21	48	5
G119		11	31	64	0
G120		14	23	54	0
G121		13	20	55	0
G122		15	15	59	0
G123		-5	12	50	4
G124		10	17	64	0
G125		3	9	59	13
G126		3	7	53	0

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G127		10	29	61	0
G128		15	41	78	23
G129		11	19	51	0
G130		1	11	77	0
G131		6	16	58	0
G132		9	27	52	0
G133		-6	13	47	0
G134		60	38	89	1
G135		2	25	56	0
G136		-4	12	49	0
G137		2	25	59	0
G138		7	17	65	0













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G139		6	17	56	0
G140		15	19	54	0
G141		0	11	50	0
G142		11	20	46	0
G143		2	27	65	0
G144		0	11	57	0
G145		19	18	68	0
G146		6	7	52	1
G147		10	35	65	0
G148		5	14	48	0
G149		23	36	82	10
G150		15	25	57	0













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G151		16	7	94	27
G152		16	7	53	5
G153		18	15	66	0
G154		10	34	91	17
G155		32	38	67	0
G156		24	24	61	0
G157		18	30	87	0
G158		22	40	73	0
G159		6	5	74	0
G160		24	6	87	0
G161		13	7	57	0
G162		25	20	84	3













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G163		16	21	67	0
G164		13	19	59	0
G165		1	22	61	0
G166		22	4	90	8
G167		19	29	53	26
G168		4	17	52	25
G169		38	49	71	32
G170		26	32	78	44
G171		4	16	47	30
G172		36	54	61	37
G173		14	21	54	18
G174		0	15	56	28













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G175		15	31	47	11
G176		22	21	62	32
G177		10	23	45	24
G178		20	54	83	43
G179		-5	10	42	21
G180		22	29	68	25
G181		30	36	75	28
G182		-2	6	52	23
G183		23	38	48	11
G184		34	36	58	17
G185		7	20	49	10
G186		7	30	51	14















Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G187		-2	12	48	16
G188		29	39	51	17
G189		10	18	56	13
G190		5	4	55	9
G191		14	15	51	3
G192		8	23	49	13
G193		19	26	56	17
G194		1	19	49	20
G195		20	26	47	17
G196		7	13	48	13
G197		14	9	61	26
G198		19	42	44	9













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G199		18	22	52	12
G200		15	23	50	19
G201		6	26	43	17
G202		5	25	48	20
G203		7	20	53	18
G204		6	10	49	19
G205		1	7	52	12
G206		13	19	58	13
G207		15	15	66	15
G208		36	36	56	11
G209		9	16	50	9
G210		13	20	45	15













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G211		23	23	45	13
G212		16	20	53	17
G213		-1	9	49	16
G214		11	16	52	15
G215		18	27	73	22
G216		29	36	54	16
G217		17	34	51	8
G218		8	19	52	11
G219		10	13	52	8
G220		10	12	54	18
G221		29	34	74	20
G222		30	27	61	8

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G223		14	19	50	14
G224		16	28	53	10
G225		36	41	51	0
G226		48	44	56	15
G227		28	17	84	17
G228		38	36	75	24
G229		12	9	44	8
G230		8	20	52	19
G231		15	30	55	6
G232		9	19	53	9
G233		10	21	48	16
G234		35	39	63	20













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G235		14	14	51	10
G236		39	36	66	16
G237		9	12	49	20
G238		18	26	58	22
G239		15	27	50	21
G240		11	27	52	15
G241		22	29	57	28
G242		43	49	69	34
G243		18	11	40	31
G244		35	27	67	30
G245		36	34	55	18
G246		32	35	62	20













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G247		17	23	54	12
G248		8	17	55	13
G249		39	40	80	37
G250		39	39	72	31
G251		10	11	48	16
G252		29	31	67	33
G253		16	26	62	19
G254		28	20	75	35
G255		-25	0	63	25
G256		-12	11	49	19
G257		-43	2	51	33
G258		25	20	84	41













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G259		-30	-20	46	21
G260		-27	1	59	30
G261		30	27	69	26
G262		1	28	59	24
G263		2	24	54	26
G264		-9	13	53	22
G265		-24	1	48	21
G266		0	7	30	28
G267		-23	-14	44	24
G268		-11	2	63	26
G269		6	2	65	19
G270		26	17	45	15

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G271		32	27	49	8
G272		-20	0	47	10
G273		-33	-2	45	17
G274		-4	5	47	15
G275		27	19	63	25
G276		2	-2	58	25
G277		8	26	56	11
G278		25	27	76	6
G279		-2	17	52	2
G280		-7	9	70	7
G281		-9	12	63	6
G282		-14	6	56	1



























Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G283		-6	10	75	16
G284		-30	-12	69	4
G285		13	18	60	1
G286		23	25	60	10
G287		-8	7	32	9
G288		-10	11	60	16
G289		-6	9	62	7
G290		-12	-2	39	2
G291		6	14	53	5
G292		-31	-15	60	0
G293		23	32	56	10
G294		30	29	74	15













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G295		8	22	78	10
G296		0	22	43	0
G297		-3	8	45	0
G298		-9	30	46	0
G299		-11	-7	49	10
G300		-18	-10	70	18
G301		13	25	79	18
G302		17	29	63	17
G303		2	45	72	18
G304		0	20	54	15
G305		-6	11	81	29
G306		-17	0	44	9













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G307		-3	12	54	8
G308		-23	5	60	5
G309		17	30	65	5
G310		-5	10	53	14
G311		-1	46	43	5
G312		-10	5	58	8
G313		11	12	57	10
G314		-7	0	47	9
G315		-1	11	50	7
G316		-2	11	56	1
G317		27	33	87	27
G318		-14	1	48	16

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G319		-5	14	54	17
G320		-15	5	46	12
G321		1	19	51	11
G322		-12	4	50	13
G323		-14	7	72	14
G324		28	26	60	5
G325		13	23	51	6
G326		12	22	13	11
G327		-7	11	55	8
G328		19	29	89	33
G329		10	19	56	9
G330		-1	12	55	2













Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G331		1	37	93	54
G332		17	14	66	21
G333		-16	-5	51	4
G334		-2	8	41	6
G335		2	21	42	3
G336		36	36	51	12
G337		7	11	52	13
G338		31	27	54	18
G339		6	34	48	9
G340		-12	9	40	28
G341		-11	10	93	23
G342		-10	13	55	20

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.03 $\mu$ M	% Inhibition at 0.25 $\mu$ M	% Inhibition at 0.1 $\mu$ M
G343		1	12	71	27
G344		25	18	50	24
G345		-2	6	36	22
G346		11	11	52	15
G347		13	15	53	21
G348		-21	14	51	23
G349		-6	11	48	31
G350		4	9	41	24
G351		0	-1	62	24
G352		4	0	64	24
G353		35	46	55	0
G354		9	3	60	31

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G355		29	13	54	28
G356		-16	5	50	27
G357		-14	17	88	17
G358		-5	7	55	21
G359		-7	-5	69	25
G360		31	9	73	18
G361		-3	-1	48	26
G362		12	6	57	9
G363		27	16	63	24
G364		-12	9	40	20
G365		8	22	93	34
G366		1	8	55	26

Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G367		38	15	71	22
G368		3	-3	50	25
G369		26	8	36	21
G370		22	17	52	19
G371		11	7	53	27
G372		-21	6	51	19
G373		-7	10	48	31
G374		-10	13	41	21
G375		10	-3	62	26
G376		20	-6	64	27
G377		10	-1	55	25
G378		40	8	60	27



Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G379		13	8	54	31
G380		-19	7	50	26
G381		5	38	88	43
G382		-17	8	47	17
G383		15	-10	51	33
G384		45	23	84	38
G385		1	-1	50	28
G386		19	12	51	28
G387		10	11	52	32
G388		3	8	45	9
G389		-12	4	61	11
G390		-14	2	40	10















Ex.	R	CDK2	CDK4	HCT-116	HCT-116
		% Inhibition	% Inhibition	% Inhibition	% Inhibition
		at 0.03 $\mu$ M	at 0.03 $\mu$ M	at 0.25 $\mu$ M	at 0.1 $\mu$ M
G391		1	-7	50	11
G392		12	-2	49	8
G393		8	0	46	8
G394		35	11	73	14
G395		17	15	48	19
G396		-6	9	67	14

Table 3


Example	Structure	K <sub>i</sub> CDK1/B (μM)
A14		0.042
A15		0.068
A20		0.15
A21		0.12
A28		0.062
A31		0.082
A37		0.044
A44		0.072

The examples above illustrate compounds according to Formula (I) and assays that may readily be performed to determine their activity levels against the various enzymes and for cell growth inhibition. It will be apparent that such assays or other suitable assays known in the art may be used to select an inhibitor having a desired level of activity against a selected target.

While the invention has been illustrated by reference to specific and preferred embodiments, the invention is intended not to be limited by the foregoing description, but to be defined by the appended claims and their equivalents.

What is claimed is:

1. A compound represented by Formula (I):



wherein:

$R^1$  and  $R^2$  are each independently hydrogen, or an alkyl, alkenyl, alkynyl, heteroalkyl, alkoxy, amino alkyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl group unsubstituted or substituted with one or more substituents selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N}-\text{OH}$ ,  $\text{N}-\text{OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{OR}_b$ ,  $-\text{CO}-\text{R}_c$ ,  $\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{OR}_c$ ,  $=\text{O}$ ,  $=\text{S}$ ,  $\text{SO}_2-\text{R}_c$ ,  $-\text{SO}-\text{R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{R}_e$ ,  $\text{NR}_c-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{SO}_2-\text{R}_c$ ,  $-\text{O}-\text{SO}-\text{R}_c$ ,  $-\text{O}-\text{S}-\text{R}_c$ ,  $-\text{S}-\text{CO}-\text{R}_c$ ,  $-\text{SO}-\text{CO}-\text{OR}_c$ ,  $-\text{SO}_2-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{SO}_3$ ,  $-\text{NR}_c-\text{SR}_d$ ,  $-\text{NR}_c-\text{SO}-\text{R}_d$ ,  $\text{NR}_c-\text{SO}_2-\text{R}_d$ ,  $-\text{CO}-\text{SR}_c$ ,  $-\text{CO}-\text{SO}-\text{R}_c$ ,  $\text{CO}-\text{SO}_2-\text{R}_c$ ,  $-\text{CS}-\text{R}_c$ ,  $-\text{CSO}-\text{R}_c$ ,  $-\text{CSO}_2-\text{R}_c$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{O}-\text{CS}-\text{R}_c$ ,  $-\text{O}-\text{CSO}-\text{R}_c$ ,  $\text{O}-\text{CSO}_2-\text{R}_c$ ,  $-\text{SO}_2-\text{NR}_d\text{R}_e$ ,  $-\text{SO}-\text{NR}_d\text{R}_e$ ,  $-\text{S}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_d-\text{CSO}_2-\text{R}_d$ ,  $-\text{NR}_c-\text{CSO}-\text{R}_d$ ,  $\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{SH}$ ,  $-\text{S}-\text{R}_b$ , and  $-\text{PO}_2-\text{OR}_c$ , where  $\text{R}_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $\text{R}_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $\text{R}_c$ ,  $\text{R}_d$  and  $\text{R}_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_f$ ,  $-\text{O}-\text{CO}-\text{R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $\text{R}_d$  and  $\text{R}_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $\text{R}_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ ,  $=\text{O}$ ,  $-\text{N}-\text{OH}$ ,  $\text{N}-\text{OR}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{C}-\text{CO}-\text{OR}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{SH}$ ,  $-\text{O}-\text{R}_b$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{S}-\text{R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl,

unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are defined above, where  $R^1$  and  $R^2$  are not both hydrogen; or

$R^1$  or  $R^2$ , together with the  $N-\overset{\text{O}}{\underset{\parallel}{C}}$  and two adjacent carbon atoms of the phenyl ring of Formula (I), forms a 5- or 6-membered ring structure fused to the phenyl ring of Formula (I) and unsubstituted or substituted with one or more substituents independently selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N}-\text{OH}$ ,  $-\text{N}-\text{OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{OR}_b$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{OR}_c$ ,  $=\text{O}$ ,  $=\text{S}$ ,  $-\text{SO}_2-\text{R}_c$ ,  $-\text{SO}-\text{R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{SO}_2-\text{R}_c$ ,  $-\text{O}-\text{SO}-\text{R}_c$ ,  $-\text{O}-\text{S}-\text{R}_c$ ,  $-\text{S}-\text{CO}-\text{R}_c$ ,  $-\text{SO}-\text{CO}-\text{OR}_c$ ,  $-\text{SO}_2-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{SO}_3$ ,  $-\text{NR}_c-\text{SR}_d$ ,  $-\text{NR}_c-\text{SO}-\text{R}_d$ ,  $-\text{NR}_c-\text{SO}_2-\text{R}_d$ ,  $-\text{CO}-\text{SR}_c$ ,  $-\text{CO}-\text{SO}-\text{R}_c$ ,  $-\text{CO}-\text{SO}_2-\text{R}_c$ ,  $-\text{CS}-\text{R}_c$ ,  $-\text{CSO}-\text{R}_c$ ,  $-\text{CSO}_2-\text{R}_c$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{O}-\text{CS}-\text{R}_c$ ,  $-\text{O}-\text{CSO}-\text{R}_c$ ,  $-\text{O}-\text{CSO}_2-\text{R}_c$ ,  $-\text{SO}_2-\text{NR}_d\text{R}_e$ ,  $-\text{SO}-\text{NR}_d\text{R}_e$ ,  $-\text{S}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_d-\text{CSO}_2-\text{R}_d$ ,  $-\text{NR}_c-\text{CSO}-\text{R}_d$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{SH}$ ,  $-\text{S}-\text{R}_b$ , and  $-\text{PO}_2-\text{OR}_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_f$ ,  $-\text{O}-\text{CO}-\text{R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ ,  $=\text{O}$ ,  $-\text{N}-\text{OH}$ ,  $-\text{N}-\text{OR}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{C}-\text{CO}-\text{OR}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{SH}$ ,  $-\text{O}-\text{R}_b$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{S}-\text{R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are as defined above; or

$R^1$  and  $R^2$ , taken together with the nitrogen atom to which they are bonded, form a monocyclic or fused or non-fused polycyclic structure which may contain one to three additional heteroatoms, the structure being unsubstituted or substituted with one or more substituents selected from the group consisting alkyl, heteroalkyl, haloalkyl, haloaryl,

halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, -O-R<sub>a</sub>-O-, -OR<sub>b</sub>, -CO-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-OR<sub>c</sub>, -O-CO-O-CO-R<sub>c</sub>, -O-OR<sub>c</sub>, =O, =S, -SO<sub>2</sub>-R<sub>c</sub>, -SO-R<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-R<sub>e</sub>, -NR<sub>c</sub>-CO-OR<sub>e</sub>, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>, -O-SO<sub>2</sub>-R<sub>c</sub>, -O-SO-R<sub>c</sub>, -O-S-R<sub>c</sub>, -S-CO-R<sub>c</sub>, -SO-CO-OR<sub>c</sub>, -SO<sub>2</sub>-CO-OR<sub>c</sub>, -O-SO<sub>3</sub>, -NR<sub>c</sub>-SR<sub>d</sub>, -NR<sub>c</sub>-SO-R<sub>d</sub>, NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>, -CO-SR<sub>c</sub>, -CO-SO-R<sub>c</sub>, -CO-SO<sub>2</sub>-R<sub>c</sub>, -CS-R<sub>c</sub>, -CSO-R<sub>c</sub>, -CSO<sub>2</sub>-R<sub>c</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -O-CS-R<sub>c</sub>, -O-CSO-R<sub>c</sub>, -O-CSO<sub>2</sub>-R<sub>c</sub>, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>c</sub>-CSO-R<sub>d</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, -COR<sub>f</sub>, -COOR<sub>f</sub>, -O-CO-O-R<sub>f</sub>, -O-CO-R<sub>f</sub>, -OH, Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where R<sub>d</sub> and R<sub>e</sub> can cyclize to form a heteroaryl or heterocycloalkyl group, and R<sub>f</sub> is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, =O, -N-OH, N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are defined above;

R<sup>3</sup> is an aryl, heteroaryl, alkyl, or cycloalkyl group, unsubstituted or substituted with one or more substituents independently selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, -O-R<sub>a</sub>-O-, -OR<sub>b</sub>, -CO-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-OR<sub>c</sub>, -O-CO-O-CO-R<sub>c</sub>, -O-OR<sub>c</sub>, =O, =S, -SO<sub>2</sub>-R<sub>c</sub>, -SO-R<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-R<sub>e</sub>, -NR<sub>c</sub>-CO-OR<sub>e</sub>, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>, -O-SO<sub>2</sub>-R<sub>c</sub>, -O-SO-R<sub>c</sub>, -O-S-R<sub>c</sub>, -S-CO-R<sub>c</sub>, -SO-CO-OR<sub>c</sub>, -SO<sub>2</sub>-CO-OR<sub>c</sub>, -O-SO<sub>3</sub>, -NR<sub>c</sub>-SR<sub>d</sub>, -NR<sub>c</sub>-SO-R<sub>d</sub>, NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>, -CO-SR<sub>c</sub>, -CO-SO-R<sub>c</sub>, -CO-SO<sub>2</sub>-R<sub>c</sub>, -CS-R<sub>c</sub>, -CSO-R<sub>c</sub>, -CSO<sub>2</sub>-R<sub>c</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -O-CS-R<sub>c</sub>, -O-CSO-R<sub>c</sub>, -O-CSO<sub>2</sub>-R<sub>c</sub>, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>c</sub>-CSO-R<sub>d</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl,

alkenyl, alkynyl, halogen,  $-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl,  $\text{R}_c$ ,  $\text{R}_d$  and  $\text{R}_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_f$ ,  $-\text{O}-\text{CO}-\text{R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $\text{R}_d$  and  $\text{R}_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $\text{R}_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ ,  $=\text{O}$ ,  $-\text{N}-\text{OH}$ ,  $\text{N}-\text{OR}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{C}-\text{CO}-\text{OR}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{SH}$ ,  $-\text{O}-\text{R}_b$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{S}-\text{R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $\text{R}_a$ ,  $\text{R}_b$ ,  $\text{R}_c$ ,  $\text{R}_d$ , and  $\text{R}_e$  are defined above; and

Y is hydrogen, alkyl, heteroalkyl, haloalkyl, halocycloalkyl, haloheterocycloalkyl, cycloalkyl, heterocycloalkyl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N}-\text{OH}$ ,  $\text{N}-\text{OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O}-\text{R}_a-\text{O}-$ ,  $-\text{OR}_b$ ,  $-\text{CO}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{OR}_c$ ,  $=\text{O}$ ,  $=\text{S}$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{O}-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{R}_e$ ,  $-\text{NR}_c-\text{CO}-\text{OR}_c$ ,  $-\text{CO}-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{O}-\text{SO}_2-\text{R}_c$ ,  $-\text{O}-\text{SO}-\text{R}_c$ ,  $-\text{O}-\text{S}-\text{R}_c$ ,  $-\text{S}-\text{CO}-\text{R}_c$ ,  $-\text{SO}-\text{CO}-\text{OR}_c$ ,  $-\text{SO}-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{SO}_3$ ,  $-\text{NR}_c-\text{SR}_d$ ,  $-\text{NR}_c-\text{SO}-\text{R}_d$ ,  $\text{NR}_c-\text{SO}_2-\text{R}_d$ ,  $-\text{CO}-\text{SR}_c$ ,  $-\text{CO}-\text{SO}-\text{R}_c$ ,  $-\text{CO}-\text{SO}_2-\text{R}_c$ ,  $-\text{CS}-\text{R}_c$ ,  $-\text{CSO}-\text{R}_c$ ,  $-\text{CSO}_2-\text{R}_c$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{O}-\text{CS}-\text{R}_c$ ,  $-\text{O}-\text{CSO}-\text{R}_c$ ,  $-\text{O}-\text{SO}_2-\text{R}_c$ ,  $-\text{SO}_2-\text{NR}_d\text{R}_e$ ,  $-\text{SO}-\text{NR}_d\text{R}_e$ ,  $-\text{S}-\text{NR}_d\text{R}_e$ ,  $-\text{NR}_d-\text{CSO}_2-\text{R}_d$ ,  $-\text{NR}_c-\text{CSO}-\text{R}_d$ ,  $-\text{NR}_c-\text{CS}-\text{R}_d$ ,  $-\text{SH}$ ,  $-\text{S}-\text{R}_b$ , and  $-\text{PO}_2-\text{OR}_c$ , where  $\text{R}_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $\text{R}_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO}-\text{R}_c$ ,  $-\text{CO}-\text{OR}_c$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_c$ ,  $-\text{O}-\text{CO}-\text{R}_c$ ,  $-\text{NR}_c-\text{CO}-\text{R}_d$ ,  $-\text{CO}-\text{NR}_d\text{R}_e$ ,  $-\text{OH}$ , heterocycloalkyl, and cycloalkyl,  $\text{R}_c$ ,  $\text{R}_d$  and  $\text{R}_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O}-\text{CO}-\text{O}-\text{R}_f$ ,  $-\text{O}-\text{CO}-\text{R}_f$ ,  $-\text{OH}$ , cycloalkyl, and heterocycloalkyl, where  $\text{R}_d$  and  $\text{R}_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $\text{R}_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl;

or a pharmaceutically acceptable salt of said compound, pharmaceutically acceptable prodrug of said compound, pharmaceutically active metabolite of said compound, or pharmaceutically acceptable salt of said metabolite.

2. A compound, salt, prodrug, or metabolite according to claim 1, wherein Y is hydrogen, OH, a halogen or an alkoxy group.

3. A compound, salt, prodrug, or metabolite according to claim 1, wherein  $R^3$  is an aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen, alkoxy, alkyl, nitro,  $-OH$ , amide, and  $-SO_2$ -alkyl.
4. A compound, salt, prodrug, or metabolite according to claim 1, wherein the  $-C(O)NR^1R^2$  moiety is meta or para to the amine linking the phenyl and thiazole rings.
5. A compound or salt according to claim 1, wherein the  $-C(O)NR^1R^2$  moiety is para to the amine linking the phenyl and thiazole rings.
6. A compound or salt according to claim 5, wherein Y is hydrogen, or an alkyl, alkoxy, or halogen group; and  
 $R^3$  is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen, alkyl, alkoxy,  $-OH$ , amide, and  $-SO_2$ -alkyl.
7. A compound or salt according to claim 5, wherein Y is hydrogen; and  
 $R^3$  is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen and alkyl.
8. A compound or salt according to claim 1, wherein the  $-C(O)NR^1R^2$  moiety is para and Y is meta to the amine linking the phenyl and thiazole rings.
9. A compound or salt according to claim 8, wherein Y is hydrogen, or an alkyl, alkoxy, or halogen group; and  
 $R^3$  is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen, alkyl, alkoxy,  $-OH$ , amide, and  $-SO_2$ -alkyl.
10. A compound or salt according to claim 8, wherein Y is hydrogen; and  
 $R^3$  is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen and alkyl.
11. A compound, salt, prodrug, or metabolite according to claim 1, wherein the  $-C(O)NR^1R^2$  moiety is para to the NH moiety;  
 $R^1$  and  $R^2$  are each independently hydrogen, or an alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl group unsubstituted or substituted with one or more substituents independently selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-NO_2$ ,  $-NH_2$ ,  $-N-OH$ ,  $N-OR_c$ ,  $-CN$ ,  $-(CH_2)_z-CN$  where z is 0-4, halogen,  $-OH$ ,  $-O-R_a-O-$ ,  $-OR_b$ ,  $-CO-R_c$ ,  $-O-CO-R_c$ ,  $-CO-OR_c$ ,  $-O-CO-OR_c$ ,  $-O-CO-O-CO-R_c$ ,  $-O-OR_c$ , keto ( $=O$ ), thioketo ( $=S$ ),  $-SO_2-R_c$ ,  $-SO-R_c$ ,  $-NR_dR_e$ ,  $-CO-NR_dR_e$ ,  $-O-CO-NR_dR_e$ ,  $-NR_c-CO-NR_dR_e$ ,  $-NR_c-CO-R_e$ ,  $-NR_c-CO-OR_e$ ,  $-CO-NR_c-CO-R_d$ ,  $-O-SO_2-R_c$ ,  $-O-SO-R_c$ ,  $-O-S-R_c$ ,  $-S-CO-R_c$ ,  $-SO-CO-OR_c$ ,  $-SO_2-CO-OR_c$ ,  $-O-SO_3$ ,  $-NR_c-SR_d$ ,  $-NR_c-SO-R_d$ ,  $-NR_c-SO_2-R_d$ ,  $-CO-SR_c$ ,  $-CO-SO-R_c$ ,  $-CO-SO_2-R_c$ ,  $-CS-R_c$ ,  $-CSO-R_c$ ,  $-CSO_2-R_c$ ,  $-NR_c-CS-R_d$ ,  $-O-CS-R_c$ ,  $-O-CSO-R_c$ ,  $-O-CSO_2-R_c$ ,  $-SO_2-$



NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>e</sub>-CSO-R<sub>d</sub>, -NR<sub>e</sub>-CS-R<sub>d</sub>, -SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, -COR<sub>f</sub>, -COOR<sub>f</sub>, -O-CO-O-R<sub>f</sub>, -O-CO-R<sub>f</sub>, -OH, Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where R<sub>d</sub> and R<sub>e</sub> can cyclize to form a heteroaryl or heterocycloalkyl group, and R<sub>f</sub> is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, =O, -N-OH, N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are defined above; where R<sup>1</sup> and R<sup>2</sup> are not both hydrogen.

12. A compound, salt, prodrug, or metabolite according to claim 11, wherein Y is hydrogen, or an alkyl, alkoxy, or halogen group; and

R<sup>3</sup> is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen, alkyl, alkoxy, -OH, amide, and -SO<sub>2</sub>-alkyl.

13. A compound, salt, prodrug, or metabolite according to claim 11, wherein Y is hydrogen; and

R<sup>3</sup> is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen and alkyl.

14. A compound, salt, prodrug, or metabolite according to claim 1, wherein the -C(O)NR<sup>1</sup>R<sup>2</sup> moiety is para to the NH moiety;

R<sup>1</sup> is hydrogen; and

R<sup>2</sup> is an alkyl, alkenyl, alkynyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl group unsubstituted or substituted with one or more substituents independently selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, -O-R<sub>a</sub>-O-, -OR<sub>b</sub>, -CO-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-OR<sub>c</sub>, -O-CO-O-CO-R<sub>c</sub>, -O-OR<sub>c</sub>, =O, =S, -SO<sub>2</sub>-R<sub>c</sub>, -SO-R<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-R<sub>e</sub>, -NR<sub>c</sub>-CO-OR<sub>c</sub>, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>, -O-

SO<sub>2</sub>-R<sub>c</sub>, -O-SO-R<sub>c</sub>, -O-S-R<sub>c</sub>, -S-CO-R<sub>c</sub>, -SO-CO-OR<sub>c</sub>, -SO<sub>2</sub>-CO-OR<sub>c</sub>, -O-SO<sub>3</sub>, -NR<sub>c</sub>-SR<sub>d</sub>, -NR<sub>c</sub>-SO-R<sub>d</sub>, -NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>, -CO-SR<sub>c</sub>, -CO-SO-R<sub>c</sub>, -CO-SO<sub>2</sub>-R<sub>c</sub>, -CS-R<sub>c</sub>, -CSO-R<sub>c</sub>, -CSO<sub>2</sub>-R<sub>c</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -O-CS-R<sub>c</sub>, -O-CSO-R<sub>c</sub>, -O-CSO<sub>2</sub>-R<sub>c</sub>, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>c</sub>-CSO-R<sub>d</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, -COR<sub>f</sub>, -COOR<sub>f</sub>, -O-CO-O-R<sub>f</sub>, -O-CO-R<sub>f</sub>, -OH, Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where R<sub>d</sub> and R<sub>e</sub> can cyclize to form a heteroaryl or heterocycloalkyl group, and R<sub>f</sub> is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, =O, -N-OH, -N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are as defined above.


15. A compound, salt, prodrug, or metabolite according to claim 14, wherein Y is selected from the group consisting of hydrogen or a hydroxy, halogen, alkyl or alkoxy group;

R<sup>3</sup> is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen, alkyl, alkoxy, -OH, amide, and -SO<sub>2</sub>-alkyl.

16. A compound, salt, prodrug, or metabolite according to claim 14, wherein Y is hydrogen; and

R<sup>3</sup> is a monocyclic aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halogen and alkyl.

17. A compound represented by Formula II:



wherein:

$R^1$  is selected from the group consisting of hydrogen, or an alkyl, alkenyl, alkynyl, heteroalkyl, alkoxy, amino-alkyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl group unsubstituted or substituted with one or more substituents selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-NO_2$ ,  $-NH_2$ ,  $-N-OH$ ,  $N-OR_c$ ,  $-CN$ ,  $-(CH_2)_z-CN$  where  $z$  is 0-4, halogen,  $-OH$ ,  $-O-R_a-O-$ ,  $-OR_b$ ,  $-CO-R_c$ ,  $-O-CO-R_c$ ,  $-CO-OR_c$ ,  $-O-CO-OR_c$ ,  $-O-CO-O-CO-R_c$ ,  $-O-OR_c$ ,  $=O$ ,  $=S$ ,  $-SO_2-R_c$ ,  $-SO-R_c$ ,  $-NR_dR_e$ ,  $-CO-NR_dR_e$ ,  $-O-CO-NR_dR_e$ ,  $-NR_c-CO-NR_dR_e$ ,  $-NR_c-CO-R_e$ ,  $-NR_c-CO-OR_e$ ,  $-CO-NR_c-CO-R_d$ ,  $-O-SO_2-R_c$ ,  $-O-SO-R_c$ ,  $-O-S-R_c$ ,  $-S-CO-R_c$ ,  $-SO_2-CO-OR_c$ ,  $-SO-CO-OR_c$ ,  $-O-SO_3$ ,  $-NR_c-SR_d$ ,  $-NR_c-SO-R_d$ ,  $NR_c-SO_2-R_d$ ,  $-CO-SR_c$ ,  $-CO-SO-R_c$ ,  $-CO-SO_2-R_c$ ,  $-CS-R_c$ ,  $-CSO-R_c$ ,  $-CSO_2-R_c$ ,  $-NR_c-CS-R_d$ ,  $-O-CS-R_c$ ,  $-O-CSO-R_c$ ,  $-O-CSO_2-R_c$ ,  $-SO_2-NR_dR_e$ ,  $-SO-NR_dR_e$ ,  $-S-NR_dR_e$ ,  $-NR_d-CSO_2-R_d$ ,  $-NR_c-CSO-R_d$ ,  $-NR_c-CS-R_d$ ,  $-SH$ ,  $-S-R_b$ , and  $-PO_2-OR_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-CO-R_c$ ,  $-CO-OR_c$ ,  $-O-CO-O-R_c$ ,  $-O-CO-R_c$ ,  $-NR_c-CO-R_d$ ,  $-CO-NR_dR_e$ ,  $-OH$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-COR_f$ ,  $-COOR_f$ ,  $-O-CO-O-R_f$ ,  $-O-CO-R_f$ ,  $-OH$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $NO_2$ ,  $-NH_2$ ,  $-CN$ ,  $-(CH_2)_z-CN$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-OH$ ,  $=O$ ,  $-N-OH$ ,  $N-OR_c$ ,  $-NR_dR_e$ ,  $-CO-NR_dR_e$ ,  $-CO-OR_c$ ,  $-CO-R_c$ ,  $-NR_c-CO-NR_dR_e$ ,  $-C-CO-OR_c$ ,  $-NR_c-CO-R_d$ ,  $-O-CO-O-R_c$ ,  $-O-CO-NR_dR_e$ ,  $-SH$ ,  $-O-R_b$ ,  $-O-R_a-O-$ ,  $-S-R_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  as are defined above, where  $R^1$  and  $R^2$  are not both hydrogen;

$R^4$ ,  $R^5$ , and  $R^6$  are each independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z\text{-CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O-R}_a\text{-O-}$ ,  $-\text{OR}_b$ ,  $-\text{CO-R}_c$ ,  $-\text{O-CO-R}_c$ ,  $-\text{CO-OR}_c$ ,  $-\text{O-CO-OR}_c$ ,  $-\text{O-CO-O-CO-R}_c$ ,  $\text{O-OR}_c$ ,  $=\text{O}$ ,  $=\text{S}$ ,  $-\text{SO}_2\text{-R}_c$ ,  $-\text{SO-R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{O-CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c\text{-CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c\text{-CO-R}_e$ ,  $-\text{NR}_c\text{-CO-OR}_e$ ,  $-\text{CO-NR}_c\text{-CO-R}_d$ ,  $-\text{O-SO}_2\text{-R}_c$ ,  $-\text{O-SO-R}_c$ ,  $-\text{O-S-R}_c$ ,  $-\text{S-CO-R}_c$ ,  $-\text{SO-CO-OR}_c$ ,  $-\text{SO}_2\text{-CO-OR}_c$ ,  $-\text{O-SO}_3$ ,  $-\text{NR}_c\text{-SR}_d$ ,  $-\text{NR}_c\text{-SO-R}_d$ ,  $\text{NR}_c\text{-SO}_2\text{-R}_d$ ,  $-\text{CO-SR}_c$ ,  $-\text{CO-SO-R}_c$ ,  $-\text{CO-SO}_2\text{-R}_c$ ,  $-\text{CS-R}_c$ ,  $-\text{CSO-R}_c$ ,  $-\text{CSO}_2\text{-R}_c$ ,  $-\text{NR}_c\text{-CS-R}_d$ ,  $-\text{O-CS-R}_c$ ,  $-\text{O-CSO-R}_c$ ,  $-\text{O-CSO}_2\text{-R}_c$ ,  $-\text{SO}_2\text{-NR}_d\text{R}_e$ ,  $-\text{SO-NR}_d\text{R}_e$ ,  $-\text{S-NR}_d\text{R}_e$ ,  $-\text{NR}_d\text{-CSO}_2\text{-R}_d$ ,  $-\text{NR}_c\text{-CSO-R}_d$ ,  $-\text{NR}_c\text{-CS-R}_d$ ,  $-\text{SH}$ ,  $-\text{S-R}_b$ , and  $-\text{PO}_2\text{-OR}_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO-R}_c$ ,  $-\text{CO-OR}_c$ ,  $-\text{O-CO-O-R}_c$ ,  $-\text{O-CO-R}_c$ ,  $-\text{NR}_c\text{-CO-R}_d$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{OH}$ ,  $\text{Ar}$ , heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O-CO-O-R}_f$ ,  $-\text{O-CO-R}_f$ ,  $-\text{OH}$ ,  $\text{Ar}$ , heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z\text{-CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ ,  $=\text{O}$ ,  $-\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{CO-OR}_c$ ,  $-\text{CO-R}_c$ ,  $-\text{NR}_c\text{-CO-NR}_d\text{R}_e$ ,  $-\text{C-CO-OR}_c$ ,  $-\text{NR}_c\text{-CO-R}_d$ ,  $-\text{O-CO-O-R}_c$ ,  $-\text{O-CO-NR}_d\text{R}_e$ ,  $-\text{SH}$ ,  $-\text{O-R}_b$ ,  $-\text{O-R}_a\text{-O-}$ ,  $-\text{S-R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are as defined above; or a pharmaceutically acceptable salt thereof, pharmaceutically acceptable prodrug thereof, pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.

18. A compound, salt, prodrug, or metabolite according to claim 17, wherein  $R^1$  is an alkyl unsubstituted or substituted with one or more substituents selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z\text{-CN}$  where  $z$  is 0-4, halogen,  $-\text{OH}$ ,  $-\text{O-R}_a\text{-O-}$ ,  $-\text{OR}_b$ ,  $-\text{CO-R}_c$ ,  $-\text{O-CO-R}_c$ ,  $-\text{CO-OR}_c$ ,  $-\text{O-CO-OR}_c$ ,  $-\text{O-CO-O-CO-R}_c$ ,  $-\text{O-OR}_c$ ,  $=\text{O}$ ,  $=\text{S}$ ,  $-\text{SO}_2\text{-R}_c$ ,  $-\text{SO-R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{O-CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c\text{-CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c\text{-CO-R}_e$ ,  $-\text{NR}_c\text{-CO-OR}_e$ ,  $-\text{CO-NR}_c\text{-CO-R}_d$ ,  $-\text{O-SO}_2\text{-R}_c$ ,  $-\text{O-SO-R}_c$ ,  $-\text{O-S-R}_c$ ,  $-\text{S-CO-R}_c$ ,  $-\text{SO-CO-OR}_c$ ,  $-\text{SO}_2\text{-CO-OR}_c$ ,  $-\text{O-SO}_3$ ,  $-\text{NR}_c\text{-SR}_d$ ,  $-\text{NR}_c\text{-SO-R}_d$ ,  $\text{NR}_c\text{-SO}_2\text{-R}_d$ , -

CO-SR<sub>c</sub>, CO-SO-R<sub>c</sub>, -CO-SO<sub>2</sub>-R<sub>c</sub>, -CS-R<sub>c</sub>, -CSO-R<sub>c</sub>, -CSO<sub>2</sub>-R<sub>c</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -O-CS-R<sub>c</sub>, -O-CSO-R<sub>c</sub>, O-CSO<sub>2</sub>-R<sub>c</sub>, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>c</sub>-CSO-R<sub>d</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, -COR<sub>f</sub>, -COOR<sub>f</sub>, -O-CO-O-R<sub>f</sub>, -O-CO-R<sub>f</sub>, -OH, Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where R<sub>d</sub> and R<sub>e</sub> can cyclize to form a heteroaryl or heterocycloalkyl group, and R<sub>f</sub> is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of NO<sub>2</sub>, -NH<sub>2</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, haloalkyl, haloaryl, -OH, =O, -N-OH, N-OR<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -CO-OR<sub>c</sub>, -CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -C-CO-OR<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -SH, -O-R<sub>b</sub>, -O-R<sub>a</sub>-O-, -S-R<sub>b</sub>, and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, and R<sub>e</sub> are as defined above.

19. A compound, salt, prodrug, or metabolite according to claim 18, wherein R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are each independently selected from hydrogen and halogen.

20. A compound, salt, prodrug, or metabolite according to claim 18, wherein said alkyl, when substituted, is alkyl-(heterocycloalkyl group) unsubstituted or substituted with one or more substituents independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, -NO<sub>2</sub>, -NH<sub>2</sub>, -N-OH, N-OR<sub>c</sub>, -CN, -(CH<sub>2</sub>)<sub>z</sub>-CN where z is 0-4, halogen, -OH, -O-R<sub>a</sub>-O-, -OR<sub>b</sub>, -CO-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-OR<sub>c</sub>, -O-CO-O-CO-R<sub>c</sub>, -O-OR<sub>c</sub>, =O, =S, -SO<sub>2</sub>-R<sub>c</sub>, -SO-R<sub>c</sub>, -NR<sub>d</sub>R<sub>e</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -O-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>c</sub>-CO-R<sub>e</sub>, -NR<sub>c</sub>-CO-OR<sub>e</sub>, -CO-NR<sub>c</sub>-CO-R<sub>d</sub>, -O-SO<sub>2</sub>-R<sub>c</sub>, -O-SO-R<sub>c</sub>, -O-S-R<sub>c</sub>, -S-CO-R<sub>c</sub>, -SO-CO-OR<sub>c</sub>, -SO<sub>2</sub>-CO-OR<sub>c</sub>, -O-SO<sub>3</sub>, -NR<sub>c</sub>-SR<sub>d</sub>, -NR<sub>c</sub>-SO-R<sub>d</sub>, NR<sub>c</sub>-SO<sub>2</sub>-R<sub>d</sub>, -CO-SR<sub>c</sub>, -CO-SO-R<sub>c</sub>, -CO-SO<sub>2</sub>-R<sub>c</sub>, -CS-R<sub>c</sub>, -CSO-R<sub>c</sub>, -CSO<sub>2</sub>-R<sub>c</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -O-CS-R<sub>c</sub>, -O-CSO-R<sub>c</sub>, -O-CSO<sub>2</sub>-R<sub>c</sub>, -SO<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -SO-NR<sub>d</sub>R<sub>e</sub>, -S-NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub>-CSO<sub>2</sub>-R<sub>d</sub>, -NR<sub>c</sub>-CSO-R<sub>d</sub>, -NR<sub>c</sub>-CS-R<sub>d</sub>, -SH, -S-R<sub>b</sub>, and -PO<sub>2</sub>-OR<sub>c</sub>, where R<sub>a</sub> is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl, R<sub>b</sub> is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen, -CO-R<sub>c</sub>, -CO-OR<sub>c</sub>, -O-CO-O-R<sub>c</sub>, -O-CO-R<sub>c</sub>, -NR<sub>c</sub>-CO-R<sub>d</sub>, -CO-NR<sub>d</sub>R<sub>e</sub>, -OH, Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and R<sub>c</sub>, R<sub>d</sub> and R<sub>e</sub> are each independently

selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O-CO-O-R}_f$ ,  $-\text{O-CO-R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $\text{R}_d$  and  $\text{R}_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $\text{R}_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ , keto ( $=\text{O}$ ),  $-\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{CO-OR}_c$ ,  $-\text{CO-R}_c$ ,  $-\text{NR}_c-\text{CO-NR}_d\text{R}_e$ ,  $-\text{C-CO-OR}_c$ ,  $-\text{NR}_c-\text{CO-R}_d$ ,  $-\text{O-CO-O-R}_c$ ,  $\text{O-CO-NR}_d\text{R}_e$ ,  $-\text{SH}$ ,  $-\text{O-R}_b$ ,  $-\text{O-R}_a-\text{O}-$ ,  $-\text{S-R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $\text{R}_a$ ,  $\text{R}_b$ ,  $\text{R}_c$ ,  $\text{R}_d$ , and  $\text{R}_e$  are as defined above, alkyl-(aryl group) unsubstituted or substituted with one or more substituents independently selected from the group consisting of hydrogen, alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $-\text{CO-OR}_c$ ,  $-\text{O-CO-O-CO-R}_c$ ,  $-\text{O-OR}_c$ ,  $=\text{O}$ ,  $=\text{S}$ ,  $-\text{SO}_2-\text{R}_c$ ,  $-\text{SO-R}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{O-CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO-NR}_d\text{R}_e$ ,  $-\text{NR}_c-\text{CO-R}_c$ ,  $-\text{NR}_c-\text{CO-OR}_c$ ,  $-\text{CO-NR}_c-\text{CO-R}_d$ ,  $-\text{O-SO}_2-\text{R}_c$ ,  $-\text{O-SO-R}_c$ ,  $-\text{O-S-R}_c$ ,  $-\text{S-CO-R}_c$ ,  $-\text{SO-CO-OR}_c$ ,  $-\text{SO}_2-\text{CO-OR}_c$ ,  $-\text{O-SO}_3$ ,  $-\text{NR}_c-\text{SR}_d$ ,  $-\text{NR}_c-\text{SO-R}_d$ ,  $\text{NR}_c-\text{SO}_2-\text{R}_d$ ,  $-\text{CO-SR}_c$ ,  $-\text{CO-SO-R}_c$ ,  $-\text{CO-SO}_2-\text{R}_c$ ,  $-\text{CS-R}_c$ ,  $-\text{CSO-R}_c$ ,  $-\text{CSO}_2-\text{R}_c$ ,  $-\text{NR}_c-\text{CS-R}_d$ ,  $-\text{O-CS-R}_c$ ,  $-\text{O-CSO-R}_c$ ,  $-\text{O-CSO}_2-\text{R}_c$ ,  $-\text{SO}_2-\text{NR}_d\text{R}_e$ ,  $-\text{SO-NR}_d\text{R}_e$ ,  $-\text{S-NR}_d\text{R}_e$ ,  $-\text{NR}_d-\text{CSO}_2-\text{R}_d$ ,  $-\text{NR}_c-\text{CSO-R}_d$ ,  $-\text{NR}_c-\text{CS-R}_d$ ,  $-\text{SH}$ ,  $-\text{S-R}_b$ , and  $-\text{PO}_2-\text{OR}_c$ , where  $\text{R}_a$  is independently selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $\text{R}_b$  is independently selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-\text{CO-R}_c$ ,  $-\text{CO-OR}_c$ ,  $-\text{O-CO-O-R}_c$ ,  $-\text{O-CO-R}_c$ ,  $-\text{NR}_c-\text{CO-R}_d$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{OH}$ , Ar, heteroaryl, heterocycloalkyl, and cycloalkyl, and  $\text{R}_c$ ,  $\text{R}_d$  and  $\text{R}_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-\text{COR}_f$ ,  $-\text{COOR}_f$ ,  $-\text{O-CO-O-R}_f$ ,  $-\text{O-CO-R}_f$ ,  $-\text{OH}$ , Ar, heteroaryl, cycloalkyl, and heterocycloalkyl, where  $\text{R}_d$  and  $\text{R}_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $\text{R}_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{CN}$ ,  $-(\text{CH}_2)_z-\text{CN}$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-\text{OH}$ ,  $=\text{O}$ ,  $-\text{N-OH}$ ,  $\text{N-OR}_c$ ,  $-\text{NR}_d\text{R}_e$ ,  $-\text{CO-NR}_d\text{R}_e$ ,  $-\text{CO-OR}_c$ ,  $-\text{CO-R}_c$ ,  $-\text{NR}_c-\text{CO-NR}_d\text{R}_e$ ,  $-\text{C-CO-OR}_c$ ,  $-\text{NR}_c-\text{CO-R}_d$ ,  $-\text{O-CO-O-R}_c$ ,  $\text{O-CO-NR}_d\text{R}_e$ ,  $-\text{SH}$ ,  $-\text{O-R}_b$ ,  $-\text{O-R}_a-\text{O}-$ ,  $-\text{S-R}_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl,

unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are as defined above.

21. A compound, salt, prodrug, or metabolite according to claim 21, wherein  $R^4$ ,  $R^5$  and  $R^6$  are each independently selected from hydrogen and halogen.

22. A compound, salt, prodrug, or metabolite according to claim 17, wherein  $R^1$  is selected from the group consisting of aryl, heteroaryl, and heterocycloalkyl, unsubstituted or substituted with one or more substituents selected from the group consisting of alkyl, heteroalkyl, haloalkyl, haloaryl, halocycloalkyl, haloheterocycloalkyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl,  $-NO_2$ ,  $-NH_2$ ,  $-N-OH$ ,  $N-OR_c$ ,  $-CN$ ,  $-(CH_2)_z-CN$  where  $z$  is 0-4, halogen,  $-OH$ ,  $-OR_a-O-$ ,  $-OR_b$ ,  $-CO-R_c$ ,  $-O-CO-R_c$ ,  $-CO-OR_c$ ,  $-O-CO-OR_c$ ,  $-O-CO-O-CO-R_c$ ,  $-O-OR_c$ ,  $=O$ ,  $=S$ ,  $-SO_2-R_c$ ,  $-SO-R_c$ ,  $-NR_dR_e$ ,  $-CO-NR_dR_e$ ,  $-O-CO-NR_dR_e$ ,  $-NR_c-CO-NR_dR_e$ ,  $-NR_c-CO-R_e$ ,  $-NR_c-CO-OR_c$ ,  $-CO-NR_c-CO-R_d$ ,  $-O-SO_2-R_c$ ,  $-O-SO-R_c$ ,  $-O-S-R_c$ ,  $-S-CO-R_c$ ,  $-SO-CO-OR_c$ ,  $-SO_2-CO-OR_c$ ,  $-O-SO_3$ ,  $-NR_c-SR_d$ ,  $-NR_c-SO-R_d$ ,  $NR_c-SO_2-R_d$ ,  $-CO-SR_c$ ,  $-CO-SO-R_c$ ,  $-CO-SO_2-R_c$ ,  $-CS-R_c$ ,  $-CSO-R_c$ ,  $-CSO_2-R_c$ ,  $-NR_c-CS-R_d$ ,  $-O-CS-R_c$ ,  $-O-CSO-R_c$ ,  $-O-CSO_2-R_c$ ,  $-SO_2-NR_dR_e$ ,  $-SO-NR_dR_e$ ,  $-S-NR_dR_e$ ,  $-NR_d-CSO_2-R_d$ ,  $-NR_c-CSO-R_d$ ,  $-NR_c-CS-R_d$ ,  $-SH$ ,  $-S-R_b$ , and  $-PO_2-OR_c$ , where  $R_a$  is selected from the group consisting of alkyl, heteroalkyl, alkenyl, and alkynyl,  $R_b$  is selected from the group consisting of alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl, halogen,  $-CO-R_c$ ,  $-CO-OR_c$ ,  $-O-CO-O-R_c$ ,  $-O-CO-R_c$ ,  $-NR_c-CO-R_d$ ,  $-CO-NR_dR_e$ ,  $-OH$ ,  $Ar$ , heteroaryl, heterocycloalkyl, and cycloalkyl, and  $R_c$ ,  $R_d$  and  $R_e$  are each independently selected from the group consisting of hydrogen, halogen, alkyl, heteroalkyl, haloalkyl, alkenyl, alkynyl,  $-COR_f$ ,  $-COOR_f$ ,  $-O-CO-O-R_f$ ,  $-O-CO-R_f$ ,  $-OH$ ,  $Ar$ , heteroaryl, cycloalkyl, and heterocycloalkyl, where  $R_d$  and  $R_e$  can cyclize to form a heteroaryl or heterocycloalkyl group, and  $R_f$  is selected from the group consisting of hydrogen, alkyl, and heteroalkyl, and where any of the alkyl, heteroalkyl, alkylene, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl moieties present in the above substituents may be further substituted with one or more substituents independently selected from the group consisting of  $NO_2$ ,  $-NH_2$ ,  $-CN$ ,  $-(CH_2)_z-CN$  where  $z$  is 0-4, halogen, haloalkyl, haloaryl,  $-OH$ ,  $=O$ ,  $-N-OH$ ,  $N-OR_c$ ,  $-NR_dR_e$ ,  $-CO-NR_dR_e$ ,  $-CO-OR_c$ ,  $-CO-R_c$ ,  $-NR_c-CO-NR_dR_e$ ,  $-C-CO-OR_c$ ,  $-NR_c-CO-R_d$ ,  $-O-CO-O-R_c$ ,  $O-CO-NR_dR_e$ ,  $-SH$ ,  $-O-R_b$ ,  $-O-R_a-O-$ ,  $-S-R_b$ , and unsubstituted alkyl, unsubstituted aryl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, and unsubstituted heteroaryl, where  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  are as defined above.

23. A compound, salt, prodrug, or metabolite according to claim 20, wherein  $R^4$ ,  $R^5$  and  $R^6$  are each independently selected from hydrogen and halogen.

24. A pharmaceutical composition comprising an amount of an agent effective to modulate cellular proliferation and a pharmaceutically acceptable carrier, said agent being

selected from the group consisting of compounds, pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, pharmaceutically metabolites as defined in claim 1.

25. A pharmaceutical composition comprising an amount of an agent effective to inhibit a protein kinase and a pharmaceutically acceptable carrier, said agent being selected from the group consisting of compounds, pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, or pharmaceutically metabolites as defined in claim 1.

26. A pharmaceutical composition according to claim 22, wherein said protein kinase is a CDK1, CDK1/cyclin complex, CDK2, CDK2/cyclin complex, CDK4, CDK4/cyclin complex, CDK6, CDK6/cyclin complex, FGF, or LCK.

27. A pharmaceutical composition comprising an amount of an agent effective to modulate cellular proliferation and a pharmaceutically acceptable carrier, said agent being selected from the group consisting of compounds, pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, or pharmaceutically metabolites as defined in claim 17.

28. A pharmaceutical composition comprising an amount of an agent effective to inhibit a protein kinase and a pharmaceutically acceptable carrier, said agent being selected from the group consisting of a compound, pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, or pharmaceutically metabolites as defined in claim 17.

29. A pharmaceutical composition according to claim 25, wherein said protein kinase is a CDK1, CDK1/cyclin complex, CDK2, CDK2/cyclin complex, CDK4, CDK4/cyclin complex, CDK6, CDK6/cyclin complex, VEGF, FGF, or LCK.






30. A method of treating a disease condition or disorder associated with uncontrolled cellular proliferation, comprising administering to a subject in need thereof a therapeutically effective amount of a compound, pharmaceutically acceptable salt, pharmaceutically acceptable prodrug or a pharmaceutically metabolite as defined in claim 1.


31. A method according to claim 27, wherein the disease condition or disorder is tumor growth, angiogenesis, viral infection, autoimmune disease or neurodegenerative disorder.


32. A method of modulating or inhibiting the activity of a protein kinase receptor, comprising delivering to the kinase receptor an effective amount of a compound, pharmaceutically acceptable salt, pharmaceutically acceptable prodrugs or pharmaceutically metabolites as defined in claim 1.



33. A method according to claim 29, wherein the protein kinase is a CDK complex, VEGF, FGF, or LCK.
34. A method of treating a disease condition or disorder associated with uncontrolled cellular proliferation, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound, pharmaceutically acceptable salt, pharmaceutically acceptable prodrug or pharmaceutically metabolite as defined in claim 17.
35. A method according to claim 31, wherein the disease condition or disorder is tumor growth, angiogenesis, viral infection, autoimmune disease or neurodegenerative disorder.
36. A method of modulating or inhibiting the activity of a protein kinase receptor, comprising delivering to the kinase receptor an effective amount of a compound, pharmaceutically acceptable salt, pharmaceutically acceptable prodrug or pharmaceutically metabolite as defined in claim 17.
37. A method according to claim 33, wherein the protein kinase is a CDK complex, VEGF, FGF or LCK.
38. A compound selected from the group consisting of:







or a pharmaceutically acceptable salt of said compound.

39. A compound selected from the group consisting of examples A1 through A62, B1 through B5, C, D1, D2, E, F, and G1 through G396, or a pharmaceutical acceptable salt of said compound.

**THIS PAGE BLANK (USPTO)**